arXiv:math/0312484v2 [math.CO] 18 Jan 2004

Uniform chains of small transfinite rank

F.P.Weber
R.M.Vollmer

School of Earth Sciences
University of Leeds
LS2 9JT
United Kingdom

Abstract

We extend F.Pastjin's construction of uniform decomposable chains of finite rank to those of rank 'finite over a limit' and investigate infinite (length ω) products and unions of such chains. We derive an extension of Pastjin's characterisation to uniform decomposable chains of small transfinite rank ($\leq \omega + \omega$). We conclude by indicating how the resulting descriptions can be iterated to chains of rank $\leq \omega^\omega$.

Contents

[1] Pastjin's Chains
[2] Tails of first limit rank
[3] Chains of rank finite over a limit
[4] Products of uniform chains
[5] Chains of first limit rank
[6] Chains of next limit rank
[7] Short representations
[8] Short representations of tails
[9] Extensions

[1] Pastjin's Chains

[1.1] Decomposable chains

Let A be a chain (i.e., a linear, or total, order). Define an equivalence relation $\rho = \rho^A$ by: $x\rho y$ if the interval between x and y in A is finite. Let $[x] = [x]^A = \{y \in A : x\rho y\}$ (the component of x in A) and $A/\rho = \{[x] : x \in A\}$.

Iterate the definition through all ordinals: Set $\rho_1 = \rho$; suppose ρ_α defined, set $A/\rho_\alpha = \{[x]_\alpha : x \in A\}$ and $x\rho_{\alpha+1}y$ in A if $[x]_\alpha \rho [y]_\alpha$ in A/ρ_α (where $[x]_\alpha = [x]^A_\alpha = \{y \in A : x\rho_\alpha y\}$ is the *α-component* of x (in A)); suppose ρ_β defined for all $\beta < \alpha$, for limit α, set $A/\rho_\beta = \{[x]_\beta : x \in A\}$ and set $x\rho_\alpha y$ in A if for some $\beta < \alpha$ it is the case that $[x]_\beta \rho [y]_\beta$ in A/ρ_β. In the following, write A/α for A/ρ_α. and $[x]_\alpha$ for $[x]_\alpha$. The *rank* of A is the least ordinal α such that $A/\alpha \cong 1$ (the singleton chain) and is denoted $\mathrm{rk}(A)$. A chain is *decomposable* if $\mathrm{rk}(A)$ is defined. Subsequently, all chains are assumed decomposable.

(1.2) **Definition** (Uniform chains) Let $A^{\geq x} \cong \{y \in A : x \leq y\}$. A chain A is (right) *uniform* if for all $x, y \in A$ the chains $A^{\geq x}$ and $A^{\geq y}$ are order isomorphic.

In this case denote by $\mathrm{f}A$ (a representative sample of) the tail of A. (We shall also use $\mathrm{f}_x A$ to denote $A^{\geq x}$ and use f and f_x interchangably with uniform chains.) f is an idempotent operation, $\mathrm{f}^2 = \mathrm{f}$. Let $\mathcal{U}(\alpha)$ denote the set of uniform chains of rank α and $\mathcal{U}_0(\alpha)$ the set of tails of chains in $\mathcal{U}(\alpha)$. $\mathcal{U}(<\alpha) = \bigcup_{i<\alpha} \mathcal{U}(i)$. If not specified otherwise, letters A and B refer in the following to uniform (decomposable) chains.

[1.2] Pastjin's uniform chain generator

Let N be the chain of non-negative integers and Z the chain of all integers. There is a scheme to generate all uniform chains of finite rank as chains of form $A\mathrm{Z}$ or $B + A\mathrm{N}$ from uniform chains of smaller finite rank, starting with the singleton chain. (AB is the anti-lexicographically ordered cartesian product of chains A and B, effectively $AB = \sum_{b \in B} A$.)

(1.3) **Definition** (Sub-diagonal strings) For an ordinal α, let $\mathcal{S}(\alpha) = \{s \in {}^\alpha\alpha : s(i) \leq i \text{ for all } i < \alpha\}$. Also let $\mathcal{S}(<\alpha) = \bigcup_{i<\alpha} \mathcal{S}(i)$ and, e.g., (for ordinal β) $\mathcal{S}([\beta, \alpha)) = \bigcup_{\beta \leq i < \alpha} \mathcal{S}(i)$, etc. (Note that $\mathcal{S}(1) = \{\langle 0 \rangle\}$.)

(1.4) **Definition** (Pastjin's finite rank unifom chain constructor) Set $A(\langle 0 \rangle) = 1 = \langle 0 \rangle$, and for any $s \in \mathcal{S}$

$$A(s \wedge \langle a, b \rangle) = \begin{cases} A(s \wedge \langle a \rangle)\mathrm{Z} & \text{if } b = |s \wedge \langle a \rangle| \\ A(s \wedge \langle b \rangle) + A(s \wedge \langle a \rangle)\mathrm{N} & \text{else, i.e., if } b < |s \wedge \langle a \rangle| \end{cases}$$

(We use '$\wedge\langle\rangle$' to denote concatenation of finite strings, while retailing the conventional '+' for chains.) Letting $A(\langle\ \rangle) = \langle\ \rangle = \mathbf{0}$ (:= the empty chain), by finite induction,

$$A(s) = A(s\lceil k(s))\mathrm{Z} + \sum_{k(s)<i<n(s)} A(s\lceil i)\mathrm{N}$$

where $n(s) = |s| - 1$ and $k(s) = \max\{i \leq n : s(n(s)) = i\}$.

Here is a basic list:

$$\begin{array}{lll}
A(\langle 0\rangle) \cong 1 & & \\
A(\langle 0,0\rangle) \cong 1+1\mathrm{N} \cong \mathrm{N} & A(\langle 0,1\rangle) \cong 1\mathrm{Z} \cong \mathrm{Z} & \\
A(\langle 0,0,0\rangle) \cong 1+1\mathrm{N}+\mathrm{NN} \cong \mathrm{N}^2 & A(\langle 0,0,1\rangle) \cong \mathrm{Z}+\mathrm{NN} \cong \mathrm{Z}+\mathrm{N}^2 & A(\langle 0,0,2\rangle) \cong N\mathrm{Z} \\
A(\langle 0,1,0\rangle) \cong \mathrm{N}+\mathrm{ZN} & A(\langle 0,1,1\rangle) \cong \mathrm{ZN} & A(\langle 0,1,2\rangle) \cong \mathrm{Z}^2
\end{array}$$

(1.5) **Theorem** (Pastjin) The chains $A(s), s \in \mathcal{S}(n+1)$ exhaust all uniform chains of rank n (up to isomorphism).

(1.6) **Notation** Define operation $s \to s^+$ and $s \to s^-$ on binary strings s by:

$$(s \wedge \langle a\rangle)^- = s, \qquad (s \wedge \langle a,b\rangle)^+ = s \wedge \langle b\rangle.$$

Thus, where $n = |s|-1$,

$$A(s) = \begin{cases} A(s^-)\mathrm{Z} & \text{if } s(n)=n \\ A(s^+)+A(s^-)\mathrm{N} & \text{if } s(n)<n. \end{cases} \tag{1}$$

Define iterations of $()^+$ and $()^-$ by $s^{+(k+1)} = (s^{+k})^+$ and $s^{-(k+1)} = (s^{-k})^-$. Note that $s^{+k} \in \mathcal{S}$ just if $s(n) \le n-k$ (where $n=|s|-1$), since $n-k = |s^{+k}|-1$, and $s^{+k}(n-k) = s(n)$. In particular, $s^{+n} \in \mathcal{S}$ just if $s(n)=0$, where $n=|s|-1$. For convenience, let $s^{=} = s^{--} = (s^-)^-$, and $s^{\equiv} = s^{---} = (s^{=})^-$. (However, $s^{\pm}$ continues to read "s^- or s^+".) Note that $s^{+-} = s^{--} = s^{=}$.

[1.3] Uniform tails

(1.7) **Lemma** (Finite tail lemma) $\mathrm{f}A(s) \cong A(s^- \wedge \langle 0\rangle) = \sum_{i<|s|} A(s\lceil i)\mathrm{N}$.

Proof The second equation is immediate. For the first, use induction. The base cases are trivial, e.g.,

$$\mathrm{f}A(\langle 0,1\rangle) = \mathrm{fZ} \cong \mathrm{N} = A(\langle 0,0\rangle) = A(\langle 0\rangle \wedge \langle 0\rangle)$$

In the induction step use the following observation:

$$\mathrm{f}_{(a,b)}AB = \mathrm{f}_a A + A\mathrm{f}_b B. \tag{2}$$

If $s(n)=n$ for $n=|s|-1$,

$$\mathrm{f}A(s) = \mathrm{f}A(s^-) + A(s^-)\mathrm{fZ} \cong \mathrm{f}A(s^-) + A(s^-)\mathrm{N} = A(s^{=} \wedge \langle 0\rangle) + A(s^-)\mathrm{N} = A(s^- \wedge \langle 0\rangle),$$

from the induction hypothesis and the fact that $(s^- \wedge \langle 0\rangle)^+ = s^{=} \wedge \langle 0\rangle$.

If $s(n)<n$ distinguish the case where the cut-off is in $A(s^+)$ or in $A(s^-)\mathrm{N}$. In the respective case we get tails of form

$$\mathrm{f}A(s) = \begin{cases} \mathrm{f}A(s^+) + A(s^-)\mathrm{N} = A(s^{=} \wedge \langle 0\rangle) + A(s^-)\mathrm{N}, & \text{(cut-off in } A(s^+)) \\ \mathrm{f}A(s^-) + A(s^-)\mathrm{fN} = A(s^{=} \wedge \langle 0\rangle) + A(s^-)\mathrm{N}, & \text{(cut-off in some } A(s^-) \times \{n\}), \end{cases}$$

again from the induction hypothesis and the fact that $(s^{\pm})^- \wedge \langle 0\rangle = s^{=} \wedge \langle 0\rangle)$. As before, either case reduces to $A(s^- \wedge \langle 0\rangle)$. •

This suggests extending f to an operation on $\mathcal{S}(\alpha)$ for successor ordinals α by $\mathrm{f}s = s^- \wedge \langle 0 \rangle$. Thus, $\mathrm{f}A(s) = A(\mathrm{f}s)$ for $s \in \mathcal{S}(n+1)$, $n \in \omega$. As a direct corollary:

(1.8) **Lemma** (String lemma)
(i) $(\mathrm{f}s)^+ = (s^- \wedge \langle 0 \rangle)^+ = s^= \wedge \langle 0 \rangle = \mathrm{f}s^- = \mathrm{f}s^+$;
(ii) $(\mathrm{f}s)^- = (s^- \wedge \langle 0 \rangle)^- = s^-$. •

(1.9) **Remark** How do we identify elements in chains $A(s)$? For instance, in the second case of the decomposition of $A(s)$ in (1), we could represent elements of $A(s)$ in the form:

$$x = \begin{cases} \langle a, 0 \rangle & \text{for } a \text{ in } A(s^+) \\ \langle b, n+1 \rangle & \text{for } b \text{ in } A(s^-), \end{cases}$$

and we frequently do. Similarly, we use '$B \times \{n\}$' to refer to the $n+1$st hyper-component of $A + B\mathrm{N}$, where $\mathrm{rk}(A) = \mathrm{rk}(B)$. On the other hand, since, e.g., a hyper-component of $A(s)$ is isomorphic to $A(t)$ for $t = s^{\pm}$, we say e.g., of $x \in A(s) = A(s^+) + A(s^-)\mathrm{N}$ that it is 'in $A(s^+)$' to indicate that it is in the initial $|s| - 1$ component of $A(s)$, or that it is 'in some $A(s^-)$' to indicate that it is in one of the successor $|s| - 1$ components of $A(s)$, and that it is 'in $A(s^-) \times \{n\}$' to indicate that it is in the nth successor $|s|$ component of $A(s)$. I.e., when considering embeddings between chains, we assume them to be 'canonical', as defined below.

(1.10) **Lemma** (Sub-component lemma) For $x \in A \in \mathcal{U}$, each $[x]_i^A$, for $i \leq \mathrm{rk}(A)$ is of the form $A(s_i)$, for some $s_i \in \mathcal{S}(i+1)$ such that
(a) $s_i \in \{s_{i+1}^+, s_{i+1}^-\}$, hence
(b) $s_i^- = s_{i+1}^=$, and
(c) $s_i^- \subset s$ for $s \in \mathcal{S}(< \omega)$ such that $A = A(s)$

Proof (a) and (b) are clear from the corollary to Pastjin's definition and the preceding remark. Setting $n = \mathrm{rk}(A) = \mathrm{rk}(A(s)) = |s| - 1 \in \omega$, we get $s_n = s$, and for all $0 < i \leq n$, $s_{i-1} \in \{s_i^+, s_i^-\}$, by (a), so $s_{i-1}^- \in \{s_i^{+-}, s_i^{--}\} = \{s_i^=\}$, and so by finite ('downward') induction: $s_{i-1}^- \subset s_i^- \subset s_n^- = s^-$. •

(1.11) **Definition** (Canonical embeddings) Let $s \in \mathcal{S}(n+2), t \in \mathcal{S}(n+1)$ such that $t \in \{s^+, s^-\}$. Call an embedding of $A(t)$ into $A(s)$ *canonical*, if
(a) $A(t)$ is mapped onto the initial n component of $A(s)$ of form $A(s^+)$ if $t = s^+$, and
(b) $A(t)$ is mapped onto one of the successor n components of $A(s)$ of form $A(s^-)$, if $t = s^-$.

[2] Tails of first limit rank

The modest goal here is a characterisation of the tails (of uniform chains) of rank ω.

[2.1] Tail-types and successor-types of hyper-components

We start with a basic observation: Let $A \in \mathcal{U}$, $x \in A$ and let $C_n = [x]^A_n$, $n \in \mathrm{rk}(A)$. Then

$$[x]^{\mathrm{f}_x A}_{n+1} = [x]^{\mathrm{f}_x A}_n + A_n \mathrm{N} = \mathrm{f}_x C_n + A_n \mathrm{N} \tag{2.1}$$

where A_n is (a representative of the isomorphism class of) the successor n-component (to C_n) in C_{n+1}, and $\mathrm{f}A_n = \mathrm{f}C_n$.

(2.1) **Definition** (nth α hyper-component) If $\mathrm{rk}(A) = \alpha + 1$, and $A/\alpha \cong \mathrm{N}$ let

$$H^\alpha_n(A) := \text{the } n\text{th } \alpha \text{ hyper-component of } A.$$

If $\mathrm{rk}(A) = \alpha + 1$, and $A/\alpha \cong \mathrm{N}$ or $A/\alpha \cong \mathrm{Z}$ let

$$H^\alpha_n(A, x) := \text{the } n\text{th } \alpha \text{ hyper-component of } A \text{ relative to } [x]^A_\alpha$$

(i.e., such that that $H^\alpha_0(A, x) = [x]^A_\alpha$). Where x is fixed in a context, write $H^\alpha_n(A)$ again for $H^\alpha_n(A, x)$. If $A/\alpha \cong \mathrm{N}$, x can always be arranged to lie in the zeroth hyper-component. In particular,

$$H^\alpha_n([x]^A_{\alpha+1}) := H^\alpha_n([x]^A_{\alpha+1}, x).$$

(2.2) **Lemma** (Hyper-component isomorphism) For $x, y \in A \in \mathcal{U}$, $\alpha < \mathrm{rk}(A)$,

$$(\forall n, m)[\mathrm{f}H^\alpha_n([x]^A_{\alpha+1}) \cong \mathrm{f}H^\alpha_m([y]^A_{\alpha+1})], \qquad (\forall n, m > 0)[H^\alpha_n([x]^A_{\alpha+1}) \cong H^\alpha_m([y]^A_{\alpha+1})]$$

Proof Fix x in A. In $\mathrm{f}_x([x]^A_{\alpha+1})$,

$$\mathrm{f}_x([x]^A_\alpha) = \mathrm{f}H^\alpha_0([x]^A_{\alpha+1}) \text{ is the zeroth } \alpha \text{ component,}$$
$$(\text{for } n > 0)\ H^\alpha_n([x]^A_{\alpha+1}) \text{ is the } n\text{th } \alpha \text{ component.}$$

For any y in A, since $\mathrm{f}_x A \cong \mathrm{f}_y A$ and since isomorphisms preserve components and component ordering,

$$\begin{aligned} \mathrm{f}H^\alpha_0([x]^A_{\alpha+1}) &\cong \mathrm{f}H^\alpha_0([y]^A_{\alpha+1}) \\ H^\alpha_n([x]^A_{\alpha+1}) &\cong H^\alpha_n([y]^A_{\alpha+1}) \end{aligned} \tag{a}$$

wherever either is defined.

Fix n such that $H_n^\alpha([x]^A_{\alpha+1})$ is defined and let $z \in H_n^\alpha([x]^A_{\alpha+1}) \subseteq [x]^A_{\alpha+1}$. In $\mathrm{f}_z([x]^A_{\alpha+1})$,

$$\begin{aligned} \mathrm{f}_z([z]^A_\alpha) = \mathrm{f}H_0^\alpha([x]^{,}_{\alpha+1}z) = \mathrm{f}H_n^\alpha([x]^{)}_{\alpha+1} &\text{ is the zeroth } \alpha \text{ component,} \\ (\text{for } k > 0)\ H_k^\alpha([x]^A_{\alpha+1}, z) = H^\alpha_{n+k}([x]^A_{\alpha+1}) &\text{ is the } m\text{th zeroth } \alpha \text{ component.} \end{aligned}$$

Since $[x]^A_{\alpha+1}$ is uniform, $\mathrm{f}_x([x]^A_{\alpha+1}) \cong \mathrm{f}_z([x]^A_{\alpha+1})$, and since isomorphisms preserve components and component ordering,

$$\begin{aligned} &\mathrm{f}H_0^\alpha([x]^{)}_{\alpha+1} \cong \mathrm{f}H_n^\alpha([x]^{)}_{\alpha+1} \\ &(\text{for } n, k > 0)\ H_n^\alpha([x]^A_{\alpha+1}) \cong H^\alpha_{n+k}([x]^A_{\alpha+1}). \end{aligned} \tag{b}$$

Now, (a) allows us to 'vary' x in the expressions $\mathrm{f}H_n^\alpha([x]^A_{\alpha+1})$ and $H_n^\alpha([x]^A_{\alpha+1})$ for fixed n and $n > 0$, respectively, while (b) allows us to 'vary' n and $n > 0$ in them, respectively, and always produce isomorphic components, respectively. •

We say a component is a *successor component* if it has an immediate predecessor in the ordering taken modulo its rank, i.e., if the component is of form $C = [x]^A_\alpha$ but C/α has an immediate predecessor in $[x]^A_{\alpha+1}/\alpha$ – hence in any super-component of the latter in A/α. (For illustration, a component is a *gap component* if it does not have an immediate predecessor. We also then say that *there is a gap at* C. If a gap-component is of form $C = [x]^A_\alpha$, the *depth* of the gap *relative to* C is $\delta = \beta - \alpha$ where β is minimal such that either $B = [x]^A_\beta$ has an immediate successor in A/β (hence in $[x]^A_\beta$), or that $\beta = \mathrm{rk}(A)$. (Here $\beta - \alpha = \sup\{\gamma : \alpha + \gamma \leq \beta\}$.) For all $\gamma \in (\alpha, \beta)$, C is the also *the initial* α component in $[x]^A_\gamma$. The *absolute depth* of the gap determined by that component is $\beta - \gamma$ where γ is minimal $\leq \alpha$ such that for all $\delta \in [\gamma, \alpha]$, there is a y such that $D = [y]^C_\delta$ is minimal in C/δ – and hence has no immediate predecessor, hence is a gap component. Note that for such y, D is a minimal hyper-component of $[y]^A_{\delta+1}$, as well a minimal element in $[y]^A_\gamma = [y]^B_\gamma$, for any $\gamma \in [\delta.\beta)$. Isomorphisms preserve not only immediate successor-ship but as gaps and gap-depth as well.) The preceding justifies:

(2.3) **Definition** (Tail and successor-component)

Let T^A_α denote the common isomorphism type of the tails of components of rank α in A, i.e., those representable as $\mathrm{f}[x]^A_\alpha$ for some $x \in A$ – and hence as $\mathrm{f}H_n^\alpha([x]^A_{\alpha+1})$ for some $x \in A$ and n.

Let S^A_α denote the common isomorphism type of the successor components of rank α in A, i.e., those representable as $H_n^\alpha([x]^A_{\alpha+1})$ for some $x \in A$ and $n > 0$.

Thus,

(2.4) **Lemma** (Tail lemma) For $x \in A \in \mathcal{U}$,

$$\mathrm{f}_x A \cong \sum_{n \in \mathrm{rk}(A)} A_n \mathrm{N}, \qquad \text{where} \qquad A_n \cong S^A_{n+1} \cong H_1^n([x]^A_{n+1}).$$

Proof Set $C_n = [x]^A_n$. The statement is equivalent to $\mathrm{f}_x C_n = \sum_{i \in n} A_i \mathrm{N}$, for $n = \mathrm{rk}(A)$. This follows by induction over $n < \mathrm{rk}(A)$.

For $n = 1$, $C_1 \in \{\mathrm{N}, \mathrm{Z}\}$ and $A_0 \cong \mathbf{1}$, so $\mathrm{f}C_1 \cong \mathrm{N} \cong A_0\mathrm{N}$.

Assume the lemma true for all $i < \alpha \leq \mathrm{rk}(A)$.

First let $\alpha = n+1$. Then by (2.1) $\mathrm{f}_x C_\alpha = \mathrm{f}_x C_n + A_n\mathrm{N}$, but by inductive hypothesis, $\mathrm{f}_x C_n = \sum_{i\in n} A_i\mathrm{N}$, giving $\mathrm{f}_x C_\alpha = \sum_{i\in n} A_i\mathrm{N} + A_n\mathrm{N}$, hence the required result.

Finally let α be a limit ordinal. Then for each $n < \alpha$ $\mathrm{f}_x C_n = \sum_{n\in n} A_n\mathrm{N}$, and for every pair $n < k < \alpha$, $\mathrm{f}_x C_n$ embeds as initial segment into $\mathrm{f}_x C_k$, giving again the required result. •

[2.2] Tails of rank ω

The preceding allows to classify all $A \in \mathcal{U}_0(\omega)$.

(2.5) **Definition** For $\sigma \in \mathcal{S}(\omega)$ set $A(\sigma \wedge \langle 0\rangle) = \sum_{i\in\omega} A(\sigma\lceil i+1)\mathrm{N}$

With $A(\langle\rangle) = \mathbf{0}$ (:= the empty chain), the above is tanatamount to setting $A(s \wedge \langle 0\rangle) = \sum_{i\in\omega} A(s\lceil i)\mathrm{N}$.

(2.6) **Lemma** (Rank ω tail representation lemma) For every $A \in \mathcal{U}(\omega)$ there exists $\sigma \in \mathcal{S}(\omega)$ such that $\mathrm{f}A = A(\sigma \wedge \langle 0\rangle)$.

Proof Choose any $x \in A$ and let $B = \mathrm{f}_x A$. Fix $s_n \in \mathcal{S}(n+1)$ such that $A(s_n) = [x]^B_n$. Since $[x]^B_n = \mathrm{f}_x[x]^B_n$, in fact $s_n = s^-_n \wedge \langle 0\rangle$.

Define a sequence $x_n \in A$ such that $x_0 = x$ and $[x_{n+1}]^B_{n+1}$ is the successor $n+1$ component to $[x]^B_{n+1}$ in A/ρ_{n+1}. Note that $[x_{n+1}]^B_{n+1} = [x_{n+1}]^A_{n+1}$. By uniformity of A (and hence B), $[x]^B_{n+1} = [x]^B_n + [x_n]^B_n\mathrm{N} = [x]^B_n + [x_n]^A_n\mathrm{N}$, so $A(s_{n+1}) = A(s_n) + A(t_n)\mathrm{N}$, where $s_n = s^+_{n+1}$ and $t_n \in \mathcal{S}(n+1)$ such that $t_n = s^-_{n+1} = s_{n+1}\lceil n+1$. and $s_n\lceil n = s^-_n = t^-_n = t_n\lceil n$. Setting $\sigma = \cup_{n\in\mathrm{N}} s_n\lceil n \in \mathcal{S}(\omega)$, we get $t_n = s_{n+1}\lceil n+1 = \sigma\lceil n+1$, and thus

$$\mathrm{f}_x A = B = \sum_{n\in\mathrm{N}} [x_n]^B_n\mathrm{N} = \sum_{n\in\mathrm{N}} [x_n]^A_n\mathrm{N} = \sum_{n\in\mathrm{N}} A(t_n)\mathrm{N} = \sum_{n\in\mathrm{N}} A(\sigma\lceil n+1)\mathrm{N} = A(\sigma \wedge \langle 0\rangle).$$

Let $y \in A$, $x \neq y$. Fix k such that $x\rho_k y$. Thus for $n \geq k$ the above procedure leads to the same sequences t_n, hence to the same s. •

As an example, for every $\sigma \in \mathcal{S}(\omega)$ and every n, we get $\mathrm{f}B(\sigma, n) = A(\sigma \wedge \langle 0\rangle)$ for each of the chains

$$B(\sigma, n) = A(\sigma\lceil n)\mathrm{Z} + \sum_{i\in\omega} A(\sigma\lceil n+i+1)\mathrm{N}.$$

The attempt to characterise uniform chains of rank ω will be resumed in section 5.

[3] Chains of rank finite over a limit

This is the main section of the paper. It describes a way to extend the Pastjin chain constructors to finite successors of limit ordinals λ using the chains of rank λ as 'building blocks'.

[3.1] Chains of rank ω+finite

(3.1) **Definition** (α successor) Call $x \in A$ an α *successor (in A)* if α is minimal $< \mathrm{rk}(A)$ such that $[x]_\alpha$ has an immediate predecessor in $[x]_{\alpha+1}$.

Say that x is an α *successor in* A/β, where $\alpha < \mathrm{rk}(A/\beta)$ and $\alpha < \mathrm{rk}(A/\beta)$, if $[x]_\beta$ is an α successor in A/β.

Examples:

(a) Every successor element in A (i.e., every element that has an immmediate predecessor in $[x]_1 \in \{\mathrm{N}, \mathrm{Z}\}$) is a 0-successor.

(b) If A has a least element x, then x is a $\mathrm{rk}(A)$ successor. •

(3.2) **Lemma** (Successor induction) Suppose x is an n successor in $A(t)$, $t \in \mathcal{S}$, where $n \in \{0, \ldots, |t|-1\}$ and let $t \in \{s^+, s^-\}$ for some $s \in \mathcal{S}$. Assume a canonical embedding of $A(t)$ into $A(s)$.

(a) x is an $n+1$ successor in $A(s)$ iff $n = |t|-1$ and $t = s^+$.

(b) x is an n successor in $A(s)$ in all other cases, i.e., iff $n < |t|-1$ or $t = s^-$.

In other words, x is an n successor in $A(s)$ iff x is an n successor in $A(t)$, except when $n = |t|-1$ and $t = s^+$, in which case x is an $n+1$ successor in $A(s)$ iff x is an n successor in $A(t)$.

Proof If $n = |t|-1$ and $t = s^+$, then x is minimal in $A(t)$ and $A(s) = A(t) + A(s^-)\mathrm{N}$; so x is minimal in $A(s)$, so n is a $n+1 = \mathrm{rk}(A(s))$ successor. Conversely, if x is an $n+1$ succssor, then x is minimal in $A(s)$, and since it is also minimal in $A(t)$ for $t \in \{s^+, s^-\}$ we cannot have $t = s^-$ (since otherwise $A(t)$ would be mapped onto one of the successor n components of $A(s)$ of form $A(s)$, contradicting minimality of x in $A(s)$), so $t = s^+$.

If $n < |t|-1$ it suffices to note that $[x]_i^{A(t)} = [x]_i^{A(s)}$ for $i = n, n+1$ so x is an n successor in $A(t)$ iff it is an n successor in $A(s)$.

That leaves the case $n = |t|-1$ and $t = s^-$. In this case if x is an n successor in $A(t)$ and $\mathrm{rk}(A(t) = n$, then x in minimal in $A(t)$; but $A(t)$ is embedded as a successor n component into $A(s)$, so n is now minimal such that x is a successor mod n, so x is also an n successor in $A(s)$. Conversely, if x is an n successor in $A(s)$, x is minimal in some successor n component of $A(s)$, i.e., x is minimal in some $A(s^-)$, so $t = s^-$. •

By the component tail lemma, for any $x, x' \in A \in \mathcal{U}(\omega, \omega * 2)$, $\mathrm{f}[x]_\omega \cong \mathrm{f}[x']_\omega$.

(3.3) **Lemma** (Successor isomorphism) For $x, x' \in A \in \mathcal{U}(\omega, \omega * 2)$, if both $[x]_\omega$ and $[x']_\omega$ are n successors in A/ω for $n < \mathrm{rk}(A/\omega) < \omega$, then $[x]_\omega \cong [x']_\omega$.

Proof Let y, y' be immediate predecessors of x, x' mod $\omega + n$ Now,

$$\mathrm{f}_y A = \mathrm{f}_y [y]_{\omega+n} + \sum_{i=0}^{m} [x]_{\omega+n+i} \mathrm{N}$$

where $m = \mathrm{rk}(A/\omega)$. By the block lemma, the isomorphism between $\mathrm{f}_y A$ and $\mathrm{f}_{y'} A$ maps the second $\omega + n$ component of $\mathrm{f}_y A$ isomorphically onto the second $\omega + n$ component of $\mathrm{f}_{y'} A$. Thus, $[x]_{\omega+n} \cong [x']_{\omega+n}$. By finite induction over $i = n-1, \ldots, 0$, using that x and x' have no immediate predecessor mod $\omega + i$ for $i = 0, \ldots, n-1$, it follows that $[x]_\omega \cong [x']_\omega$. The induction step is simply to note that if $[x]_{\omega+i+1} \cong [x']_{\omega+i+1}$, then also for the respective initial components of these, $[x]_{\omega+i} \cong [x']_{\omega+i}$, by another application of the block lemma. •

(3.4) **Definition** ($\mathcal{S}$-domains) Define $\mathcal{D}(s)$ to be the closure of $\{s\}$, $(s \in \mathcal{S}(<\omega))$ under $t \to t^+$ and $t \to t^-$; formally, $\mathcal{D}(\langle\rangle) = \{\langle\rangle\}$ and if $s^+ \in \mathcal{S}$, $\mathcal{D}(s) = \{s\} \cup \mathcal{D}(s^+) \cup \mathcal{D}(s^-)$, else (if $s^+ \notin \mathcal{S}$) $\mathcal{D}(s) = \{s\} \cup \mathcal{D}(s^-)$.

(3.5) **Definition** ($\mathcal{S}$-trees) The *binary tree* $\mathcal{T}(s)$ of $s \in \mathcal{S}(<\omega)$ is the tree of height $|s|$ defined by

(a) [Level 0] The top-node (root) of $\mathcal{T}(s)$ is $\langle s; 0\rangle$, i.e., $\mathrm{Lev}_0 \mathcal{T}(s) = \{\langle s; 0\rangle\}$.

(b) [Level $i \to$ level $i+1$] Suppose $\sigma = \langle t, k\rangle \in \mathrm{Lev}_i \mathcal{T}(s)$, where $i < |s| - 1$.

(b.0) If $t^+ \notin \mathcal{S}$, i.e., if $t(m) = m$ for $m = |t| - 1$, then $\sigma^0 = \langle t^-; 0\rangle$ is the only extension of σ in $\mathrm{Lev}_{i+1} \mathcal{T}(s)$.

(b.1) If $t^+ \in \mathcal{S}$, i.e., if $t(m) < m$ for $m = |t| - 1$, then $\sigma^1 = \langle t^+; k+1\rangle$ and $\sigma^0 = \langle t^-, 0\rangle$ are the only extensions of σ in $\mathrm{Lev}_{i+1} \mathcal{T}(s)$.

Order the extensions of $\sigma \in \mathcal{T}(s)$ by setting $\sigma^1 < \sigma^0$ whenever σ^1 is defined. This will induce an ordering on each level of $\mathcal{T}(s)$, and hence on the leaves of $\mathcal{T}(s)$, by setting in addition $\sigma^{0,1} < \tau^{0,1}$ on level $k+1$ whenever $\sigma < \tau$ on level k.

Notation Denote by $\lambda^s(i)$ the ith leaf of $\mathcal{T}(s)$.

Corollary

(a) $\mathcal{T}(s^-)$ is a sub-tree of $\mathcal{T}(s)$. - Hence:

(b) $\mathcal{T}(s^{-k})$ is a sub-tree of $\mathcal{T}(s)$, for every $k = 1, \ldots, |s| - 1$. - More specifically:

(c) Let k be maximal $\leq |s|$ such that $s^{+k} \in \mathcal{S}$. Then $\mathcal{T}(s^{-i})$ is the sub-tree of $\mathcal{T}(s)$ extending node $\langle s^{+i}; i\rangle$ via node $\langle s^{(+i)-}, 0\rangle$ ('to the right'), for every $i = 1, \ldots, k$, and $\mathcal{T}(s^{-(k+1)})$ is the sole extension of $\mathcal{T}(s)$ at node $\langle s^{+k}; k\rangle$.

Proof (a) The top-node of $\mathcal{T}(s^-)$ is $\langle s^-; 0\rangle$, which an extension (to the 'right', if $s^+ \in \mathcal{S}$, else the only) of the top-node $\langle s; 0\rangle$ of $\mathcal{T}(s)$. (b) follows by induction, and (c) follows from $s^{(+k)-} = s^{-(k+1)}$. •

(3.6) **Definition** (Association) For each $x \in A = A(s)$ define the *association* between x and nodes σ on $\mathcal{T}(s)$ as follows:

(a) $x \to \langle s, 0\rangle$

(b) Supposing $x \to \sigma = \langle t, k\rangle \in \mathcal{T}(s)$ for $|t| > 1$. Then

  (b.0) $x \to \sigma^1 = \langle t^+, k+1\rangle$ if $[x]_{|t|-1}$ is the initial $|t| - 1$ component of $[x]_{|t|}$;

  (b.1) $x \to \sigma^0 = \langle t^-, 0\rangle$ if $[x]_{|t|-1}$ is a successor $|t| - 1$ component of $[x]_{|t|}$.

Also say that x is associated to a branch in $\mathcal{T}(s)$ if x is associated to each node of that branch.

Examples

$$\mathcal{T}(\langle 0\rangle) \qquad \{ \qquad \langle\langle 0\rangle;0\rangle$$















Let $x \in A(t)$ where $t \in \mathcal{S}$ and assume $t \in \{s^+, s^-\}$ for some $s \in \mathcal{S}$ and suppose $A(t)$ canonically embedded in $A(s)$. Suppose x is associated to the nodes of a branch in $\mathcal{T}(t)$ containing a node $\langle u, 0\rangle$, $u \in \mathcal{D}(t) \subset \mathcal{D}(s)$. Then x is also associated to node $\langle u, 0\rangle$ in $\mathcal{T}(s)$. Namely, in $\mathcal{T}(s)$, x is associated to $\langle t, 1\rangle$ if $t = s^+$, else to $\langle t, 0\rangle$. In the latter case it follows immediately that the branch in $\mathcal{T}(s)$ extending $\langle t, 0\rangle$ to which x is associated is the same as the branch in $\mathcal{T}(t)$. If $t = s^+$ these branches differ in the initial segment of nodes σ for which x gets associated to the left successor σ^1 of σ. I.e., let i be such that $x \to \langle t^{+i}; i\rangle$ and $x \to \langle t^{(+i)-}; 0\rangle$. Such a node must exist if x is associated to some node of form $\langle u, 0\rangle$. Then, in $\mathcal{T}(t)$,
(a) $x \to \langle t^{+j}; j\rangle$ for $j = 0, \ldots, i$, $x \to \langle t^{(+i)-}; 0\rangle$,
while in $\mathcal{T}(s)$,
(b) $x \to \langle t^{+j}; j+1\rangle = \langle s^{+(j+1)}; j+1\rangle$ for $j = 0, \ldots, i$ if $t = s^+$;
(c) $x \to \langle t^{+j}; j\rangle = \langle s^{-(+j)}; j\rangle$ for $j = 0, \ldots, i$ if $t = s^-$.
But in either tree, $x \to \langle t^{(+i)-}; 0\rangle = \langle s^{\pm(+i)-}; 0\rangle$, so the branches associated to x extending the latter nodes in $\mathcal{T}(t)$ and $\mathcal{T}(s)$, respectively, are identical. In particular, the branches associated to x extending $\langle u, 0\rangle$ in either $\mathcal{T}(t)$ or $\mathcal{T}(s)$ are identical.

Notation (Leaf number) Let $\ell(s)$ denote the number of leaves in $\mathcal{T}(s)$. Note that $\ell(s) = \ell(s^+) + \ell(s^-)$ if s^+ is defined and $\ell(s) = \ell(s^-)$ else. $\ell(s)$ will figure as the length of certain arrays to be defined.

A basic list of $\ell(s)$ examples is:

$\ell(\langle 0\rangle) = 1$
$\ell(\langle 0,0\rangle) = \ell(\langle 0\rangle) + \ell(\langle 0\rangle) = 1 + 1 = 2$
$\ell(\langle 0,0,0\rangle) = \ell(\langle 0,0\rangle) + \ell(\langle 0,0\rangle) = 2 + 2 = 4$
$\ell(\langle 0,0,2\rangle) = \ell(\langle 0,0\rangle) = 2$
$\ell(\langle 0,1,1\rangle) = \ell(\langle 0,1\rangle) + \ell(\langle 0,1\rangle) = 1 + 1 = 2$
$\ell(\langle 0,1\rangle) = \ell(\langle 0\rangle) = 1$
$\ell(\langle 0,0,1\rangle) = \ell(\langle 0,1\rangle) + \ell(\langle 0,0\rangle) = 1 + 2 = 3$
$\ell(\langle 0,1,0\rangle) = \ell(\langle 0,0\rangle) + \ell(\langle 0,1\rangle) = 2 + 1 = 3$
$\ell(\langle 0,1,2\rangle) = \ell(\langle 0,1\rangle) = 1$

(3.7) **Definition** (Weight) For a node $\sigma \in \mathcal{T}(s)$ set $\psi^s(\sigma) = k$ just if $\sigma = \langle t, k\rangle$ for some $t \in \mathcal{D}(s)$. For $x \in A(s)$ set $\psi^s(x) = \psi^s(\sigma)$ for the leaf $\sigma \in \mathrm{Lev}_{|s|}(\mathcal{T}(s))$ associated to x. For $i < \ell(s)$ set $\psi^s(i) = \psi^s(\lambda^s(i))$.

(3.8) **Lemma** (Weight induction) Suppose $x \in A(t)$ has weight n, where $t \in \mathcal{S}$ and $n \in \{0, \ldots, |t|-1\}$ and suppose $t \in \{s^+, s^-\}$ for some $s \in \mathcal{S}$. Then for all $n \in \{0, \ldots, |t|-1\}$, $\psi^s(x) = n$ iff $\psi^t(x) = n$, except when $n = |t| - 1$ and $t = s^+$, in which case $\psi^s(x) = n + 1$ iff $\psi^t(x) = n$.

Proof Let $t \in \{s^+, s^-\}$, $n = |t| - 1$ and $x \in A(t)$. Then

$$\begin{aligned}
\psi^t(x) = n \quad & \text{iff } x \to \sigma \text{ for some } \sigma \in \mathrm{Lev}_n(\mathcal{T}(t)) \text{ such that } \psi^t(\sigma) = n \\
& \text{iff } x \to \sigma = \langle\langle 0\rangle; n\rangle \\
& \text{iff } t^{+n} = \langle 0\rangle \in \mathcal{S} \text{ and } x \to \langle t^{+n}; n\rangle = \langle\langle 0\rangle; n\rangle \\
& \text{iff } t = t^- \wedge \langle 0\rangle \text{ and } x \to \langle t^{+n}; n\rangle = \langle\langle 0\rangle; n\rangle \\
& \text{iff } x \to \langle t^{+i}; i\rangle, \text{ for each } i = 0, \ldots, n,
\end{aligned}$$

i.e., x is associated to each node of the left-most branch of $\mathcal{T}(t)$, given by $\langle t, 0\rangle, \langle t^+, 1\rangle, \ldots, \langle t^{+i}; i\rangle, \ldots, \langle\langle 0\rangle; n\rangle$.

Case (a): Suppose the $\psi^t(x) = n = |t| - 1$.

Case (a.0): If also $t = s^+$, then x is associated to each node $\langle s^{+i}; i\rangle$, for $i = 0, \ldots, n+1$, of the left-most branch of $\mathcal{T}(s)$, given by $\langle s, 0\rangle, \langle t, 1\rangle, \langle t^+, 2\rangle, \ldots, \langle t^{+i}, i+1\rangle, \ldots, \langle\langle 0\rangle; n+1\rangle$, and so $\psi^s(x) = n+1$.

Case (a.1): If on the other hand $t = s^-$, then x is associated to nodes $\langle s, 0\rangle, \langle s^-, 0\rangle = \langle t, 0\rangle$, and hence each node $\langle s^{-(+i)}; i\rangle = \langle t, i\rangle$ for $i = 0, \ldots, n$ of the branch $\langle s, 0\rangle, \langle t, 0\rangle, \langle t^+, 1\rangle, \ldots, \langle t^{+i}; i\rangle, \ldots, \langle\langle 0\rangle; n\rangle$ of $\mathcal{T}(s)$, and so $\psi^s(x) = n$.

Case (b): Suppose on the other hand that $\psi^t(x) = n < |t| - 1$. Then there is some least $i < n$ such that x is associated with each node of the branch $\langle t, 0\rangle, \ldots \langle t^{+i}; i\rangle, \langle t^{(+i)-}; 0\rangle$. (I.e., for some $i < n$, x must get associated with $\langle t^{(+i)-}; 0\rangle$, else we are back in case (a.0). For the least such i, x is associated with the branch just described.) Hence in $A(s)$, x is associated to each node of the branch $\langle s, 0\rangle, \langle t, k_0\rangle, \ldots \langle t^{+i}; k_i\rangle, \langle t^{(+i)-}; 0\rangle$, where $k_j = j+1$ if $t = s^+$ and $k_j = j$ if $t = s^-$. The remaining nodes of the branches in either $\mathcal{T}(t)$ or $\mathcal{T}(s)$ with which x gets associated are now identical. Thus $\psi^t(x) = \psi^s(x)$, as required. •

Corollary Let $s \in \mathcal{S}$ and let $n = |s| - 1$.

(a)

$$\psi^s(0) = \begin{cases} \psi^{s^+}(0) + 1 & \text{if } s^{+n} = \langle 0\rangle \in \mathcal{S} \text{ (i.e. } s = s^- \wedge \langle 0\rangle) \\ \psi^{s^+}(0) & \text{else} \end{cases}$$

(b) For $0 < i < \ell(s)$,

$$\psi^s(i) = \begin{cases} \psi^{s^+}(i) & \text{if } s^+ \in \mathcal{S} \text{ and } i < \ell(s^+) \\ \psi^{s^-}(\ell(s^+) + i) & \text{if } s^+ \in \mathcal{S} \text{ and } \ell(s^+) \leq i < \ell(s) = \ell(s^+) + \ell(s^-) \\ \psi^{s^-}(i) & \text{if } s^+ \notin \mathcal{S} \text{ and } i < \ell(s) = \ell(s^-) \end{cases}$$

Corollary

(a) If $s = s^- \wedge \langle 0\rangle$ (and so $s^+ \in \mathcal{S}$), then for all $i < \ell(s) = \ell(s^+) + \ell(s^-)$

$$\psi^s(i) = \begin{cases} \psi^{s^+}(i) + 1 & \text{for } i = 0 \\ \psi^{s^+}(i) & \text{for } 0 < i < \ell(s^+) \\ \psi^{s^-}(\ell(s^+) + i) & \text{for } \ell(s^+) \leq i < \ell(s) \end{cases}$$

(b) If $s^+ \in \mathcal{S}$ but $s \neq s^- \wedge \langle 0\rangle$, then for all $i < \ell(s) = \ell(s^+) + \ell(s^-)$

$$\psi^s(i) = \begin{cases} \psi^{s^+}(i) & \text{for } i < \ell(s^+) \\ \psi^{s^-}(\ell(s^+) + i) & \text{for } \ell(s^+) \leq i < \ell(s) \end{cases}$$

(c) If $s^+ \notin \mathcal{S}$, then for all $i < \ell(s) = \ell(s^-)$

$$\psi^s(i) = \psi^{s^-}(i) \text{ for } i < \ell(s) = \ell(s^-)$$

Examples:

(i) Compare $\mathcal{T}(s)$ for $s = \langle 0,0,1\rangle$ or $s = \langle 0,0,0,1\rangle$ with those for $s = \langle 0,1,0\rangle$ or $s = \langle 0,0,1,0\rangle$ given above. The latter are examples in which case (a) of the preceding corollary applies.

(ii) Compare $\mathcal{T}(s)$ for $s = \langle 0,0,2,1\rangle$, $s = \langle 0,0,1,2\rangle$ and $s = \langle 0,0,1,3\rangle$ with $\mathcal{T}(\langle 0,0,1\rangle)$.


$\mathcal{T}(\langle 0,0,2,1\rangle)$
$\langle\langle 0,0,2,1\rangle;0\rangle$
$\langle\langle 0,0,1\rangle;1\rangle$
$\mathcal{T}(\langle 0,0,2\rangle)$
$\langle\langle 0,1\rangle;1\rangle$
$\mathcal{T}(\langle 0,0\rangle)$
$\mathcal{T}(\langle 0\rangle)$


$\mathcal{T}(\langle 0,0,1,2\rangle)$
$\langle\langle 0,0,1,2\rangle;0\rangle$
$\langle\langle 0,0,2\rangle;1\rangle$
$\mathcal{T}(\langle 0,0,1\rangle)$
$\mathcal{T}(\langle 0,0\rangle)$


$\mathcal{T}(\langle 0,0,1,3\rangle)$
$\langle\langle 0,0,1,3\rangle;0\rangle$
$\mathcal{T}(\langle 0,0,1\rangle)$

(iii) Compare $\mathcal{T}(s)$ for $s = \langle 0,1,0,1\rangle$, $s = \langle 0,1,0,2\rangle$, and $s = \langle 0,1,0,3\rangle$ with $\mathcal{T}(\langle 0,1,0\rangle)$.


$\mathcal{T}(\langle 0,1,0,0\rangle)$
$\langle\langle 0,1,0,0\rangle;0\rangle$
$\langle\langle 0,1,0\rangle;1\rangle$
$\mathcal{T}(\langle 0,1,0\rangle)$
$\langle\langle 0,0\rangle;2\rangle$
$\mathcal{T}(\langle 0,1\rangle)$
$\langle\langle 0\rangle;3\rangle$
$\mathcal{T}(\langle 0\rangle)$


$\mathcal{T}(\langle 0,1,0,1\rangle)$
$\langle\langle 0,1,0,1\rangle;0\rangle$
$\langle\langle 0,1,1\rangle;1\rangle$
$\mathcal{T}(\langle 0,1,0\rangle)$
$\langle\langle 0,1\rangle;1\rangle$
$\mathcal{T}(\langle 0,1\rangle)$
$\mathcal{T}(\langle 0\rangle)$


$\mathcal{T}(\langle 0,1,0,3\rangle)$
$\langle\langle 0,1,0,3\rangle;0\rangle$
$\mathcal{T}(\langle 0,1,0\rangle)$

(3.9) **Lemma** (Weight isomorphism)

(a) Every $x \in A(s)$ is a $\psi^s(x)$ successor.

(b) Suppose $A/\omega \cong A(s)$. Then for all $x, x' \in A$, if $\psi^s([x]_\omega) = \psi^s([x']_\omega) < \mathrm{rk}(A/\omega) = \mathrm{rk}(A(s)) < \omega$ then $[x]_\omega \cong [x']_\omega$.

Proof (i) By induction on $\mathrm{rk}(A(s)) = |s| - 1$.

For $s = \langle 0 \rangle$ the claim is vacuously true.

For $|s| = 2$, $\psi^s(x) = 1$ iff $x \to \langle\langle 0\rangle; 1\rangle = \langle s, 0\rangle^+$ iff x is minimal in $A(s)$ iff x is a $\mathrm{rk}(A(s)) = 1$ successor, and $\psi^s(x) = 0$ iff x is a 0 successor (i.e., has an immediate predecessor in $[x]_1 = \mathrm{N}$).

For $s = \langle 0, 0\rangle$, there is exactly one $x \in A(s) = \mathrm{N}$ for which each statement in the first chain of implications holds. For any other element in N, and any element in $A(\langle 0, 1\rangle) = \mathrm{Z}$, both statements of the second implication apply.

Let $s \in \mathcal{S}$ and suppose the statement is true for $t \in \{s^+, s^-\} \cap \mathcal{S}$, i.e., x is an n successor in $A(t)$ iff $\psi^t(x) = n$, for all $n \in \{0, \ldots, |s| - 1\}$. Assume $A(t)$ embeds canonically into $A(s)$. By the successor induction lemma, for all $n \in \{0, \ldots, |t| - 1\}$, x is an n successor in $A(s)$ iff x is an n successor in $A(t)$, except when $n = |t| - 1$ and $t = s^+$, in which case x is an $n + 1$ successor in $A(s)$ iff x is an n successor in $A(t)$. By the weight induction lemma, for all such n, $\psi^s(x) = n$ iff $\psi^t(x) = n$, except when $n = |t| - 1$ and $t = s^+$, in which case $\psi^s(x) = n + 1$ iff $\psi^t(x) = n$. Thus, by inductive hypothesis, x is an n successor in $A(s)$ iff $\psi^s(x) = n$, for all $n \in \{0, \ldots, |s| - 1\}$.

(ii) Follows from (i) and the successor isomorphism lemma: Suppose $A/\omega \cong A(s)$. If $\psi^s([x]_\omega) = \psi^s([x']_\omega) = n$ for $x, x' \in A$ and $n < \mathrm{rk}(A/\omega) \in \omega$, then by (i), $[x]_\omega$ and $[x']_\omega$ are n successors in A/ω. Again by the successor isomorphism lemma, $[x]_\omega \cong [x']_\omega$, as required. •

(3.10) **Definition** (Plus finite rank Pastjin uniform chain constructor — long version) Let $M_i \in \mathcal{U}(\omega)$, $i \in \omega$. Define

$$A(\langle 0\rangle)[M_0] = 1[M_0] = M_0$$

and for $\ell(s)$ arrays $\langle M_0, \ldots, M_{\ell(s)-1}\rangle \in {}^{\ell(s)}\mathcal{U}(\omega)$,

$$A(s)[M_0, \ldots, M_{\ell(s)-1}] = \begin{cases} A(s^-)[M_0, \ldots, M_{\ell(s^-)-1}].\mathrm{Z} & \text{if } s(n) = n \\ A(s^+)[M_0, \ldots, M_{\ell(s^+)-1}] + A(s^-)[M_{\ell(s^+)}, \ldots, M_{\ell(s)-1}].\mathrm{N} & \text{if } s(n) < n \end{cases}$$

where $n = |s| - 1$ (and where the '.' is meant to regulate precedence of operations in the usual fashion). The sequence $\mathbf{M} = \mathbf{M}_{[0,\ell(s))} = \langle M_0, \ldots, M_{\ell(s)-1}\rangle$ is subsequently called the *array* (of building blocks) of the constructor, and $A(s^+)[M_0, \ldots, M_{\ell(s^+)-1}]$, respectively $A(s^-)[M_0, \ldots, M_{\ell(s^-)-1}]$ or $A(s^-)[M_{\ell(s^+)}, \ldots, M_{\ell(s)-1}]$ (whichever applies) are referred to as the 'plus', respectively 'minus' sub-components of $A(s)[M_0, \ldots, M_{\ell(s)-1}]$.

Note that $A(s)[\mathbf{M}]/\omega \cong A(s)$.

A basic list of such chains of rank finite over ω contructed with $\ell(s)$ arrays is:

$A(\langle 0\rangle)[M_0] = M_0$

$A(\langle 0, 0\rangle)[M_0, M_1] = M_0 + M_1\mathrm{N}$ $\quad$ $A(\langle 0, 1\rangle)[M_0] = M_0\mathrm{Z}$

$A(\langle 0, 0, 0\rangle)[M_0, M_1, M_2, M_3] = M_0 + M_1\mathrm{N} + (M_2 + M_3\mathrm{N})\mathrm{N}$

$A(\langle 0, 0, 1\rangle)[M_0, M_1, M_2] = M_0\mathrm{Z} + (M_1 + M_2\mathrm{N})\mathrm{N}$

$A(\langle 0, 0, 2\rangle)[M_0, M_1] = (M_0 + M_1\mathrm{N})\mathrm{Z}$ $\quad$ $A(\langle 0, 1, 0\rangle)[M_0, M_1, M_2] = M_0 + M_1\mathrm{N} + M_2\mathrm{Z}\mathrm{N}$

$A(\langle 0, 1, 1\rangle)[M_0, M_1] = M_0\mathrm{Z} + M_1\mathrm{Z}\mathrm{N}$ $\quad$ $A(\langle 0, 1, 2\rangle)[M_0] = M_0\mathrm{Z}^2$

Arbitrary chains of the above form will not in general be uniform again (even with $M_i \in \mathcal{U}(\omega)$).

1. Order-isomorphisms between tails of chains induce (restrict to) order isomorphisms between the initial components of these tails. The set of initial ω components of tails of $A = A(s)[M_0, \ldots, M_{\ell(s)-1}]$ is the set of tails $\mathrm{f}M_0, \ldots, \mathrm{f}M_{\ell(s)-1}$. These must be pairwise isomorphic, if A is uniform.
2. Order-isomorphisms between tails of chains induce isomorphisms between the components that are 'successors' to the initial components of the tail chains. This yields a partition of the set $M_0, \ldots, M_{\ell(s)-1}$ into subsets of pairwise isomorphic members. For instance, any two M_i, M_j that are be successor ω components in (tails) of A must be isomorphic, any two M_i, M_j that can be initial ω components in successor $\omega + 1$ components in tails of A must be isomorphic, etc.

(3.11) **Lemma** (Uniformity over limit rank) If $A = A(s)[M_0, \ldots, M_{\ell(s)-1}]$ is uniform, then

$$\begin{array}{ll} \text{(a)} & \mathrm{f}M_i = \mathrm{f}M_j \text{ for all } i, j < \ell(s), \text{ and} \\ \text{(b)} & M_i = M_j \text{ whenever } \psi^s(i) = \psi^s(j). \end{array} \tag{3.1}$$

Proof (a) is immediate by the fact that for any $x, x' \in A$, $\mathrm{f}[x]_\omega \cong \mathrm{f}[x']_\omega$ by the component tail isomorphism lemma.

(b) By the weight isomorphism lemma it suffices to show that $\psi^s(M_i) = \psi^s(i)$ for all $i < \ell(s)$. For this in turn it suffices to show that $M_i \to \lambda^s(i)$ for all $i < \ell(s)$.

This follows by induction on $\mathrm{rk}(A(s))$, $s \in \mathcal{S}(< \omega)$. By definition of the association of $x \in A(s)$ to nodes $\sigma \in \mathcal{T}(s)$,

(a) $M_i \to \langle s, 0\rangle \in \mathcal{T}(s)$ for all $i < \ell(s)$, and

(b.0) if $s^+ \in \mathcal{S}$, then $M_i \to \langle s^+, 1\rangle \in \mathcal{T}(s)$ for all $i < \ell(s^+)$, $M_i \to \langle s^-, 0\rangle \in \mathcal{T}(s)$ for all $\ell(s)^+ \le i < \ell(s)$, i.e., $M_{lf(s^+)+i} \to \langle s^-, 0\rangle \in \mathcal{T}(s)$ for all $i < \ell(s^-)$, while

(b.1) if $s^+ \notin \mathcal{S}$, then $M_i \to \langle s^-, 0\rangle \in \mathcal{T}(s)$ for all $i < \ell(s) = \ell(s^-)$.

Suppose $s^+ \in \mathcal{S}$, then, if $\lambda^{s^+}(i) = \langle\langle 0\rangle; k\rangle$ for $i < \ell(s^+)$, then $\lambda^s(i) = \langle\langle 0\rangle; k+1\rangle$ for $i = 0$, and $\lambda^s(i) = \langle\langle 0\rangle; k\rangle$ for $0 < i < \ell(s^+)$; and if $\lambda^{s^-}(i) = \langle\langle 0\rangle; k\rangle$ for $i < \ell(s^-)$, then $\lambda^s(\ell(s^+) + i) = \langle\langle 0\rangle; k\rangle$ for $i < \ell(s^-)$.

Suppose $s^+ \in \mathcal{S}$, then, if $\lambda^{s^-}(i) = \langle\langle 0\rangle; k\rangle$ for $i < \ell(s^-)$, then $\lambda^s(i) = \langle\langle 0\rangle; k\rangle$ for $i < \ell(s)$.

In case (b.1) it follows immediately from the inductive hypothesis that $\psi^s(M_i) = \psi^{s^-}(M_i) = \psi^{s^-}(i) = \psi^s(i)$.

In case (b.0),

(a) $\psi^s(M_0) = \psi^{s^+}(M_0) + 1 = \psi^{s^-}(0) + 1 = \psi^s(0)$,

(b) $\psi^s(M_i) = \psi^{s^+}(M_i) + 1 = \psi^{s^+}(i) = \psi^s(i)$ for $0 < i < \ell(s^+)$, and

(c) $\psi^s(M_{\ell(s^+)+i}) = \psi^{s^+}(M_i) = \psi^{s^+}(i) = \psi^s(\ell(s^+) + i)$ for $\ell(s^+) \le i \le \ell(s)$,

again by inductive hypothesis. •

Examples: By earlier calculations,

(i) $\ell(\langle 0, 1, 0\rangle) = \ell(\langle 0, 0\rangle) + \ell(\langle 0, 1\rangle) = (\ell(\langle 0\rangle) + \ell(\langle 0\rangle)) + \ell(\langle 0\rangle) = 3$,

(ii) $\ell(\langle 0, 0, 1\rangle) = \ell(\langle 0, 1\rangle) + \ell(\langle 0, 0\rangle) = \ell(\langle 0\rangle) + (\ell(\langle 0\rangle) + \ell(\langle 0\rangle)) = 3$.

Thus, for (i),

$$\begin{aligned} A(\langle 0,1,0\rangle)[M_0,M_1,M_2] &= A(\langle 0,0\rangle)[M_0,M_1] + A(\langle 0,1\rangle)[M_2].\mathrm{N} \\ &= (1+\mathrm{N})[M_0,M_1] + \mathrm{Z}[M_2].\mathrm{N} \\ &= M_0 + M_1\mathrm{N} + M_2\mathrm{ZN}, \end{aligned}$$

$\mathcal{T}(\langle 0,1,0\rangle)$



So, $\psi^{\langle 0,1,0\rangle}(M_0) = 2$, $\psi^{\langle 0,1,0\rangle}(M_1) = \psi^{\langle 0,1,0\rangle}(M_2) = 0$, so $M_1 \cong M_2$. Hence, the uniform sets contucted from $A(\langle 0,1,0\rangle)$ over uniform rank ω chains are of form $A(\langle 0,1,0\rangle)[M_0,M_1,M_1] = M_0 + M_1\mathrm{N} + M_1\mathrm{ZN}$. For (ii):

$$\begin{aligned} A(\langle 0,0,1\rangle)[M_0,M_1,M_2] &= A(\langle 0,1\rangle)[M_0] + A(\langle 0,0\rangle)[M_1,M_2].\mathrm{N} \\ &= \mathrm{Z}[M_0] + (1+\mathrm{N})[M_1,M_2].\mathrm{N} \\ &= M_0\mathrm{Z} + (M_1 + M_2\mathrm{N})\mathrm{N}, \end{aligned}$$

$\mathcal{T}(\langle 0,0,1\rangle)$



So, $\psi^{\langle 0,0,1\rangle}(M_0) = \psi^{\langle 0,0,1\rangle}(M_2) = 0$, $\psi^{\langle 0,0,1\rangle}(M_1) = 1$, so $M_0 \cong M_2$. Hence, the uniform sets contucted from $A(\langle 0,0,1\rangle)$ over uniform rank ω chains are of form $A(\langle 0,0,1\rangle)[M_0,M_1,M_0] = M_0\mathrm{Z} + (M_1 + M_0\mathrm{N})\mathrm{N}$. •

To show that condition (a) and (b) is also sufficient for uniformity of A we need to calculate its tails, fA.

[3.2] Tails of rank ω+finite

3.12 **Definition** (Tail array) Given an array $\mathbf{M}_{\ell(s)} = [M_0,\ldots,M_{\ell(s)-1}]$, define the array $\mathrm{f}_s(\mathbf{M}_{\ell(s)})$ by induction over $\mathcal{D}(s)$:

First, define

$$\begin{aligned} \mathrm{f}_{\langle 0\rangle}(\mathbf{M}_{\ell(\langle 0\rangle)}) &= \mathrm{f}([M_0]) = [\mathrm{f}M_0] \\ \mathrm{f}_{\langle 0,0\rangle}(\mathbf{M}_{\ell(\langle 0,0\rangle)}) &= \mathrm{f}([M_0,M_1]) = [\mathrm{f}M_0,M_1] \\ \mathrm{f}_{\langle 0,1\rangle}(\mathbf{M}_{\ell(\langle 0,1\rangle)}) &= \mathrm{f}([M_0]) = [\mathrm{f}M_0,M_0] \end{aligned}$$

Next suppose $\mathrm{f}_t\mathbf{M}_{\ell(s)}$ is defined for all $t \in \mathcal{D}(s)$. For a given $\ell(s)$ array $\mathbf{M}_{\ell(s)} = \mathbf{M}_{[0,\ell(s))}$ set

$$\mathrm{f}_s(\mathbf{M}_{\ell(s)}) = \begin{cases} \mathrm{f}_{s^+}(\mathbf{M}_{\ell(s^+)}) \ \wedge \ \mathbf{M}_{[\ell(s^+),\ell(s))} & \text{if } s^+ \in \mathcal{S}, \\ \mathrm{f}_{s^-}(\mathbf{M}_{\ell(s)}) \ \wedge \ \mathbf{M}_{\ell(s^-)} & \text{else (i.e., } s^+ \notin \mathcal{S}). \end{cases}$$

Tail-operation shall take priority over concatenation.

Note that the definition is consistent with the fact that

$$\mathbf{M}_{\ell(s)} = \begin{cases} \mathbf{M}_{\ell(s^+)} \ \wedge \ \mathbf{M}_{[\ell(s^+),\ell(s))} = \mathbf{M}_{[0,\ell(s^+))} \ \wedge \ \mathbf{M}_{[\ell(s^+),\ell(s))} & \text{if } s^+ \in \mathcal{S} \\ \mathbf{M}_{\ell(s^-)} = \mathbf{M}_{[0,\ell(s^-))} & \text{else.} \end{cases}$$

The index 's' on the f operator is necessary in order to disambiguate definiens $\mathrm{f}_{s^-}\mathbf{M}_{\ell(s^-)}$ from definiendum $\mathrm{f}_s\mathbf{M}_{\ell(s)}$ when $s^+ \notin \mathcal{S}$, as in this case $\mathbf{M}_{\ell(s)} = \mathbf{M}_{\ell(s^-)}$. (E.g., we may have $\mathbf{M}_{\ell(\langle 0\rangle)} = [M] = \mathbf{M}_{\ell(\langle 0,1\rangle)}$, but the array of ω blocks for $\mathrm{f}A(\langle 0,1\rangle)\mathbf{M}_{\ell(\langle 0,1\rangle)} = \mathrm{f}M + M\mathrm{N}$ contains two elements ($\mathrm{f}M$ and M), while that for $\mathrm{f}A(\langle 0\rangle)\mathbf{M}_{\ell(\langle 0\rangle)} = \mathrm{f}M$ contains only one ($\mathrm{f}M$), i.e., f-operations on $[M]$ depend on the sequence for which $[M]$ is the array.)

Example: Let's calculate $\mathrm{f}A$ for $A = A(\langle 0,0,1,2\rangle)[M_0, M_1, M_2, M_3, M_4]$. Note that

$$\begin{aligned}
\ell(\langle 0,0,1,2\rangle) &= \ell(\langle 0,0,2\rangle) + \ell(\langle 0,0,1\rangle)\\
&= \ell(\langle 0,0\rangle) + \ell(\langle 0,1\rangle) + \ell(\langle 0,0\rangle)\\
&= \ell(\langle 0\rangle) + \ell(\langle 0\rangle) + \ell(\langle 0\rangle) + \ell(\langle 0\rangle) + \ell(\langle 0\rangle) = 5
\end{aligned}$$

and

$$\begin{aligned}
A(\langle 0,0,1,2\rangle)[M_0, M_1, M_2, M_3, M_4] &= A(\langle 0,0,2\rangle)[M_0, M_1] + A(\langle 0,0,1\rangle)[M_2, M_3, M_4].\mathrm{N}\\
&= A(\langle 0,0\rangle)[M_0, M_1].\mathrm{Z} + (A(\langle 0,1\rangle)[M_2] + A(\langle 0,0\rangle)[M_3, M_4].\mathrm{N}).\mathrm{N}\\
&= (M_0 + M_1\mathrm{N})\mathrm{Z} + (M_2\mathrm{Z} + (M_3 + M_4\mathrm{N})\mathrm{N})\mathrm{N}.
\end{aligned}$$

But $\mathcal{T}(\langle 0,0,1,2\rangle)$ is:


$\langle\langle 0,0,1,2\rangle;0\rangle$
$\langle\langle 0,0,2\rangle;1\rangle$
$\langle\langle 0,0,1\rangle;0\rangle$
$\langle\langle 0,0\rangle,0\rangle$
$\langle\langle 0,1\rangle,1\rangle$
$\langle\langle 0,0\rangle,0\rangle$
$\langle\langle 0\rangle,1\rangle$
$\langle\langle 0\rangle,0\rangle$
$\langle\langle 0\rangle,0\rangle$
$\langle\langle 0\rangle,1\rangle$
$\langle\langle 0\rangle,0\rangle$

Thus $\psi^s(0) = \psi^s(3) = 1$ and $\psi^s(1) = \psi^s(2) = \psi^s(4) = 0$ for $s = \langle 0,0,1,2\rangle$ so assuming A satisfies the uniformity conditions,

$$\mathrm{f}M_i \cong \mathrm{f}M_j \text{ for all } i,j < 5, \quad M_0 \cong M_3, \quad M_1 \cong M_2 \cong M_4$$

A cut-off point x in A may be chosen in five locations generically described by

(1) $x = \langle a, k\rangle \in M_0 \times k$ for $k \in \mathrm{Z}$ from an ω sub-component of form M_0 of the first $\omega + 2$ component of A (of form $(M_0 + M_1\mathrm{N})\mathrm{Z}$).

(2) $x = \langle\langle a, n\rangle, k\rangle \in M_1 \times n, k$ for $n \in \mathrm{N}$ and $k \in \mathrm{Z}$, from an ω sub-component of form $M_1 \times n$ of the first $\omega + 2$ component of A.

(3) $x = \langle\langle a, k\rangle, n\rangle \in M_2 \times k, n$ for $n \in \mathrm{N}$ and $k \in \mathrm{Z}$, from an ω sub-component of form $M_2 \times k$ of a successor $\omega + 2$ component of A (of form $(M_2\mathrm{Z} + (M_3 + M_4\mathrm{N})\mathrm{N}) \times n$).

(4) $x = \langle\langle a, m\rangle, n\rangle \in M_3 \times m, n$ for $m, n \in \mathrm{N}$, from a an ω sub-component of form $M_3 \times m$ of a successor $\omega + 2$ component of A.

(5) $x = \langle\langle\langle a, l\rangle, m\rangle, n\rangle \in M_4 \times l, m, n$ for $l, m, n \in \mathrm{N}$, from an ω sub-component of form $M_4 \times l$ of a successor $\omega + 2$ component of A.

Correspondingly,

$$\mathrm{f}A = \begin{cases} (\mathrm{f}M_0 + M_1\mathrm{N}) + (M_0 + M_1\mathrm{N})\mathrm{N} + (M_2\mathrm{Z} + (M_3 + M_4\mathrm{N})\mathrm{N})\mathrm{N} & \text{cut-off (1)} \\ (\mathrm{f}M_1 + M_1\mathrm{N}) + (M_0 + M_1\mathrm{N})\mathrm{N} + (M_2\mathrm{Z} + (M_3 + M_4\mathrm{N})\mathrm{N})\mathrm{N} & \text{cut-off (2)} \\ (\mathrm{f}M_2 + M_2\mathrm{N}) + (M_3 + M_4\mathrm{N})\mathrm{N} + (M_2\mathrm{Z} + (M_3 + M_4\mathrm{N})\mathrm{N})\mathrm{N} & \text{cut-off (3)} \\ (\mathrm{f}M_3 + M_4\mathrm{N}) + (M_3 + M_4\mathrm{N})\mathrm{N} + (M_2\mathrm{Z} + (M_3 + M_4\mathrm{N})\mathrm{N})\mathrm{N} & \text{cut-off (4)} \\ (\mathrm{f}M_4 + M_4\mathrm{N}) + (M_3 + M_4\mathrm{N})\mathrm{N} + (M_2\mathrm{Z} + (M_3 + M_4\mathrm{N})\mathrm{N})\mathrm{N} & \text{cut-off (5)} \end{cases}$$

which are pairwise isomorphic under the uniformity conditions.

3.13 **Lemma** (Tail uniformity) If $\mathbf{M}_{\ell(s)} = [M_0, \ldots, M_{\ell(s)-1}]$ satisfies the conditions of the uniformity over limit rank, then $\mathrm{f}_{s^+}\mathbf{M}_{[0,\ell(s^+))} = \mathrm{f}_{s^-}\mathbf{M}_{[\ell(s^-),\ell(s))}$.

Proof By definition

$$\mathrm{f}_{s^+}\mathbf{M}_{[0,\ell(s^+))} = \begin{cases} \mathrm{f}_{s^{++}}(\mathbf{M}_{[0,\ell(s^{++}))}) \ \wedge \ \mathbf{M}_{[\ell(s^{++}),\ell(s^+))} & \text{if } s^{++} \in \mathcal{S}, \\ \mathrm{f}_{s^{+-}}(\mathbf{M}_{[0,\ell(s^{+-}))}) \ \wedge \ \mathbf{M}_{[0,\ell(s^{+-}))} & \text{else (i.e., if } s^{++} \notin \mathcal{S}); \end{cases}$$

while

$$\mathrm{f}_{s^-}(\mathbf{M}_{[\ell(s^+),\ell(s))}) = \begin{cases} \mathrm{f}_{s^{-+}}(\mathbf{M}_{[\ell(s^+),\ell(s^+)+\ell(s^{-+}))}) & \text{if } s^{-+} \in \mathcal{S} \\ \mathrm{f}_{s^{--}}(\mathbf{M}_{[\ell(s^+),\ell(s))}) \ \wedge \ \mathbf{M}_{[\ell(s^+),\ell(s))} \ \wedge \ \mathbf{M}_{[\ell(s^+)+\ell(s^{-+}),\ell(s))} & \text{else (i.e., if } s^{-+} \notin \mathcal{S}), \end{cases}$$

the latter by applying the definition to the array $\mathbf{M}_{[0,\ell(t))} = \mathbf{M}_{[\ell(s^+),\ell(s))}$ for $t = s^-$.

We shall be tacitly using the fact that $s^{+-} = s^{--}$, and hence $\ell(s^{+-}) = \ell(s^{--})$ and $\mathrm{f}_{s^{+-}} = \mathrm{f}_{s^{--}}$. If $\mathbf{A} = \langle A_1, \ldots, A_n\rangle$ and $\mathbf{B} = \langle B_1, \ldots, B_n\rangle$ are arrays of chains, we write $\mathbf{A} \cong \mathbf{B}$ to express that $A_i \cong B_i$ for $i = 1, \ldots, n$.

If the uniformity over limit rank conditions apply to $\mathbf{M}_{\ell(s)}$, then the following hold:

$$\begin{array}{ll} \text{Case (a)} & s^{++} \in \mathcal{S}, s^{-+} \in \mathcal{S} \ : \ \mathbf{M}_{[\ell(s^{++}),\ell(s^+))} \cong \mathbf{M}_{[\ell(s^+)+\ell(s^{-+}),\ell(s))} \\ \text{Case (b)} & s^{++} \notin \mathcal{S}, s^{-+} \notin \mathcal{S} \ : \ \mathbf{M}_{[0,\ell(s^{+-}))} \cong \mathbf{M}_{[\ell(s^+),\ell(s^+)+\ell(s^-))} \\ \text{Case (c)} & s^{++} \in \mathcal{S}, s^{-+} \notin \mathcal{S} \ : \ \mathbf{M}_{[\ell(s^{++}),\ell(s^+))} \cong \mathbf{M}_{[\ell(s^+),\ell(s))} \\ \text{Case (d)} & s^{++} \notin \mathcal{S}, s^{-+} \in \mathcal{S} \ : \ \mathbf{M}_{[0,\ell(s^{+-}))} \cong \mathbf{M}_{[\ell(s^+)+\ell(s^{-+}),\ell(s))} \end{array}$$

In case (a), note that by leaf-length arithmetic,

$$\mathbf{M}_{[\ell(s^{++}),\ell(s^+))} = \mathbf{M}_{[\ell(s^{++}),\ell(s^{++})+\ell(s^{+-}))}$$
$$\mathbf{M}_{[\ell(s^+)+\ell(s^{-+}),\ell(s))} = \mathbf{M}_{[\ell(s^+)+\ell(s^{-+}),\ell(s^+)+\ell(s^{-+})+\ell(s^{--}))}.$$

This follows from the partial 'tree isomorphisms' between the subtree of $\mathcal{T}(s)$ extending $\langle s^{+-}, 0\rangle$ and the one extending $\langle s^{--}, 0\rangle$, which in turn is immediate form $s^{+-} = s^{--}$. (The aforemetioned subtrees are both isomorphic to the tree $\mathcal{T}(t)$, for $t = s^{+-} = s^{--}$ if we consider trees $\mathcal{T}(s)$ as partial orders over a set of nodes $\langle t, k\rangle \in \mathcal{D}(s) \times |s|$.) Call these subtrees $\mathcal{T}^{+-} = \mathrm{Ext}(\mathcal{T}(s, \langle s^{+-}, 0\rangle)$ and $\mathcal{T}^{--} = \mathrm{Ext}(\mathcal{T}(s, \langle s^{--}, 0\rangle)$, respectively. Let temporarily '$\hookrightarrow$' denote the embeddings of $\mathcal{T}^{+-}$ and $\mathcal{T}^{--}$ into $\mathcal{T} = \mathcal{T}(s)$, and let $[a, b)$

denote the 'interval' between leaves a (inclusive) and b (exclusive) of $\mathrm{Lev}_{|s|}(\mathcal{T})$. Then

$$\begin{aligned}
&[\mathrm{a}] \quad s^{++} \in \mathcal{S}, s^{-+} \in \mathcal{S} \ : \ && \mathrm{Lev}_{|s^{+-}|}(\mathcal{T}^{+-}) \hookrightarrow [\lambda(\ell(s^{++})), \lambda(\ell(s^{++}) + \ell(s^{+-}))),\\
& && \qquad = [\lambda(\ell(s^{++})), \lambda(\ell(s^{+})))\\
& && \mathrm{Lev}_{|s^{--}|}(\mathcal{T}^{--}) \hookrightarrow [\lambda(\ell(s^{+}) + \ell(s^{-+}))), \lambda(\ell(s^{+}) + \ell(s^{-+}) + \ell(s^{--}))),\\
& && \qquad = [\lambda(\ell(s^{+}) + \ell(s^{-+})), \lambda(\ell(s)))\\
&[\mathrm{b}] \quad s^{++} \notin \mathcal{S}, s^{-+} \notin \mathcal{S} \ : \ && \mathrm{Lev}_{|s^{+-}|}(\mathcal{T}^{+-}) \hookrightarrow [\lambda(0), \lambda(\ell(s^{+-}))),\\
& && \mathrm{Lev}_{|s^{--}|}(\mathcal{T}^{--}) \hookrightarrow [\lambda(\ell(s^{+})), \lambda(\ell(s^{+}) + \ell(s^{-})))\\
&[\mathrm{c}] \quad s^{++} \in \mathcal{S}, s^{-+} \notin \mathcal{S} \ : \ && \mathrm{Lev}_{|s^{+-}|}(\mathcal{T}^{+-}) \hookrightarrow [\lambda(\ell(s^{++})), \lambda(\ell(s^{+}))),\\
& && \mathrm{Lev}_{|s^{--}|}(\mathcal{T}^{--}) \hookrightarrow [\lambda(\ell(s^{+})), \lambda(\ell(s)))\\
&[\mathrm{d}] \quad s^{++} \notin \mathcal{S}, s^{-+} \in \mathcal{S} \ : \ && \mathrm{Lev}_{|s^{+-}|}(\mathcal{T}^{+-}) \hookrightarrow [\lambda(0), \lambda(\ell(s^{+-}))),\\
& && \mathrm{Lev}_{|s^{--}|}(\mathcal{T}^{--}) \hookrightarrow [\lambda(\ell(s^{+}) + \ell(s^{-+})), \lambda(\ell(s)))
\end{aligned}$$

Schematically, where the underlines indicate isomorphisms between leaf-intervals:
Case $s^{++}, s^{-+} \in \mathcal{S}$:


$\langle s; 0\rangle$
$\langle s^{+}; 1\rangle$
$\langle s^{-}; 0\rangle$
$\langle s^{++}; 2\rangle$
$\underline{\mathcal{T}(s^{+-})}$
$\langle s^{-+}; 1\rangle$
$\underline{\mathcal{T}(s^{--})}$

Case $s^{++}, s^{-+} \notin \mathcal{S}$: $(s^{+-} = s^{--})$


$\langle s; 0\rangle$
$\langle s^{+}; 1\rangle$
$\langle s^{-}; 0\rangle$
$\underline{\mathcal{T}(s^{--})}$
$\underline{\mathcal{T}(s^{--})}$

Case $s^{++} \in \mathcal{S}, s^{-+} \notin \mathcal{S}$:


$\langle s; 0\rangle$
$\langle s^{+}; 1\rangle$
$\langle s^{-}; 0\rangle$
$\langle s^{++}; 2\rangle$
$\underline{\mathcal{T}(s^{+-})}$
$\underline{\mathcal{T}(s^{--})}$

Case $s^{++} \notin \mathcal{S}, s^{-+} \in \mathcal{S}$:


$\langle s; 0\rangle$
$\langle s^{+}; 1\rangle$
$\langle s^{-}; 0\rangle$
$\underline{\mathcal{T}(s^{+-})}$
$\langle s^{-+}; 1\rangle$
$\underline{\mathcal{T}(s^{--})}$

Using the inductive hypothesis and $s^{+-} = s^{--}$,

$$\begin{aligned}
\mathrm{f}_{s^{++}}\mathbf{M}_{[0,\ell(s^{++}))} &\cong \mathrm{f}_{s^{+-}}\mathbf{M}_{[\ell(s^{++}),\ell(s^{+}))} \\
&= \mathrm{f}_{s^{+-}}\mathbf{M}_{[\ell(s^{++}),\ell(s^{++})+\ell(s^{+-}))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{++}),\ell(s^{++})+\ell(s^{--}))}, \\
\mathrm{f}_{s^{-+}}\mathbf{M}_{[0,\ell(s^{-+}))} &\cong \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{-+}),\ell(s^{-}))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{-+}),\ell(s^{-+})+\ell(s^{--}))}, \\
\mathrm{f}_{s^{-+}}\mathbf{M}_{[\ell(s^{+}),\ell(s^{+})+\ell(s^{-+}))} &\cong \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+})+\ell(s^{-+}),\ell(s^{+})+\ell(s^{-}))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+})+\ell(s^{-+}),\ell(s))}.
\end{aligned}$$

So,

$$\begin{aligned}
\mathrm{f}_{s^{+}}\mathbf{M}_{[0,\ell(s^{+}))} &= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{++}),\ell(s^{+}))} \ \wedge\ \mathbf{M}_{[\ell(s^{++}),\ell(s^{+}))}, \\
\mathrm{f}_{s^{-}}\mathbf{M}_{[\ell(s^{+}),\ell(s))} &= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+})+\ell(s^{-+}),\ell(s))} \ \wedge\ \mathbf{M}_{[\ell(s^{+})+\ell(s^{-+}),\ell(s))}.
\end{aligned}$$

But by (a), the 'second halves' are isomorphic $\mathbf{M}_{[\ell(s^{++}),\ell(s^{+}))} \cong \mathbf{M}_{[\ell(s^{+})+\ell(s^{-+})+\ell(s))}$. In the immediately preceding equations, the respective 'first halves' of $\mathrm{f}_{s^{+}}\mathbf{M}_{[0,\ell(s^{+}))}$ and $\mathrm{f}_{s^{-}}\mathbf{M}_{[\ell(s^{+}),\ell(s))}$ are isomorphic to tails of the 'second' ones, hence also isomorphic to each other, hence so are the respective left-hand terms, as required.

Case $s^{++}, s^{-+} \notin \mathcal{S}$: In that case,

$$\begin{aligned}
\mathrm{f}_{s^{+}}\mathbf{M}_{[0,\ell(s^{+}))} &= \mathrm{f}_{s^{+-}}\mathbf{M}_{[0,\ell(s^{+-}))} \ \wedge\ \mathbf{M}_{[0,\ell(s^{+-}))} \\
\mathrm{f}_{s^{-}}\mathbf{M}_{[\ell(s^{+}),\ell(s))} &= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+}),\ell(s^{+})+\ell(s^{-}))} \ \wedge\ \mathbf{M}_{[\ell(s^{+}),\ell(s^{+})+\ell(s^{-}))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+}),\ell(s))} \ \wedge\ \mathbf{M}_{[\ell(s^{+}),\ell(s))}
\end{aligned}$$

Now by (b), $\mathbf{M}_{[0,\ell(s^{+-}))} \cong \mathbf{M}_{[\ell(s^{+}),\ell(s^{+})+\ell(s^{-}))} = \mathbf{M}_{[\ell(s^{+}),\ell(s))}$. it follows immediately that the respective first and second 'halves' are isomorphic, whence so are the respective left-hand terms.

Case $s^{++} \in \mathcal{S}, s^{-+} \notin \mathcal{S}$: Using the inductive hypothesis ans $s^{+-} = s^{--}$,

$$\begin{aligned}
\mathrm{f}_{s^{+}}\mathbf{M}_{[0,\ell(s^{+}))} &= \mathrm{f}_{s^{++}}\mathbf{M}_{[0,\ell(s^{++}))} \ \wedge\ \mathbf{M}_{[\ell(s^{++}),\ell(s^{+}))} \\
&= \mathrm{f}_{s^{+-}}\mathbf{M}_{[\ell(s^{++}),\ell(s^{+})} \ \wedge\ \mathbf{M}_{[\ell(s^{++}),\ell(s^{+}))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{++}),\ell(s^{+})} \ \wedge\ \mathbf{M}_{[\ell(s^{++}),\ell(s^{+}))} \\
\mathrm{f}_{s^{-}}\mathbf{M}_{[\ell(s^{+}),\ell(s))} &= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+}),\ell(s^{+})+\ell(s^{-}))} \ \wedge\ \mathbf{M}_{[\ell(s^{+}),\ell(s^{+})+\ell(s^{-}))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+}),\ell(s))} \ \wedge\ \mathbf{M}_{[\ell(s^{+}),\ell(s))}
\end{aligned}$$

By (c), $\mathbf{M}_{[\ell(s^{++}),\ell(s^{+})} \cong \mathbf{M}_{[\ell(s^{+}),\ell(s))}$, it follows again that the respective first and second 'halves' are isomorphic, whence so are the respective left-hand terms.

Case $s^{++} \notin \mathcal{S}, s^{-+} \in \mathcal{S}$: From $s^{+-} = s^{--}$ and the inductive hypothesis,

$$\begin{aligned}
\mathrm{f}_{s^{+}}\mathbf{M}_{[0,\ell(s^{+}))} &= \mathrm{f}_{s^{+-}}\mathbf{M}_{[0,\ell(s^{+-}))} \ \wedge\ \mathbf{M}_{[0,\ell(s^{+-}))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[0,\ell(s^{+-}))} \ \wedge\ \mathbf{M}_{[0,\ell(s^{+-}))} \\
\mathrm{f}_{s^{-}}\mathbf{M}_{[\ell(s^{+}),\ell(s))} &= \mathrm{f}_{s^{-+}}\mathbf{M}_{[\ell(s^{+}),\ell(s^{+})+\ell(s^{-+}))} \ \wedge\ \mathbf{M}_{[\ell(s^{+})+\ell(s^{-+}),\ell(s))} \\
&= \mathrm{f}_{s^{--}}\mathbf{M}_{[\ell(s^{+})+\ell(s^{-+}),\ell(s))} \ \wedge\ \mathbf{M}_{[\ell(s^{+})+\ell(s^{-+}),\ell(s))},
\end{aligned}$$

whence (d), $\mathbf{M}_{[0,\ell(s^{+-}))} \cong \mathbf{M}_{[\ell(s^+)+\ell(s^{-+},\ell(s))}$ implies that the respective segments, and hence the respective left-hand terms are isomorphic. •

As a corollary, we get

3.14 **Lemma** (Uniformity over limit lemma - sufficiency part) If $\mathbf{M}_{\ell(s)} = [M_0, \ldots, M_{\ell(s)-1}]$ satisfies the conditions of the uniformity over limit rank, then $A = A(s)[M_0, \ldots, M_{\ell(s)-1}]$ is uniform and in fact,

$$\mathrm{f}(A(s)[\mathbf{M}_{\ell(s)}]) = A(\mathrm{f}s)[\mathrm{f}_s(\mathbf{M}_{\ell(s)})].$$

Proof Base-cases: The condition is vacuous for $s = \langle 0 \rangle$. Cleary, $A(\langle 0,0 \rangle)[M_0, M_1] \in U(\omega+1)$ just if $\mathrm{f}M_0 \cong \mathrm{f}M_1$ (there are no further conditions on $[M_0, M_1]$ as $\psi^{\langle 0,0 \rangle}(0) \neq \psi^{\langle 0,0 \rangle}(1)$), $\mathrm{f}_{\langle 0,0 \rangle}[M_0, M_1] = [\mathrm{f}M_0, M_1]$, and

$$\mathrm{f}(A(\langle 0,0 \rangle)[M_0, M_1]) = \mathrm{f}(M_0 + M_1\mathrm{N}) = M + M_1\mathrm{N} = A(\langle 0,0 \rangle)[M, M_1]$$

for $M \cong \mathrm{f}M_0 \cong \mathrm{f}M_1$. For $A(\langle 0,1 \rangle)[M_0] \in U(\omega+1)$ there are no further conditions on $[M_0]$ (except the underlying that $M_0 \in \mathcal{U}(\omega)$), $\mathrm{f}_{\langle 0,1 \rangle}[[M_0] = [\mathrm{f}M_0, M_0]$, and

$$\mathrm{f}(A(\langle 0,1 \rangle)[M_0]) = \mathrm{f}(M_0\mathrm{Z}) = M + M_0\mathrm{N} = A(\langle 0,0 \rangle)[M, M_0] = A(\mathrm{f}\langle 0,1 \rangle)[M, M_0]$$

for $M \cong \mathrm{f}M_0 \cong \mathrm{f}M_1$.

Let now $s \in \mathcal{S}$ and assume the lemma is true for all $t \in \mathcal{D}(s)$.

Case $s^+ \notin \mathcal{S}$: Here,

$$\begin{aligned}
\mathrm{f}(A(s)[\mathbf{M}_{\ell(s)}]) &= \mathrm{f}(A(s^-)[\mathbf{M}_{[\ell(s^+),\ell(s))}].\mathrm{Z}) \\
&= \mathrm{f}(A(s^-)[\mathbf{M}_{\ell(s^-)}]) + A(s^-)[\mathbf{M}_{[\ell(s^+),\ell(s))}].\mathrm{N} \\
&= A(\mathrm{f}s^-)[\mathrm{f}_{s^-}(\mathbf{M}_{\ell(s^-))}] + A(s^-)[\mathbf{M}_{[\ell(s^+),\ell(s))}].\mathrm{N} \\
&= A(\mathrm{f}s^{)}[\mathrm{f}_{s^-}(\mathbf{M}_{\ell(s^-))} \ \wedge \ +[\mathbf{M}_{\ell(s^-)}].\mathrm{N} \\
&= A(\mathrm{f}s)[\mathrm{f}_s\mathbf{M}_{\ell(s)}],
\end{aligned}$$

using the inductive hypothesis in step 3, the definition of $A(t)[\mathbf{M}_{\ell(t)}]$ for $t = \mathrm{f}s$ in step 4, and the definition of $\mathrm{f}_s(\mathbf{M}_{\ell(s)})$ in step 5.

Case $s^+ \in \mathcal{S}$: Here,

$$\begin{aligned}
\mathrm{f}(A(s)[\mathbf{M}_{\ell(s)}]) &= \mathrm{f}(A(s^+)[\mathbf{M}_{\ell(s^+)}] + A(s^-)[\mathbf{M}_{[\ell(s^+),\ell(s))}].\mathrm{N}) \\
&= \begin{cases} \mathrm{f}(A(s^+)[\mathbf{M}_{\ell(s^+)}]) + A(s^-)[\mathbf{M}_{[\ell(s^+),\ell(s))}].\mathrm{N} & \text{cut-off in } A(s^+)[\mathbf{M}_{\ell(s^+)}] \\ \mathrm{f}(A(s^-)[\mathbf{M}_{\ell(s^-)}]) + A(s^-)[\mathbf{M}_{[\ell(s^+),\ell(s))}].\mathrm{N} & \text{cut-off in some } A(s^-)[\mathbf{M}_{[\ell(s^-),\ell(s)}] \times \{n\}. \end{cases}
\end{aligned}$$

By inductive hypothesis,

$$\begin{aligned}
\mathrm{f}(A(s^+)[\mathbf{M}_{\ell(s^+)}]) &= A(\mathrm{f}s^+)[\mathrm{f}_{s^+}(\mathbf{M}_{\ell(s^+)}], \\
\mathrm{f}(A(s^-)[\mathbf{M}_{[\ell(s^+),\ell(s))}]) &= A(\mathrm{f}s^-)[\mathrm{f}_{s^-}(\mathbf{M}_{[\ell(s^+),\ell(s))}].
\end{aligned}$$

But $\mathrm{f}s^+ = \mathrm{f}s^- = s^= \wedge \langle 0 \rangle$ by the tail string lemma, and $\mathrm{f}_{s^+}(\mathbf{M}_{[0,\ell(s^+))}) = \mathrm{f}_{s^-}(\mathbf{M}_{[\ell(s^+),\ell(s))})$ by the tail uniformity lemma. •

3.15 **Remark** The following description yields an equivalent definition of f_s. The base case is as in definition 3.12. For the inductive step, suppose $\mathrm{f}_t(\mathbf{M}_{\ell(t)})$ is defined for every $t \in \mathcal{D}(s)$. Let $n(s) = |s| - 1$ and $k(s) = \max\{i \leq n(s) : s(i) = i\}$

(a) Case $k(s) = n(s)$. Then $\ell(s) = \ell(s^-)$, while $\ell(\mathrm{f}s) = \ell((\mathrm{f}s)^+) + \ell((\mathrm{f}s)^-) = \ell(\mathrm{f}s^-) + \ell(s^-)$. By hypothesis, $\mathrm{f}(\mathbf{M}_{\ell(s^-)})$ is defined. So

$$\mathrm{f}_s(\mathbf{M}_{\ell(s)}) = \mathrm{f}_{s^-}(\mathbf{M}_{\ell(s^-)}) \ \wedge \ \mathbf{M}_{\ell(s)}$$

(b) Case $k(s) < n(s)$. Then $\ell(s) = \ell(s^+) + \ell(s^-)$, and $\ell(\mathrm{f}s) = \ell((\mathrm{f}s)^+) + \ell((\mathrm{f}s)^-) = \ell(\mathrm{f}s^-) + \ell(s^-)$. Let k maximal $\leq n(s)$ such that $s^{-k} \in \mathcal{S}$. By induction

$$\ell(s) = \ell(s^{-k}) + \ell(s^{-(k-1)}) + \ldots + \ell(s^-),$$

and by hypothesis, $\mathrm{f}_{s^{-i}}(\mathbf{M}_{\ell(s^{-i})})$ are defined for $i = 1, \ldots, k$ and $\mathrm{f}_{s^{-k}}(\mathbf{M}_{\ell(s^{-k})})$ is defined by case (a). So

$$\mathrm{f}_s(\mathbf{M}_{\ell(s)}) = \mathrm{f}_{s^{-k}}(\mathbf{M}_{\ell(s^{-k})}) \ \wedge \ \mathbf{M}_{[\ell(s^{-(k-1)}),\ell(s^{-(k-2)}))} \ \wedge \ \ldots \ \wedge \ \mathbf{M}_{[\ell(s^{-2}),\ell(s^{-}))}$$

[3.3] Short representations

Set $w^s = \mathrm{ran}(\psi^s)$. Given $A = A(s)[M_i]_{i<\ell^s}$, by the ω+finite uniformity lemma, there are chains $U_k \in \mathcal{U}(\omega)$, $k \in w^s$ such that

$$(\forall i \in \ell(s))[M_i = U_{\psi^s(i)}]. \tag{3.2}$$

(i.e., the set $\{U_k : k \in w^s\}$ contains just one representative (suitably tagged) under isomorphism, for each of the possibly non-isomorphic ω components.) Set $A(s)\{\langle U_k, k\rangle\}_{k\in w^s} := A(s)[M_i]_{i<\ell(s)}$ whenever (6.3) holds and refer to this as the *tagged short representation* of $A = A(s)[M_i]_{i<\ell(s)}$. Also refer to $\mathbf{U} = \{\langle U_k, k\rangle\}_{k\in w^s}$ as the *tagged short array* of that representation and to $\mathbf{U}$ as the *tagged reduction* of $\mathbf{M} = \langle M_i\rangle_{i<\ell(s)}$.

Corollary For any $s \in \mathcal{S}(n+1)$ and any $\{U_k : k \in w^s\}$, $U_k \in \mathcal{U}(\omega)$ such that

$$\mathrm{f}U_k \cong \mathrm{f}U_l \text{ for any } k, l \in w^s, \tag{3.3}$$

$A(s)\{\langle U_k, k\rangle : k \in w^s\}$ is a uniform chain of rank $\omega + n$ and every $A \in \mathcal{U}(\omega, \omega+\omega)$ has a repsentation of this form.

Whenever referring to a set $\{\langle U_k, k\rangle : k \in w^s\}$, $U_k \in \mathcal{U}(\omega)$ as a tagged short array, condition (3.3) is assumed to hold.

Example Consider $s = \langle 0, 0, 1, 0\rangle$: $\mathcal{T}(s)$, fully expanded, is



so $\psi^s = \langle 3,0,1,0,0,1,0\rangle$ and with a tagged short array $\{\langle U_0,0\rangle,\langle U_1,1\rangle,\langle U_3,3\rangle\}$, $U_0,U_1,U_3 \in \mathcal{U}(\omega)$

$$A(\langle 0,0,1,0\rangle)\{\langle U_0,0\rangle,\langle U_1,1\rangle,\langle U_3,3\rangle\} = A(\langle 0,0,1,0\rangle)[U_3,U_0,U_1,U_0,U_0,U_1,U_0]$$

If it seems that the price in tags seems high compared to the apparent gain in length-reduction, observe for example that for all $s_n = \langle 0\rangle \wedge \langle 1^{(n+1)}\rangle$, $\ell(s_n) = 2^n$, while $|w^{s_n}| = 1$. This will be become relevant in situations where sequences s_n arise naturally (such as when approximating $A \in \mathcal{U}(\omega+\omega)$).

Moreover, the tagging can be made dispensible in turn, by defining a canonical ordering. There is an obvious association between the nodes of form $\langle t,k\rangle$ of $\mathcal{T}^s$ and components of $A = A[M_i]_{i<\ell(s)}$ of form $A(t)[M_i]_{i\in\mathcal{J}}$. (Here $\mathcal{J}$ is a sub-interval of $[0,\ell(s))$, i.e., $\langle M_i\rangle_{i\in\mathcal{J}}$ is the set of building-blocks $\langle M'_i\rangle_{i<\ell(t)}$ used for that component). Namely, starting with A at the root and proceeding dow the tree, associate the 'plus'/'minus' subcomponent(s) arived at to the 'plus'/'minus' extension(s) of the node arrived at. Next, extend that association to the tagged short arrays of these components. It then turns out that along any branch $\sigma_0,\ldots,\sigma_j,\ldots,\sigma_n = \langle\langle 0\rangle,k_0\rangle,\ldots,\langle t,k\rangle,\ldots,\langle s,0\rangle$ of $\mathcal{T}^s$, proceeding from the leaves to the root, the arrays $\mathbf{U}_{t,k}$ associated to the nodes $\langle t,k\rangle$ in that way form a chain whose members increase in size by at most 1 over their predecessors. That generates an ordering of the elements in the union over such a chain (e.g., the order of 'aquisition'). But that union is always $\mathbf{U} = \mathbf{U}_{s,0}$, the array for A itself (that gets associated with the top node). For a 'canonical' ordering it then suffices that we can always single out a unique such branch, say by chosing the right-most one (i.e., following consistently the 'minus' extensions as considered from the root).

Example Here is the 'associated' tree of tagged short arrays for $t \in \mathcal{D}(s)$ for $s = \langle 0,0,1,0\rangle$ of the preceding example – note that the tags on the chains in $\mathbf{U}_{t,k}$ now represent the weights from the weight-sequence of sequence t of the corresponding node $\langle t,k\rangle$, not the weight k of the node:



The 'canonical' ordering of the chains in the tagged short array would thus be $\langle U_0,U_1,U_3\rangle$, corresponding to the right-most branch, starting at leaf $i = 6$. (E.g., using the left-most branch, starting at leaf $i = 0$ would result in $\langle U_3,U_0,U_1\rangle$ and starting at leaf $i = 1$ would result in $\langle U_0,U_3,U_1\rangle$. Etc.) The canonically ordered array shall be the desired *short array*. Since the size of $\mathbf{U}_{t,k}$ is equal to that of w^t, some facts are required about how $|w^s|$ relates to $|w^{s^\pm}|$ in order to implement this.

(3.16) **Definition** (Leaf-intervals of components and component-node association) Suppose $A = A[M_i]_{i<\ell(s)} \in \mathcal{U}(\omega+n)$, where $n = |s|-1$.

- (i) Set $\mathcal{L}_{s,0} = [0,\ell(s))$, the leaf interval of A. Associate A to $\langle s,0\rangle$.
- (ii) Let $\langle t,k\rangle \in \mathcal{T}^s$ and supose $\mathcal{L}_{t,k} \subseteq \mathcal{L}_{s,0}$ is the leaf-interval (of length $\ell(t)$) of $A_{t,k} = A(t)[M_i]_{i\in\mathcal{L}_{t,k}}$ and suppose $A_{t,k}$ is associated to that node. Let $m = |t|-1$.

(a) If $t(m) = m$ then $\langle t^-, 0\rangle$ is the only extension of $\langle t, k\rangle$ $\ell(t) = \ell(t^-)$, and $A(t)[M_i]_{i\in\mathcal{L}_{t,k}} = A(t^-)[M_i]_{i\in\mathcal{L}_{t,k}}$.Z, so set $\mathcal{L}^-_{t,k} := \mathcal{L}_{t^-,0} := \mathcal{L}_{t,k}$ and associate $A(t^-)[M_i]_{i\in\mathcal{L}_{t,k}}$ to $\langle t^-, 0\rangle$,

(b) Else ($t(m) < m$), there are two extensions $\langle t^+, k+1\rangle$ and $\langle t^-, 0\rangle$ and $\ell(t) = \ell(t^+) + \ell(t^-)$. Set $\mathcal{L}^+_{t,k} := \mathcal{L}_{t^+,k+1} :=$ the initial segment of $\mathcal{L}_{t,k}$ of length $\ell(t^+)$, and $\mathcal{L}^-_{t,k} := \mathcal{L}_{t^-,0} :=$ the final segment of $\mathcal{L}_{t,k}$ of length $\ell(t^-)$. Then $A(t)[M_i]_{i\in\mathcal{L}_{t,k}} = A(t^+)[M_i]_{i\in\mathcal{L}^+_{t,k}} + A(t^-)[M_i]_{i\in\mathcal{L}^-_{t,k}}$.N, so associate $A(t^+)[M_i]_{i\in\mathcal{L}^+_{t,k}}$ to $\langle t^+, k+1\rangle$ and $A(t^-)[M_i]_{i\in\mathcal{L}^-_{t,k}}$ to $\langle t^-, 0\rangle$.

Let $\mathbf{M} = \langle M_i\rangle_{i<\ell(s)} \in {}^{\ell(s)}\mathcal{U}(\omega)$, be the array in the representation of some $A \in \mathcal{U}(\omega)$. Implicit in the definition of A as $A(s)[\mathbf{M}]$, is a mapping

$$[0, \ell(s)) \to \begin{cases} [0, \ell(s))^- := [0, \ell(s^-)) = [0, \ell(s)) & \text{if } s(n) = n \\ [0, \ell(s))^+, [0, \ell(s))^- := [0, \ell(s^+)), [\ell(s^+), \ell(s)) & \text{if } s(n) < n \end{cases}$$

and hence

$$\mathbf{M} \to \begin{cases} \mathbf{M}^- := \langle M_i\rangle_{i<\ell(s^-)} = \mathbf{M} & \text{if } s(n) = n \\ \mathbf{M}^+, \mathbf{M}^- := \langle M_i\rangle_{i<\ell(s)}, \langle M_i\rangle_{\ell(s^+)\le i<\ell(s)} & \text{if } s(n) < n. \end{cases}$$

Define the corresponding operations $\mathbf{U} \to \mathbf{U}^\pm$ on short arrays so that $\mathbf{U}^\pm$ is the reduction of $\mathbf{M}^\pm$; again, these will depend on s, so we effectively have an operation $\mathbf{U} \to \mathbf{U}^\pm_s$. (In case of the arrays $\mathbf{M} = \langle M_i : i < \ell(s)\rangle$ the dependency on s is implicit in the notation. The operation is assumed to be defined if $\mathbf{U}$ is of form $\{U_k\}_{k\in w^s}$ for chains $U_k \in \mathcal{U}(\omega)$). Thus:

(3.17) **Definition** (Decomposition of tagged short arrays) Where $\mathbf{U} = \{\langle U_k, k\rangle\}_{k\in w^s}$, set and $n = |s| - 1$,

$$\mathbf{U}^-_s = \{\langle U_k, k\rangle\}_{k\in w^{s^-}}$$
$$\mathbf{U}^+_s = \begin{cases} \{\langle U_n, n-1\rangle\} \cup \{\langle U_k, k\rangle\}_{k\in w^{s^+}\setminus\{n-1\}} & \text{if } s(n) = 0 \\ \{\langle U_k, k\rangle\}_{k\in w^{s^+}} & \text{if } 0 < s(n) < n \\ \text{undefined} & \text{if } s(n) = n. \end{cases}$$

(The change from $\langle U_n, n\rangle \in \mathbf{U}$ to $\langle U_n, n-1\rangle \in \mathbf{U}^+_s$ in case $s(n) = 0$ is due to the fact that $\psi^s(0) = \psi^{s^+}(0) + 1 = n$ just if $s(n) = 0$ for $n = |s| - 1$, by the second weight induction corollary). This allows us to define the chain constructor in terms of the short arrays and obtain the 'transfinite' analogue of Pastjin's uniform chain constructor:

(3.18) **Corollary** (Plus finite rank Pastjin unifom chain constructor — short version) Where $\mathbf{U} = \{\langle U_k, k\rangle\}_{k\in w^s} \subseteq \mathcal{U}(\omega) \times w^s$ is a tagged short array,

$$A(s)\{\mathbf{U}\} = \begin{cases} A(s^-)\{\mathbf{U}^-_s\}.\mathrm{Z} & \text{if } s(n) = n \text{ for } n = |s| - 1 \\ A(s^+)\{\mathbf{U}^+_s\} + A(s^-)\{\mathbf{U}^-_s\}.N & \text{if } s(n) < n \text{ for } n = |s| - 1 \end{cases}$$

Example (Comparing the short arrays $U_0, \ldots, U_{\zeta^s}$ with the full array, $M_0, \ldots, M_{\ell-1}$.)

If $s = \langle 0,0,0\rangle$, then $\tau_1 \in \{0,1\}$ and $\nu^s = \{0\}$ if $\tau_1 = 1$ and $\nu^s = \{2\}$ if $\tau_0 = 0$, so

$$
\begin{aligned}
&A(0,0,0)\{U_0,U_1,U_2\}) \\
&= A(0,0,0)[M_0,M_1,M_2,M_3] \\
&= A(0,0)[M_0,M_1] + A(0,0)[M_2,M_3].\mathrm{N} \\
&= \begin{cases}
A(0,0)[U_0,U_1] + A(0,0)[U_2,U_1].\mathrm{N} & \text{if } \tau_0 = 0 \text{ and } \tau_1 = 0 \ (\nu^{s^+} = \{1\} \text{ and } \nu^s = \{2\}) \\
A(0,0)[U_2,U_1] + A(0,0)[U_0,U_1].\mathrm{N} & \text{if } \tau_0 = 0 \text{ and } \tau_1 = 1 \ (\nu^{s^-} = \{1\} \text{ and } \nu^s = \{0\}) \\
A(0,0)[U_1,U_0] + A(0,0)[U_2,U_0].\mathrm{N} & \text{if } \tau_0 = 1 \text{ and } \tau_1 = 0 \ (\nu^{s^+} = \{0\} \text{ and } \nu^s = \{2\}) \\
A(0,0)[U_2,U_0] + A(0,0)[U_1,U_0].\mathrm{N} & \text{if } \tau_0 = 1 \text{ and } \tau_1 = 1 \ (\nu^{s^-} = \{0\} \text{ and } \nu^s = \{0\})
\end{cases} \\
&= \begin{cases}
A(0,0)\{U_0,U_1\} + A(0,0;\{U_2,U_1\}.\mathrm{N} \\
A(0,0)\{U_2,U_1\} + A(0,0)\{U_0,U_1\}.\mathrm{N} \\
A(0,0)\{U_1,U_0\} + A(0,0)\{U_2,U_0\}.\mathrm{N} \\
A(0,0)\{U_2,U_0\} + A(0,0)\{U_1,U_0\}.\mathrm{N}
\end{cases} \\
&= \begin{cases}
U_0 + U_1.\mathrm{N} + (U_2 + U_1.\mathrm{N}).\mathrm{N} \\
U_2 + U_1.\mathrm{N} + (U_0 + U_1.\mathrm{N}).\mathrm{N} \\
U_1 + U_0.\mathrm{N} + (U_2 + U_0.\mathrm{N}).\mathrm{N} \\
U_2 + U_0.\mathrm{N} + (U_1 + U_0.\mathrm{N}).\mathrm{N}
\end{cases}
\end{aligned}
$$

by following the 'line up' of the U_i along the association tree for $\langle 0,0,0\rangle$ (i.e., depending on where we select the 'base-point'. E.g., if $\tau_0 = 1$ and $\tau_1 = 0$ we start with the second leaf $\lambda^s(1) = \langle 0\rangle$ of $\mathcal{T}^s$ picking up U_0, proceed to its parent node $\langle 0,0\rangle$ on the left branch of s picking up U_1 and arriving at the top-node s, picking up U_2.)

[4] Products of uniform chains

The preceding section takes uniform chain constructions forward by only an 'infinitesimal' step, to ranks finite over a limit. The crucial step will be to complete the limit steps. The following discussion considers simple types of uniform chains of 'limit' rank, but really serves to illustrate the fact that such simple 'algebraic' type constructions (involving sums/products) won't get us anywhere near exploring the 'limit' explosion in the variety of uniform chains.

[4.1] Atomic chain products

If I is a chain (linear order) and for each $i \in I$, A_i is a chain, then the expression $A = \sum_{i \in I} A_i$ defines a chain in the usual fashion. Sums easily allow to express finite products. The (anti-lexicographically ordered cartesian) product of two chains, $A = B_1 B_2$ is expressed by $A = \sum_{b_2 \in B_2} B_1$ and this definition iterates through all finite products. Some infinite products can be equally easily expressed by infinite sums (of finite products) such as e.g.,

$$\mathrm{N}^{\mathrm{N}} = \prod_{i \in \mathrm{N}} \mathrm{N} = \sum_{i \in \mathrm{N}} \mathrm{N}^i.$$

More revealing is another example,

$$\mathrm{Z}^{\mathrm{N}} = \prod_{i \in \mathrm{N}} \mathrm{Z} = \sum_{-i \in \mathrm{N}^*} \mathrm{Z}^i + \sum_{i \in \mathrm{N}} \mathrm{Z}^i,$$

which suggests how to express infinite products of chains. Let $H_0 = 1$ and $H_i \in \{\mathrm{N}, \mathrm{Z}\}$ for $i \geq 1$. What should $\prod_{i \in \mathrm{N}} H_i$ be? Define $P_0 = 0$, and for $i > 0$,

$P_i = \prod_{k<i} H_k$

$P_{-i} = P_i$ if $H_i = \mathrm{Z}$,

$P_{-i} = 0$ otherwise (if $H_i = \mathrm{N}$).

Take $A = \prod_{i \in \mathrm{N}} H_i$ to be the chain expressed by $A = \sum_{-i \in \mathrm{N}^*} P_{-i}\mathrm{N}^* + \sum_{i \in \mathrm{N}} P_i \mathrm{N}$ or, compactly $A = \sum_{i \in \mathrm{Z}} P_i \mathrm{N}^{\mathrm{sgn}(i)}$, on the understanding that $\mathrm{N}^+ = \mathrm{N}$ $\mathrm{N}^- = \mathrm{N}^*$. This definition quickly reproduces the two special cases mentioned; in N^{N} all P_{-i} vanish and $P_{i+1} = \mathrm{N}^i$; in Z^{N} for all i, $P_{-i-1} = P_{i+1} = \mathrm{Z}^i$. It also applies in the case of finite products:

(4.1) **Lemma** $P_n = \sum_{i \in (-n,n)} P_i \mathrm{N}^{\mathrm{sgn}(i)}$ for $n > 1$.

Proof Observe that $P_0 = 0$, $P_1 = H_0 = 1$, $P_2 = H_0 H_1 = H_1 \in \{\mathrm{N}, \mathrm{Z}\}$, and

$$\sum_{i \in (-2,2)} P_i \mathrm{N}^{\mathrm{sgn}(i)} = P_{-1}\mathrm{N}^* + 0 + P_1 \mathrm{N},$$

which is N if $H_1 = \mathrm{N}$ and Z if $H_1 = \mathrm{Z}$, hence P_2. Assuming the statement of the lemma true for some $n > 1$, we get

$$P_{n+1} = \prod_{i \leq n} H_i =_{\mathrm{dfn}} (\prod_{i<n} H_i) H_n = P_n H_n \in \{P_n \mathrm{N}, P_n \mathrm{Z}\},$$

while

$$\begin{aligned}\sum_{i\in[-n,n]} P_i \mathrm{N}^{\mathrm{sgn}(i)} & \\ &= P_{-n}\mathrm{N}^* + \sum_{i\in(-n,n)} P_i\mathrm{N}^{\mathrm{sgn}(i)} + P_n\mathrm{N} \\ &=_{\mathrm{ind}} P_{-n}\mathrm{N}^* + P_n + P_n\mathrm{N} \\ &= P_{-n}\mathrm{N}^* + P_n\mathrm{N},\end{aligned}$$

which is $P_n\mathrm{N}$ if $H_n = \mathrm{N}$ and $P_n\mathrm{Z}$ if $H_n = \mathrm{Z}$, hence equal to P_{n+1}. •

Thus, for each $n > 1$,

$$\begin{aligned}\sum_{i\in\mathrm{Z}} P_i\mathrm{N}^{\mathrm{sgn}(i)} &= \sum_{-i\in\mathrm{N}^*} P_{-i}\mathrm{N}^* + \sum_{i\in\mathrm{N}} P_i\mathrm{N} \\ &= \sum_{-i\leq -n} P_{-i}\mathrm{N}^* + P_n + \sum_{i\geq n} P_i\mathrm{N},\end{aligned}$$

and in this representation the middle summand is called the 'central' P_n.

(4.2) **Lemma** $A = \prod_{i\in\mathrm{N}} H_i$ is (right) uniform, and

$$\mathrm{f}A = \sum_{i\in\mathrm{N}} P_i\mathrm{N}.$$

Proof By the observation on tails of finite products,

$$\mathrm{f}P_n = \mathrm{f}\prod_{i<n} H_i = \sum_{i<n}(\prod_{k<i} H_i)\mathrm{f}H_i = \sum_{i<n}(\prod_{k<i} H_i)\mathrm{N} = \sum_{i<n} P_i\mathrm{N}.$$

Let $x \in A$. By (2.3) chose $n > 1$ such that x is in the 'central' P_n. Thus,

$$\mathrm{f}_x A = \mathrm{f}P_n + \sum_{i\geq n} P_i\mathrm{N} = \sum_{i\in\mathrm{N}} P_i\mathrm{N}$$

as claimed. Alternatively, let $P_n\mathrm{N}^{\mathrm{sgn}(n)}, n \in \mathrm{Z}$ be the summand containing x and set $x = (x', x'')$ in the cartesian representation of $P_n\mathrm{N}^{\mathrm{sgn}(n)}$. If $n \geq 0$,

$$\mathrm{f}_x A = \mathrm{f}_x P_n\mathrm{N} + \sum_{i>n} P_i\mathrm{N} = \mathrm{f}_{x'}P_n + P_n\mathrm{f}_{x''}\mathrm{N} + \sum_{i>n} P_i\mathrm{N} = \mathrm{f}P_n + P_n\mathrm{N} + \sum_{i>n} P_i\mathrm{N} = \sum_{i\in\mathrm{N}} P_i\mathrm{N}.$$

If $n = -m < 0$ (and consequently $P_{-m} = P_m$),

$$\begin{aligned}\mathrm{f}_x A &= \mathrm{f}_x P_{-m}\mathrm{N}^* + \sum_{i\in(-m,m)} P_i\mathrm{N}^{\mathrm{sgn}(i)} + \sum_{i\geq m} P_i\mathrm{N} \\ &= \mathrm{f}_{x'}P_m + P_{-m}\mathrm{f}_{x''}\mathrm{N}^* + P_m + \sum_{i\geq m} P_i\mathrm{N} \\ &= \mathrm{f}P_m + P_m \times \mathrm{fin} + \sum_{i>m} P_i\mathrm{N} \\ &= \sum_{i\in\mathrm{N}} P_i\mathrm{N},\end{aligned}$$

where $P_m \times \mathrm{fin}$ indicates 'finitely many copies' of P_m. •

[4.2] Infinite products of uniform chains

Given chains $A_i, i \in \mathrm{N}$, define $A = \prod_{i\in\mathrm{N}}\langle A_i, x_i\rangle$ by fixing a 'zero' $x_i \in A_i$ and setting $A = \sum_{k\in\mathrm{N}^*} P_k \mathrm{j}_{x_i} A_k + \sum_{k\in\mathrm{N}} P_k \mathrm{f}_{x_i} A_k$, where $P_k \prod_{i<k} A_i$. In general this 'product' depends on the choice of 'zeros'. In case of finite products it doesn't, e.g. (using right distributivity of ordered cartesian products),

$$P_2 = A_1A_2 = A_1(\mathrm{j}_{x_2}A_2 + \mathrm{f}_{x_2}A_2) = A_1\mathrm{j}_{x_2}A_2 + A_1\mathrm{f}_{x_2}A_2,$$

and using the 'telescope efect' $\mathrm{f}A = 1 + \mathrm{f}A$,

$$\begin{aligned} P_3 &= A_1A_2A_3 \\ &= A_1A_2(\mathrm{j}_{x_3}A_3 + 1 + \mathrm{f}_{x_3}A_3) \\ &= A_1A_2\mathrm{j}_{x_3}A_3 + A_1A_2 + A_1A_2\mathrm{f}_{x_3}A_3) \\ &= A_1A_2\mathrm{j}_{x_3}A_3 + A_1(\mathrm{j}_{x_2}A_2 + \mathrm{f}_{x_2}A_2) + A_1A_2\mathrm{f}_{x_3}A_3) \\ &= A_1A_2\mathrm{j}_{x_3}A_3 + A_1\mathrm{j}_{x_2}A_2 + A_1\mathrm{f}_{x_2}A_2 + A_1A_2\mathrm{f}_{x_3}A_3), \end{aligned}$$

etc, for any, hence independently of the, choice of $x_i \in A_i$. Thus, assuming for some arbitrary n, $P_n = \prod_{i<n} A_i = \sum_{-n<-k<0} P_k\mathrm{j}_{x_i}A_k + \sum_{0\le k<n} P_k\mathrm{f}_{x_i}A_k$, it follows that

$$\begin{aligned} P_{n+1} &= \prod_{i\le n} A_i \\ &= P_nA_n \\ &= P_n(\mathrm{j}_{x_n}A_n + 1 + \mathrm{f}_{x_n}A_n) \\ &= P_n\mathrm{j}_{x_n}A_n + P_n + P_n\mathrm{f}_{x_n}A_n) \\ &=_{\mathrm{ind}} P_n\mathrm{j}_{x_n}A_n + \sum_{-n<-k<0} P_k\mathrm{j}_{x_i}A_k + \sum_{0\le k<n} P_k\mathrm{f}_{x_i}A_k + P_n\mathrm{f}_{x_n}A_n) \\ &= \sum_{-n\le -k<0} P_k\mathrm{j}_{x_i}A_k + \sum_{0\le k\le n} P_k\mathrm{f}_{x_i}A_k, \end{aligned}$$

for any choice of $x_n \in A_n$. In the case of uniform chains A_i there is a 'canonical' choice of zeros:

(4.3) **Lemma** For every $A \in \mathcal{U}(<\omega)$ there are $x \in A$ and $B_i \in \mathcal{U}(i) \cup \{0\}$ such that

$$\mathrm{j}_xA = \sum_{i<\mathrm{rk}(A)} B_i\mathrm{N}^*.$$

Proof In case $\mathrm{rk}(A) = 1$ ($A \in \{\mathrm{N}, \mathrm{Z}\}$) the 'ordinary' zeros are candidates for x (in case of $A = \mathrm{Z}$, any element will do). Then $\mathrm{j}_xA \in \{0, \mathrm{N}^* = 1\dot{\mathrm{N}}^*\}$ satisfies the conclusion with $B_0 \in \{0, 1\}$. Assume the statement holds for all $A \in \mathcal{U}(<n)$. Given $A \in \mathcal{U}(n)$ fix $s \in \mathcal{S}(n)$ such that $A = A(s)$.

If $s(n) = n$, $A(s) = A(s^-)\mathrm{Z}$. Set $A' = A(s^-)$. By hypothesis there are $x' \in A'$ and $B'_i \in \mathcal{U}(i) \cup \{0\}$ for $i < n$ such that $\mathrm{j}_{x'}A' = \sum_{i<n} B'_i\mathrm{N}^*$. Let $x'' \in \mathrm{Z}$ (e.g., $x'' = 0$) and set $x = (x', x'') \in A'\mathrm{Z}$. Then

$$\begin{aligned} \mathrm{j}_xA &= \mathrm{j}_{(x',x''}A'\mathrm{Z} \\ &= A'\mathrm{j}_{x''}\mathrm{Z} + \mathrm{j}_{x'}A' \\ &= A'\mathrm{N}^* + \sum_{i<n} B'_i\mathrm{N}^* \\ &= \sum_{i\le n} B_i\mathrm{N}^* \end{aligned}$$

upon setting $B_i = B'_i$ for $i < n$ and $B_n = A'$.

If $s(n) < n$, $A(s) = A(s^+) + A(s^-)\mathrm{N}$. Set $A' = A(s^+)$ but now let $x' \in A'$ and $B'_i \in \mathcal{U}(i) \cup \{0\}$ for $i < n$ such that $\mathrm{j}_{x'}A' = \sum_{i<n} B'_i\mathrm{N}^*$. Then

$$\mathrm{j}_x A = \mathrm{j}_x A' = \sum_{i<n} B'_i \mathrm{N}^* = \sum_{i \le n} B_i \mathrm{N}^*,$$

upon setting $B_i = B'_i$ for $i < n$ and $B_n = 0$. •

This also shows that in the representation of $\mathrm{j}_x A$ we have

$$B_i = \begin{cases} 0 & \text{if } s(i) < i \\ A(s\lceil i) & \text{if } s(i) = i \end{cases}$$

for the $s \in \mathcal{S}(\mathrm{rk}(A))$ such that $A = A(s)$, so that the representing B_i are independent of the choice of x (subject to the condition that they allow such a representation).

[4.3] Infinite uniform chain products

The infinite uniform chain product $A = \prod_{i \in \mathrm{N}} H_i$ is defined to be

$$A = \sum_{-i \in \mathrm{N}^*} P_i \mathrm{j}_{x_i} H_i + \sum_{i \in \mathrm{N}} P_i \mathrm{f}_{x_i} H_i$$

(with $P_i = \prod_{k<i} H_k$ as before) for a canonical choice of $x_i \in H_i$. Namely, then,

$$A = \sum_{-i \in \mathrm{N}^*} P_i K_i + \sum_{i \in \mathrm{N}} P_i \mathrm{f} H_i$$

where the chains $K_i = \mathrm{j}_{x_i} H_i = \sum_{k < \mathrm{rk}(H_i)} B_k^{(i)} \mathrm{N}^*$ are independent of the choice of the x_i, viz,

$$B_k^{(i)} = \begin{cases} 0 & \text{if } s_i(k) < k \\ A(s_i\lceil k) & \text{if } s_i(k) = k \end{cases}$$

for the $s_i \in \mathcal{S}(\mathrm{rk}(()H_i)$ such that $H_i = A(s_i)$.

(4.4) **Corollary**

$$\prod_{i \in \mathrm{N}} H_i = \sum_{n \in \mathrm{Z}} P_n K_n$$

where for $s_i \in \mathcal{S}(\mathrm{rk}(H_i)))$ such that $H_i = A(s_i)$,

$$P_{-i} = P_i = \prod_{k<i} H_i,\ K_{-i} = \sum_{k<i, s_i(k)=k} A(s_i\lceil k)\mathrm{N},\ K_i = \sum_{k<i} A(s_i\lceil k)\mathrm{N}.$$

(4.5) **Lemma** (ω-product lemma) Infinite uniform chain products of 'length' N are uniform. In fact,

$$\mathrm{f} \prod_i H_i = \sum_i (\prod_{k<i} H_k) \mathrm{f} H_i.$$

Proof Choose a representation

$$A = \prod_i H_i = \sum_{-i \in \mathrm{N}^*} P_i K_i + \sum_{i \in \mathrm{N}} P_i \mathrm{f} H_i$$

where P_i, K_i are as before. For $x \in A$, $x = (x', x'') = (\pm n, x'')$ for $x' \in \mathrm{Z}$ and $x'' \in P_n K_n$ if $x' = -n$ or $x'' \in P_n \mathrm{f} H_n$ if $x' = n$.

If $z = n \geq 0$, since all terms involved in the right summand are themselves uniform chains,

$$\begin{aligned}
\mathrm{f}_x A &= \mathrm{f}(P_n \mathrm{f} H_n) + \sum_{i>n} P_i \mathrm{f} H_i \\
&= \mathrm{f} P_n + P_n \mathrm{f}^2 H_n + \sum_{i>n} P_i \mathrm{f} H_i \\
&= \sum_{i<n} P_i \mathrm{f} H_i + P_n \mathrm{f} H_n \sum_{i>n} P_i \mathrm{f} H_i \\
&= \sum_{i \in \mathrm{N}} P_i \mathrm{f} H_i.
\end{aligned}$$

Suppose $z = -n < 0$. Then (with canonical zeros $x_i \in H_i$)

$$\begin{aligned}
\mathrm{f}_x A &= \mathrm{f}_{x''} P_n K_n + \sum_{-n<-k<0} P_i K_i + \sum_{i \in \mathrm{N}} P_i \mathrm{f} H_i \\
&= \mathrm{f}_{x''} P_n K_n + \sum_{-n<-i<0} P_i \mathrm{j}_{x_i} H_i + \sum_{0 \leq i<n} P_i \mathrm{f} H_i + \sum_{i \geq n} P_i \mathrm{f} H_i \\
&= \mathrm{f}_{x''} P_n K_n + P_n + \sum_{i \geq n} P_i \mathrm{f} H_i \\
&= \mathrm{f}_{x''} P_n K_n + P_n \mathrm{f} H_n + \sum_{i>n} P_i \mathrm{f} H_i
\end{aligned}$$

Fix $p \in P_n, k \in K_n = \mathrm{j}_{x_n} H_n$ such that $x'' = (p, k)$, so that

$$\mathrm{f}_{x''} P_n K_n = \mathrm{f}_p P_n + P_n \mathrm{f}_k K_n = \sum_{i<n} P_i \mathrm{f} H_i + P_n \mathrm{f}_k \mathrm{j}_{x_n} H_n,$$

so

$$\begin{aligned}
\mathrm{f}_x A &= \sum_{i<n} P_i \mathrm{f} H_i + P_n \mathrm{f}_k \mathrm{j}_{x_n} H_n + P_n \mathrm{f} H_n + \sum_{i>n} P_i \mathrm{f} H_i \\
&= \sum_{i<n} P_i \mathrm{f} H_i + P_n(\mathrm{f}_k \mathrm{j}_{x_n} H_n + \mathrm{f}_{x_n} H_n) + \sum_{i>n} P_i \mathrm{f} H_i \\
&= \sum_{i<n} P_i \mathrm{f} H_i + \sum_{i \geq n} P_i \mathrm{f} H_i \\
&= \sum_{i \in \mathrm{N}} P_i \mathrm{f} H_i,
\end{aligned}$$

using right distributivity and the observation that for any uniform chain X and any $a < b \in X$,

$$\mathrm{f}_a \mathrm{j}_b X + \mathrm{f}_b X = \mathrm{f}_a X = \mathrm{f}_b X. \bullet$$

[5] Chains of first limit rank

We now resume the thread left off in section 2 and proceed to characterise (uniform) chains of rank ω.

[5.1] Chains of rank ω

Approximating $A \in \mathcal{U}(\omega)$ requires consideration of entire 'inclusion' chains of sub-components, essentially over the 'length' of the approximation given by the rank of A. I.e., we need to incorporate into any 'limit rank uniform chain constructor' the information that each such A is the union of the 'chain of chains' $\{[x]^A_n\}_{n\in\omega}$ (for any fixed $x \in A$), and hence of an 'embedding' chain of chains $\{A(s_n)\}_{n\in\omega}$ where $\{s_n\}_{n\in\omega}$ is a suitable sequence of $\mathcal{S}$ strings and where the embedding $A(s_n) \hookrightarrow A(s_{n+1})$ is the 'canonical' embedding. To state the obvious fact:

(5.1) **Lemma** (Copy lemma) For $A \in \mathcal{U}(\omega)$, $x \in A$ and $n \in \omega$,

$$[x]^A_{n+1} \cong \begin{cases} [x]^A_n + B_n.\mathrm{N} & \text{if } [x]^A_{n+1}/n = \mathrm{N} \text{ and } [x]^A_n \text{ has no predecessors in } [x]^A_{n+1} \\ B_n + [x]^A_n.\mathrm{N} & \text{if } [x]^A_{n+1}/n = \mathrm{N} \text{ and } [x]^A_n \text{ does have predecessors in } [x]^A_{n+1} \\ [x]^A_n.\mathrm{Z} & \text{if } [x]^A_{n+1}/n = \mathrm{Z} \end{cases}$$

where $B_n \in \mathcal{U}(n)$ such that $\mathrm{f}B_n \cong \mathrm{f}[x]^A_n$ and where by 'predecessor' in $[x]^A_{n+1}$ we mean 'n-component predecessor'.

We shall now turn this observation into a characterisation of $A \in \mathcal{U}(\omega)$.

(5.2) **Definition**

(a) Call a sequence $\{s_n\}_n$ of strings $s_n \in \mathcal{S}(n+1)$ *realised by* $\langle A, x\rangle$, where $x \in A \in \mathcal{U}(\omega)$, if $[x]^A_n = A(s_n)$ for all n. Call $\{s_n\}_n$ as above *realised by* $A \in \mathcal{U}(\omega)$ if it is realised by some $\langle A.x\rangle$, $x \in A$, and *realised in* $\mathcal{U}(\omega)$ if it's realised by some $A \in \mathcal{U}(\omega)$.

(b) Define the *embedding sequence* $\pi \in {}^{\omega}(\omega+1)$ of $\langle A, x\rangle$ by

$$\pi(n) = \pi^{A,x}(n) = \begin{cases} i \in \omega & \text{if } [x]^A_{n+1}/n \cong \mathrm{N} \text{ and } [x]^A_n \text{ has } i \text{ predecessors in } [x]^A_{n+1}, \\ \omega & \text{else (i.e., if } [x]^A_{n+1}/n \cong \mathrm{Z}). \end{cases}$$

(c) Define the *type sequence* $\tau \in {}^{\omega}\{0, 1, \omega\}$ of a sequence $\pi \in {}^{\omega}(\omega+1)$ by

$$\tau(n) = \begin{cases} \pi(n) & \text{if } \pi(n) \in \{0, \omega\} \\ 1 & \text{if } \pi(n) \in (0, \omega). \end{cases}$$

Define the type sequence $\tau^{A,x}(n)$ of $\langle A, x\rangle$ as the type sequence of $\pi^{A,x}(n)$.

In the following, write π_n, τ_n for $\pi(n), \tau(n)$.

Thus every $A \in \mathcal{U}(\omega)$ is determined by a pair $\langle\{s_n\}_n, \pi\rangle$ where $\{s_n\}_n$ is a sequence realised by $\langle A, x\rangle$ for some s in A, and where π is the embedding sequence for that pair $\langle A, x\rangle$. Moreover, any two pairs $\langle\{s_n\}_n, \pi\rangle$, $\langle\{s'_n\}_n, \pi'\rangle$ determining A in that fashion agree on a final segment: If the corresponding elements of A are x and x', then $s_n = s'_n$ and $\pi_n = \pi'_n$ for all $n \geq m$ such that $x\rho_m x'$.

(5.3) **Lemma** (String copy lemma) Let $\{s_n\}_n$, $s_n \in \mathcal{S}(n+1)$ be realised by $\langle A, x\rangle$, i.e., such that $[x]^A_n = A(s_n), n \in \omega$. Then,

$$s_n = \begin{cases} s^+_{n+1} & \text{if } \pi_n = 0 \text{ iff } \tau_n = 0, \\ s^-_{n+1} & \text{if } \pi_n > 0 \text{ iff } \tau_n > 0, \end{cases} \tag{5.1}$$

and in either case $s^-_n = s^=_{n+1}$.

Proof $s_n = s^+_{n+1}$ just if the first of the above cases for $[x]^A_{n+1}$ holds, else $s_n = s^-_{n+1}$; the remainder is the string lemma. •

(5.4) **Definition and Lemma** (Aligned string sequences) Call a sequence of strings $\{s_n\}_n$ *aligned* with π (resp., τ), if condition (5.1) holds, and simply *aligned*, if $s_n \in \{s^\pm_{n+1}\}$ for all n. For such an aligned sequence define $a_n = a^{A,x}_n$, $b_n = b^{A,x}_n$, $n \in \omega$ such that for all $n > 0$,

$$s_n = s^=_n \wedge \langle a_n, b_n\rangle.$$

(We let a_0 undefined and set $b_0 = 0$, so $s_0 = \langle b_0\rangle$.) Then,

$$b_n = \begin{cases} b_{n+1} & \text{if } s_n = s^+_{n+1} \text{ iff } \pi_n = 0 \text{ iff } \tau_n = 0, \\ a_{n+1} & \text{if } s_n = s^-_{n+1} \text{ iff } \pi_n > 0 \text{ iff } \tau_n > 0. \end{cases} \tag{5.2}$$

Also, if we set $A_n = A(s_n)$, then $\{A_n\}_n$ the sequence $\{A_n\}_n$ satisfies

$$A_{n+1} \cong \begin{cases} A_n\mathrm{Z} & \text{if } \pi_n = \omega, \\ A_n + B_n\mathrm{N}, & \text{for } B_n \in \mathcal{U}(n) \text{ with } \mathrm{f}B_n = \mathrm{f}A_n, \text{ if } \pi_n = 0, \\ B_n + A_n\mathrm{N} & \text{for } B_n \in \mathcal{U}(n) \text{ with } \mathrm{f}B_n = \mathrm{f}A_n, \text{ if } \pi_n \in (0, \omega). \end{cases} \tag{5.3}$$

It follows, for instance, that if $\pi_n = 0$ for all $n \geq m$ for some m, then

(a) $(\forall n)[s^-_n \subset s^-_{n+1}]$, and, for the least such m,

(b.1) $(\forall n < m)[\tau_n = 0 \rightarrow (\exists k, n < k \leq m)[\tau_k > 0 \text{ and hence } s_k \subset s_{k+1}]]$, i.e.,

$(\forall n < m)[\tau_n = 0 \rightarrow (\exists k, n < k \leq m)[\tau_k > 0 \text{ and hence } s_n(n) = b_n = a_k = s_{k+1}(k)]]$, and

(b.2) $(\forall n \geq m)[s_n(n) = b_n = b_m]$,

while if $\pi_n > 0$ for infinitely many n, then

(a) $(\forall n)[s^-_n \subset s^-_{n+1}]$, and

(c.1) $(\exists^\infty n)[\tau_n > 0 \text{ and hence } s_n \subset s_{n+1}]$, i.e.,

$(\forall n)[\tau_n = 0 \rightarrow (\exists k \geq n)[\tau_k > 0 \text{ and hence } s_n(n) = b_n = a_k = s_{k+1}(k)]]$.

Thus, in the first case the entire sequence $\{s_n\}_n$ can be recovered from

(b.1) s_m, in particular s^-_m and $b_m = s_m(m)$ (for the least such m as stipulated), and

(b.2) the chain of strings $s^-_{m+1} \subset s^-_{m+2} \subset \ldots$,

while in the second case we can recover the sequence $\{s_n\}_n$ from

(c.1) the set of m where $\tau_m > 0$ and

(c.2) the chain of strings $s^-_0 \subset s^-_1 \subset \ldots$.

In either case, the sequence $\{s_n\}_n$ can be recovered from the chain $\{s_n^-\}_n$ and τ (or π), plus possibly a single element b (the b_m of the first case). This allows to economise the descriptors of $A \in \mathcal{U}(\omega)$ (compared with the full set $\langle \{s_n\}_n, \pi \rangle$). In fact, we can replace $\{s_n\}_n$ by a single string $\sigma \in \mathcal{S}(\omega + 1)$.

(5.5) **Definition** (a) (Trunk-string) Define the *trunk* of $\{s_n\}_n$, $\sigma = \sigma^{A,x} \in \mathcal{S}(\omega + 1)$, by setting

$$\sigma \lceil \omega = \bigcup_n s_{n+1}^-$$

$$\sigma(\omega) = \begin{cases} b_m & \text{where } m \text{ is minimal such that } \pi_n = 0 \text{ for all } n \geq m, \text{ if such exists} \\ \omega & \text{else.} \end{cases}$$

(b) (Type-partition) Let $\delta(\tau) = \delta(\langle A, x \rangle) = 1 + |\{v \in \omega : \tau_{v+1} \neq \tau_v\}|$. Note that $\delta(\tau) \in \omega \cup \{\omega\}$. Define the *partition* $\omega = \bigcup_{n<\delta(\tau)} [v_n, v_{n+1})$ *induced by* $\tau \in {}^\omega\{0, 1, \omega\}$, by

(i) $v_0 = 0$,

(ii) $v_{n+1} = \min\{v > v_n : \tau_v \neq \tau_{v_n}\}$, for $n < \delta(\tau)$ and

(iii) $v_{n+1} = \omega$ if $\delta(\tau) = n + 1$, i.e., if n is such that for all $v > v_n$, $\tau_v = \tau_{v_n}$.

This is the coarsest partition of ω such that τ is constant on each block $[v_n, v_{n+1})$ (i.e., such that also adjacent intervals are of different types). Call $[v_n, v_{n+1})$ *of type* τ_{v_n}. The partition induced by π, resp., by $\langle A, x \rangle$, is defined as the partition induced by the type sequence τ of π, resp., by the type sequence $\tau^{\langle A,x \rangle}$ of $\langle A, x \rangle$.

Let now $s_k \in \mathcal{S}(k + 1)$ be such that $[x]_k^A = A(s_k)$, for $x \in A \in \mathcal{U}(\omega)$ and all $k \in \omega$.

(5.6) **Lemma**

(i) For an interval $[v_n, v_{n+1})$ of type 0, $b_{v_n} = b_{v_n+1} = \ldots = b_{v_{n+1}-1} = a_{v_{n+1}} = \sigma(v_{n+1})$; hence:

(ii) For k in an interval $[v_n, v_{n+1})$ of type 0, $s_k = \sigma^{A,x} \lceil k \wedge \langle b_{v_n} \rangle$. Moreover:

(iii) If $v_{n+1} < \omega$ for an interval $[v_n, v_{n+1})$ of type 0, then also $b_{v_n} = s_{v_{n+1}+1}(v_{n+1}))$.

(iv) For k in an interval $[v_n, v_{n+1})$ of type 1 or ω, $s_k = \sigma^{A,x} \lceil k + 1$.

Proof By induction over $i \in [v_n, v_{n+1})$ using the string copy lemma. •

(5.7) **Corollary**

(i) For k in $[v_n, v_{n+1})$ of type 0, $s_{k+1} = \sigma^{A,x} \lceil k \wedge \langle \sigma(v_{n+1}) \rangle$; and:

(ii) If $[v_n, v_{n+1})$ is of type 0 and finite, then in fact $s_{k+1} = \sigma^{A,x} \lceil k \wedge \langle s_{v_{n+1}+1}(v_{n+1}) \rangle$.

(iii) For k in $[v_n, v_{n+1})$ of type 1 or ω, $s_{k+1} \subset \sigma^{A,x}$. Moreover:

(iii) On $[v_n, v_{n+1})$ of type ω, $s_{k+1} = s_k \wedge \langle k \rangle \subset \sigma^{A,x}$ •

(5.8) **Definition and Lemma**

(a) Call $\sigma \in \mathcal{S}(\omega + 1)$ and $\pi \in {}^\omega(\omega + 1)$, *compatible* if $\sigma(k) = k$ iff $\pi_k = \omega$ and for every n, if $\pi_n = 0$ then for the least $m > n$ such that $\pi_m > 0$, $\sigma(m) \leq n$. I.e., if

$$\begin{aligned} \sigma(k) &= k && \text{iff } \pi_k = \omega \\ \sigma(v_{k+1}) &\leq v_k && \text{whenever } \pi_{v_k} = 0 \text{ (and hence } \pi_{v_{k+1}} > 0) \end{aligned} \tag{5.4}$$

Similarly, call $\sigma \in \mathcal{S}(\omega + 1)$ and $\tau \in {}^\omega\{0, 1, \omega\}$ *compatible* if the preceding condition holds with 'τ' in place 'π'.

(b) Call the sequence $\{s_n\}_n$, $s_n \in {}^{n+1}\omega$, *derived from* $\langle \sigma, \pi\rangle$ if

$$s_n = \begin{cases} \sigma\lceil n+1 & \text{if } \pi_n > 0 \\ \sigma\lceil n \wedge \langle \sigma(w_n)\rangle & \text{for } w_n = \inf\{v \geq n : \pi_v > 0\}, \text{ if } \pi_n = 0. \end{cases} \tag{5.5}$$

Similarly, call the sequence $\{s_n\}_n$, $s_n \in {}^{n+1}\omega$, *derived from* $\langle \sigma, \tau\rangle$, if the preceding condition holds with 'τ' replacing 'π' throughout.

Then

(i) $\langle \sigma, \pi\rangle$ are compatible iff $\langle \sigma, \tau\rangle$ are;

(ii) $\{s_n\}_n$, $s_n \in {}^{n+1}\omega$, is derived from $\langle \sigma, \tau\rangle$ iff it is derived from $\langle \sigma, \tau\rangle$;

(iii) $s_n \in \mathcal{S}(n+1)$ whenever $\langle \sigma, \pi\rangle$ (resp., $\langle \sigma, \tau\rangle$) are compatible and $\{s_n\}_n$ is derived from $\langle \sigma, \tau\rangle$.

(iv) $s_n \in \{s^{\pm}_{n+1}\}$ whenever $\langle \sigma, \pi\rangle$ (resp., $\langle \sigma, \tau\rangle$) are compatible and $\{s_n\}_n$ is derived from $\langle \sigma, \tau\rangle$, in fact $\{s_n\}_n$ is aligned with π (resp., τ).

Proof

(i) and (ii) are clear, since $\pi_k = \omega$ iff $\tau_k = \omega$. and $\pi_k = 0$ iff $\tau_k = 0$.

(iii) We need to show that $s_n(i) \leq i$ for all $i < n+1, n \in \omega$. This is clear if $\pi_n > 0$ since then $s_n(i) = \sigma(i) \leq i$ for all $i < n+1$. If $\pi_n = 0$ then $s_n(i) = \sigma(i) \leq i$ for all $i < n$, so it remains to show that $s_n(n) \leq n$ in that case also. We have $n \in [v_k, v_{k+1})$ of type 0, for some $k < \delta(\tau)$, and $w_m = v_{k+1}$ for all $m \in [v_k, v_{k+1})$. By compatibility, $\sigma(w_n) = \sigma(v_{k+1}) \leq v_k$, so in particular $\sigma(w_n) \leq n$, as required.

(iv) From the definition, $s_n\lceil n = \sigma\lceil n$, so $s^-_n \subset s^-_{n+1}$ for all n, and it remains to show that $b_n \in \{a_{n+1}, b_{n+1}\}$ for all n, where again, from the preceding, $a_{n+1} = \sigma(n)$ for all n.

(a) Case $\pi_n = 0$ and $\pi_{n+1} = 0$. Then $w_n = w_{n+1} > n+1$ and so $b_n = b_{n+1} = \sigma(w_n)$, where here, and subsequently, w_n is as in the definition.

(b) Case $\pi_n = 0$ and $\pi_{n+1} > 0$. Then $w_n = n+1$, and $s_{n+1} = \sigma\lceil n+2$, i.e., $b_{n+1} = \sigma(n+1)$ and so $b_n = \sigma(w_n) = \sigma(n+1) = b_{n+1}$.

(c) Case $\pi_n > 0$. Then immediately, $b_n = \sigma(n) = a_{n+1}$. •

Corollary The sequence $\{s_n\}_n$ derived from a compatible pair satisfies property (5.1) of realised sequences (string copy lemma). Hence $A_n = A(s_n)$ satisfy (5.3).

(5.9) **Definition and Lemma** Call σ, π, a, b, *compatible*, where $\sigma \in \mathcal{S}(\omega+1)$, $\pi \in {}^{\omega}(\omega+1)$ and $a, b \in \mathcal{S}(\omega)$, if σ, π compatible (i.e, $\sigma(k) = k$ whenever $\tau_k = \omega$), $\sigma\lceil k+1 = \sigma\lceil k \wedge \langle a_k\rangle$ and $b_{k+1} = b_k$ whenever $\pi_k = 0$, $a_{k+1} = b_k$ whenever $\pi_k \in \omega \setminus 0$. Call the sequence $\{s_n\}_n$, $s_n \in \mathcal{S}(n+1)$, *derived from* $\langle \sigma, \pi, a, b\rangle$ if $s_n = \sigma\lceil n-1 \wedge \langle a_n, b_n\rangle$. If $\{s_n\}_n$, $s_n \in \mathcal{S}(n+1)$, derives from $\langle \sigma, \pi, a, b\rangle$, then it derives from $\langle \sigma, \pi\rangle$, hence satifies (5.1).

Proof If $\pi_n \neq 0$, then either $\pi_n = \omega$ and $s_n = \sigma\lceil n+1$ by definition, or $\pi_n \in (0, \omega)$ and $s_n = \sigma\lceil n-1 \wedge \langle a_n, b_n\rangle = \sigma\lceil n-1 \wedge \langle a_n, a_{n+1}\rangle = \sigma\lceil n \wedge \langle a_{n+1}\rangle = \sigma\lceil n+1$, by induction. If $\pi_n = 0$ then $s_n = \sigma\lceil n-1 \wedge \langle a_n, b_n\rangle = \sigma\lceil n \wedge \langle b_n\rangle = \sigma\lceil n \wedge \langle w_n\rangle$ since $b_k = b_{k+1}$ for all $k \in [n, w_n)$ by compatibility. •

(5.10) **Corollary** (Summary) The following summarises the properties of the strings $\sigma = \sigma^{A,x}, \pi = \pi^{A,x}, a = a^{A,x}, b = b^{A,x}$.

(a) a_0 is undefined, $b_0 = 0$ and $[x]^A_0 \cong A(\sigma\lceil 0 \wedge \langle b_0\rangle) = A(\langle 0\rangle) = \mathbf{1}$;

(b) If $\pi(n) = \omega$ then

  (i) $[x]^A_{n+1} \cong [x]^A_n \mathrm{Z}$;

  (ii) $a_{n+1} = \sigma(n) = b_n$ and $b_{n+1} = n+1$, so

$$\begin{aligned} &\text{(ii.1)} \qquad [x]^A_n \cong A(\sigma\lceil n \wedge \langle b_n\rangle); \\ &\text{(ii.2)} \qquad [x]^A_{n+1} \cong A(\sigma\lceil n \wedge \langle b_n, n+1\rangle); \end{aligned}$$

(c) If $\pi(n) = 0$, then

  (i) $[x]^A_{n+1} \cong [x]^A_n + B_n\mathrm{N}$ for $B_n \in \mathcal{U}(n)$ such that $\mathrm{f}B_n \cong \mathrm{f}[x]^A_n \cong A(\sigma\lceil n \wedge \langle 0\rangle)$;

  (ii) $a_{n+1} = \sigma(n)$ and $b_{n+1} = b_n \leq n$, so

$$\begin{aligned} &\text{(ii.1)} \qquad [x]^A_n \cong A(s_n) \cong A(s^+_{n+1}) \cong A(s^=_{n+1} \wedge \langle b_{n+1}\rangle) \cong A(\sigma\lceil n \wedge \langle b_n\rangle), \\ &\qquad\qquad B_n \cong A(s^-_{n+1}) \cong A(s^=_{n+1} \wedge \langle a_{n+1}\rangle) \cong A(\sigma\lceil n \wedge \langle \sigma(n)\rangle) \\ &\text{(ii.2)} \qquad [x]^A_{n+1} \cong A(s_{n+1}) \cong A(s^=_{n+1} \wedge \langle a_{n+1}, b_{n+1}\rangle) \cong A(\sigma\lceil n \wedge \langle \sigma(n), b_n\rangle); \end{aligned}$$

  (iii) $\pi(n) = 0$ is the number of (n-component) predecessors of $[x]^A_n$ in $[x]^A_{n+1}$, and $[x]^A_n$ is an initial n component of $[x]^A_{n+1}$.

(d) If $\pi(n) \in \omega \setminus \{0\}$, then

  (i) $[x]^A_{n+1} \cong B_n + [x]^A_n\mathrm{N}$ for $B_n \in \mathcal{U}(n)$ such that $\mathrm{f}B_n \cong \mathrm{f}[x]^A_n \cong A(\sigma\lceil n \wedge \langle 0\rangle)$;

  (ii) $a_{n+1} = \sigma(n)$, but also $a_{n+1} = b_n$, and $b_{n+1} \leq n$, so

$$\begin{aligned} &\text{(ii.1)} \qquad B_n \cong A(s^+_{n+1}) \cong A(s^=_{n+1} \wedge \langle b_{n+1}\rangle) \cong A(\sigma\lceil n \wedge \langle b_{n+1}\rangle), \\ &\qquad\qquad [x]^A_n \cong A(s_n) \cong A(s^-_{n+1}) \cong A(\sigma\lceil n \wedge \langle a_{n+1}\rangle) \cong A(\sigma\lceil n \wedge \langle b_n\rangle) \\ &\text{(ii.2)} \qquad [x]^A_{n+1} \cong A(\sigma\lceil n \wedge \langle \sigma(n), b_{n+1}\rangle); \cong A(\sigma\lceil n \wedge \langle b_n, b_{n+1}\rangle); \end{aligned}$$

  (iii) $[x]^A_n$ is a successor n component of $[x]^A_{n+1}$ and $\pi(n) > 0$ is the number of (n-component) predecessors of $[x]^A_n$ in $[x]^A_{n+1}$.

We are now ready to state the characterisation of $A \in \mathcal{U}(\omega)$.

Let $\mathcal{G}(\omega+1)$ denote the set of pairs $\langle\sigma, \pi\rangle$ of compatible $\sigma \in \mathcal{S}(\omega+1)$, $\pi \in {}^\omega(\omega+1)$, and let $\mathcal{G}^o(\omega+1)$ denote the set of pairs $\langle\sigma, \tau\rangle$ of compatible $\sigma \in \mathcal{S}(\omega+1)$, $\tau \in {}^\omega\{0, 1, \omega\}$, respectively.

(5.11) **Lemma** (Realisation lemma) For $\langle\sigma, \pi\rangle \in \mathcal{G})\omega+1)$ and let $\{s_n\}_n$, $s_n \in \mathcal{S}(n+1)$, be the sequence derived from $\langle\sigma, \pi\rangle$. Then $\{s_n\}_n$ is realised in $\mathcal{U}(\omega)$.

Proof We need to construct $A \in \mathcal{U}(\omega)$ and $x \in A$, such that $\{s_n\}_n$ is realised by $\langle A, x\rangle$.

It will suffice to define A as the union over the chain of chains $A = \bigcup_{n\in\omega}\langle A_n, \phi_n\rangle$ where $\phi_n : A_n \hookrightarrow A_{n+1}$ is the inclusion of A_n in A_{n+1} as the n-component with π_n n-component predecessors. I.e., where $A_{n+1} = \sum_{i\in A_{n+1}/n} C_i$ shall be such that

(i) if $\pi_n \in \omega$ then $A_{n+1}/n = \mathrm{N}$, and $A_n = C_{\pi_n}$;

(ii) if $\pi_n = \omega$ then $A_{n+1}/n = \mathrm{Z}$, and $A_n = C_0$.

The element $x \in A$ for which, $A_n = [x]^A_n$ for all n is given as the x for which $\{x\} = A_0$ (really, by the 'thread' $\langle\phi_n(0)\rangle_n$, since $A_0 = \mathbf{1} = \{0\}$).

Set $A_n = A(s_n)$. Clearly then,

(a) $A_n \in \mathcal{U}(n)$;

(b) A_n is an n component of A_{n+1}, for each n (since $\{s_n\}_n$ are aligned with π and $\{A_n\}$ satisfy (5.3)), so A_{n+1}/n is as required; and

(c) since $s_n = s^+_{n+1}$ whenever $\pi_n = 0$ we can embed A_n as an initial n component into A_{n+1}, while $s_n = s^-_{n+1}$ if $\pi_n > 0$ allows to embed A_n as a successor n component into A_{n+1}.

Thus, let ϕ_n be the canonical embedding $A_n \hookrightarrow A_{n+1}$ that maps A_n isomorphically to the π_nth n component of A_{n+1} if $\pi_n < \omega$, and that so maps A_n to the 0th n component if $\pi_n = \omega$.

Thus, $\{s_n\}_n$ is realised by $\langle A, x\rangle$. •

Let

$$A(\sigma, \pi) \tag{5.6}$$

denote the element of $\mathcal{U}(\omega)$ represented by $\langle \sigma, \pi\rangle$ via the realisation lemma.

For such $A = A(\sigma, \pi)$ and the sequence $\{s_n\}$ derived from $\langle \sigma, \pi\rangle$:

(a) For some $x \in A$, $\pi^{\langle A,x\rangle} = \pi$ and for all n, $[x]^A_{n+1}/n \cong \mathrm{N}$ if $\pi_n < \omega$, $[x]^A_{n+1}/n \cong \mathrm{Z}$ if $\pi_n = \omega$, and $[x]^A_n = A(s_n)$.

(b) If $\pi_n < \omega$ then the initial n component of $[x]^A_{n+1}$ is (mod $\cong$) given by $A(s^+_{n+1})$. The successor n-components in $[x]^A_{n+1}$ are given by $A_n \cong A(s^-_{n+1}) = A(\sigma\lceil n+1)$, so by the tail lemma,

$$\mathrm{f}A \cong \sum_{n\in\omega} A_n\mathrm{N} \cong \sum_{n\in\omega} A(\sigma\lceil n+1)\mathrm{N}.$$

In fact, with a, b defined for $\langle \sigma, \pi\rangle \in \mathcal{G}(\omega+1)$ as in the aligned string sequence lemma, all of the summary applies to $A = A(\sigma, \pi)$.

(5.12) **Corollary** (Rank ω representation lemma) Every $A \in \mathcal{U}(\omega)$ has a representation of from (5.6). In particular, $A \in \mathcal{U}(\omega)$ is isomorphic to $A(\sigma, \pi)$ whenever $\sigma = \sigma^{\langle A,x\rangle}$ and $\pi = \pi^{\langle A,x\rangle}$ for some $x \in A$. •

Another corollary is that the representation of $A \in \mathcal{U}(\omega)$ by compatible pairs $\langle \sigma, \pi\rangle$ is unique up to finite initial segments of σ, π.

(5.13) **Corollary**

(i) If $A(\sigma, \pi) = A(\sigma', \pi')$, then $\sigma^- = \sigma'^-$ and $\pi =^* \pi'$ (i.e., $\pi\lceil(n, \omega) = \pi'\lceil(n, \omega)$ for some n).

(ii) If $\mathrm{f}A(\sigma, \pi) = \mathrm{f}A(\sigma', \pi')$, then $\sigma^- = \sigma'^-$ (i.e., $\sigma\lceil\omega = \sigma'\lceil\omega$). •

[5.2] Tails of rank ω

We shall verify that by computing tails of $A \in \mathcal{U}(\omega)$ in the representation just given, one obtains the representations produced in section 2. Observe that if $\sigma \in \mathcal{S}(\omega + 1)$ is compatible with $\pi = 0^\omega$ (where $0^\omega = \langle 0, 0, \ldots\rangle$ is a sequence of 0s of length ω), then it must be the case that $\sigma(\omega) = 0$ (since for all $n \in \omega$, $w_n = \omega$, i.e, for the partition induced by $\pi = 0^\omega$,

$$\delta(\pi) = 0, v_0 = 0, v_1 = \omega, \text{ and } [v_0, v_1) \text{ is of type } 0,$$

so $\sigma(\omega) = \sigma(v_1) \leq v_0 = 0$). Hence, for the sequence of strings $\{s_n\}_n$ derived from $\langle \sigma, 0^\omega \rangle \in \mathcal{G}(\omega + 1)$ we have $s_n = \sigma\lceil n + 1 \wedge \langle 0 \rangle$.

Now observe that for the point $x_0 \in A_0 = A(\sigma, 0^\omega)$ such that $\pi^{\langle A_0, x_0 \rangle} = 0^\omega$, we have $[A_0]_n^{x_0} = A(s_n) = A(\sigma\lceil n \wedge \langle 0 \rangle)$. On the other hand, for the left-most element x_1 in the chain $A_1 = \sum_{n \in \mathrm{N}} A(\sigma\lceil n + 1)\mathrm{N}$ we have $[A_1]_n^{x_1} = \sum_{i<n} A(\sigma\lceil i)\mathrm{N}, = A(\sigma\lceil n \wedge \langle 0 \rangle)$. By the block-lemma it follows that $A_0 \cong A_1$. Thus,

$$A(\sigma, 0^\omega) = A(\sigma^- \wedge \langle 0 \rangle, 0^\omega) = A(\sigma\lceil\omega \wedge \langle 0 \rangle, 0^\omega) \cong \sum_{n \in \mathrm{N}} A(\sigma\lceil n + 1)\mathrm{N}.$$

(5.14) **Corollary** (Rank ω tail lemma) For all compatible $\langle \sigma, \pi \rangle \in \mathcal{G}(\omega + 1)$,

$$\mathrm{f}A(\sigma, \pi) \cong A(\sigma^- \wedge \langle 0 \rangle, 0^\omega).$$

Proof Recall that for $x \in A \in \mathcal{U}$, $\mathrm{f}_x A \cong \sum_{0 \leq n < \mathrm{rk}(A)} A_n \mathrm{N}$, where A_n is (a representative of the isomorphism class of) the successor n-component of $[x]_n^A$ in $[x]_{n+1}^A$.

But, for $A = A(\sigma, \pi)$ in particular,

$$[x]_{n+1}^A \cong A(s_{n+1}) \in \{A(s_{n+1}^-)\mathrm{Z}, A(s_{n+1}^+) + A(s_{n+1}^-)\mathrm{N}\} \bmod \cong,$$

so $A_n = A(s_{n+1}^-) = A(\sigma\lceil n + 1)$.

Set now $A = A(\sigma^- \wedge \langle 0 \rangle, 0^\omega)$. Note that the sequence derived from $\langle \sigma, 0^\omega \rangle$ is $\{\sigma\lceil n \wedge \langle 0 \rangle\}_n$. Thus for the element $x = 0$ such that $H_0^1(A) = \mathbf{1} = \{\mathbf{x}\}$, it is the case that for all n, $[x]_n^A = A(\sigma\lceil n \wedge \langle 0 \rangle)$. Equally, for $B = \sum_{n \in \omega} A(\sigma\lceil n + 1)\mathrm{N}$, the element $y = 0$ such that $H_0^1(B) = \mathbf{1} = \{\mathbf{y}\}$, it is the case that for all n, $[y]_n^B = \sum_{i<n} A(\sigma\lceil i + 1)\mathrm{N} = A(\sigma\lceil n \wedge \langle 0 \rangle)$. So $A \cong B$.

Thus, for the element $x \in A$ giving rise to the embedding sequence π,

$$\mathrm{f}_x A \cong \sum_{0 \leq n < \mathrm{rk}(A)} A_n \mathrm{N} \cong \sum_{n \in \omega} A(\sigma\lceil n + 1)\mathrm{N} \cong A(\sigma^- \wedge \langle 0 \rangle, 0^\omega). \bullet$$

Corollary (Rank ω tail representation lemma, mark two) The following expressions range precisely over the set of all uniform tails of rank ω.

(a) For $\sigma \in \mathcal{S}(\omega + 1)$ set

$$A(\sigma) := A(\sigma, 0^\omega).$$

(b) For $\sigma \in \mathcal{S}(\omega)$ set

$$A(\sigma) := A(\sigma \wedge \langle 0 \rangle),$$

the latter being defined via (a) – i.e., $\mathring{A}(\sigma) = A(\sigma \wedge \langle 0 \rangle) := A(\sigma \wedge \langle 0 \rangle, 0^\omega)$.

[6] Chains of next limit rank

The plus finite rank chain constructor of section 3 combined with the rank ω representation of section 5 allows us to obtain approximations to (and a characterisation of) uniform chains of rank $\omega+\omega$, but the limitations of this procedure in having to step through all successor ordinals will also become apparent.

[6.1] Chains of rank $\omega+\omega$

Given $A \in \mathcal{U}(\omega{+}\omega)$, applying the rank ω representation twice, to A/ω and to the rank ω building blocks of the rank $\omega{+}n$ sub-components $[x]^A_{\omega+n}$ (for some fixed $x \in A$), one obtains a crude representation involving 'arrays' of compatible pairs: By the ω+finite uniformity lemma the chain constructed as $A(s)[M_0,\dots,M_{\ell(s)-1}]$ is uniform (of rank $\omega+|s|$) if for $\sigma_i \in \mathcal{S}(\omega+1)$ and $\pi_i \in {}^{\omega}(\omega+1)$ such that $M_i = A(\sigma_i,\pi_i)$, the ω-uniformity conditions (3.6) hold:

$$\begin{aligned}
&M_i \in \mathcal{U}(\omega) \text{ for all } i<\ell(s)\\
&\mathrm{f}M_i \cong \mathrm{f}M_j \text{ for all } i,j<\ell(s)\\
&M_i \cong M_j \text{ whenever } \psi^s(i)=\psi^s(j).
\end{aligned}$$

Combining this with the rank ω representation lemma it follows that $A(s)[M_0,\dots,M_{\ell(s)-1}]$ is uniform if the following derived ω-uniformity conditions are satisfied:

$$\begin{aligned}
&\langle\sigma_i,\pi_i\rangle \in \mathcal{G}(\omega+1) \text{ for all } i<\ell(s)\\
&\sigma_i^- = \sigma_j^- \text{ for all } i,j<\ell(s)\\
&\sigma_i=\sigma_j \text{ and } \pi_i =^* \pi_j \text{ whenever } \psi^s(i)=\psi^s(j).
\end{aligned}\tag{6.1}$$

Letting $A \in \mathcal{U}(\omega+\omega)$, by the rank ω representation lemma there are compatible $\sigma \in \mathcal{S}(\omega+1)$ and, $\pi \in {}^{\omega}(\omega+1)$ such that $A/\omega = A(\sigma,\pi)$. Again, by the rank ω representation lemma, for some $x \in A$ and each n,

$$[x]^A_{\omega+n} \cong A(s_n)[M^{(n)}_i]_{0\le i<\ell_n}$$

where

$$M^{(n)}_i = A(\sigma^{(n)}_i,\pi^{(n)}_i),\tag{6.2}$$

such that

(i) $\{s_n\}_{n\in\omega}$ derives from $\langle\sigma,\pi\rangle$,

(ii) the $M^{(n)}_i \in \mathcal{U}(\omega)$ satisfy the ω-uniformity conditions, and hence

(iii) the $\langle\sigma^{(n)}_i,\pi^{(n)}_i\rangle$ satisfy the derived conditions.

Consequently, there are $\sigma^{(n)} \in \mathcal{S}(\omega)$ such that

$$\begin{aligned}
&\sigma^{(n)-}_i = \sigma^{(n)} \text{ for all } i<\ell_n, \text{ and}\\
&\pi^{(n)}_i =^* \pi^{(n)}_j \text{ whenever } \psi_n(i)=\psi_n(j)
\end{aligned}$$

(where $\psi_n := \psi^{s_n}$).

We shall need compact compact for the leaf intervals of the $\mathcal{T}(s_{n+1}^{\pm})$ subtrees of $\mathcal{T}(s_{n+1})$ and the corresponding intervals of building blocks.

Notation Set $\mathcal{L}(s) = \ell(s) = [0, \ell(s))$, and $\mathcal{L}(s)^+ = \mathcal{L}(s^+)$, but set $\mathcal{L}(s)^- = [\ell(s) - \ell(s^-), \ell(s))$. Thus

$$\mathcal{L}(s)^- = \begin{cases} [\ell(s^+), \ell(s)) & \text{if } s^+ \in \mathcal{S} \\ [0, \ell(s)) = \mathcal{L}(s) = \mathcal{L}(s^-) & \text{else,} \end{cases}$$

If we also convene that $\ell(s) = 0$ whenever $s \notin \mathcal{S}$, then always $\mathcal{L}(s)^- = [\ell(s^+), \ell(s))$.

Set $\mathcal{L}_n = \mathcal{L}(s_n)$, and $\mathcal{L}_n^{\pm} = \mathcal{L}(s_n)^{\pm}$i when dealing with $\{s_n\}_n$ deriving from $\langle \sigma, \pi \rangle \in \mathcal{G}(\omega + 1)$. Note, $\mathcal{L}_n^- = \mathcal{L}(\sigma\lceil n+1)^-$, $\mathcal{L}_{n+1}^+ = \mathcal{L}_n$ and $\mathcal{L}_{n+1}^- = [\ell_n, \ell_{n+1})$ if $\tau_n = 0$, $\mathcal{L}_{n+1}^+ = [0, \ell_{n+1} - \ell_n)$ and $\mathcal{L}_{n+1}^- = [\ell_{n+1}^+, \ell_{n+1})$ if $\tau_n = 1$.

Set $\mathbf{M}_n = \mathbf{M}^{(n)} = \mathbf{M}_{\mathcal{L}_n} = \langle M_i^{(n)} : i \in \mathcal{L}_n \rangle$ and similarly, $\mathbf{M}_n^{\pm} = \mathbf{M}_{\mathcal{L}_n}^{\pm} = \mathbf{M}_{\mathcal{L}_n^{\pm}} = \langle M_i^{(n)} : i \in \mathcal{L}_n^{\pm} \rangle$, and for $\mathcal{J} \subseteq \mathcal{L}_n$, $\mathbf{M}_{\mathcal{J}}^{(n)} = \langle M_i^{(n)} : i \in \mathcal{J} \rangle$, etc.

Let now $x \in A \in \mathcal{U}(\omega+\omega)$ such that for each n setting $M_i^{(n)} = A(\sigma_i^{(n)}, \pi_i^{(n)})$. Then $[x]_{\omega+n}^A = A(s_n)[\mathbf{M}_n]$, and by plus finite chain construction,

$$A(s_{n+1})[\mathbf{M}_{n+1}] = \begin{cases} A(s_{n+1}^-)[\mathbf{M}_{n+1}^-].\mathrm{Z} & \text{if } \tau_n = \omega \\ A(s_{n+1}^+)[\mathbf{M}_{n+1}^+] + A(s_{n+1}^-)[\mathbf{M}_{n+1}^-].\mathrm{N} & \text{else,} \end{cases}$$

so:

(6.1) **Lemma** (Copy lemma) Either

$$\tau_n = 0,\ s_n = s_{n+1}^+, \text{ and } A(s_n)[\mathbf{M}_n] = A(s_{n+1}^+)[\mathbf{M}_{n+1}^+],$$

or

$$\tau_n > 0,\ s_n = s_{n+1}^- = \sigma\lceil_{n+1}, \text{ and } A(s_n)[\mathbf{M}_n] = A(s_{n+1}^-)[\mathbf{M}_{n+1}^-],$$

and in particular,

$$\mathbf{M}_{n+1} = \begin{cases} \mathbf{M}_n & \text{if } s_{n+1}^+ \notin \mathcal{S} \\ \mathbf{M}_n \ \wedge\ \mathbf{M}_{n+1}^- & \text{if } s_{n+1}^+ \in \mathcal{S} \text{ and } s_{n+1}^+ = s_n \\ \mathbf{M}_{n+1}^+ \ \wedge\ \mathbf{M}_n & \text{if } s_{n+1}^+ \in \mathcal{S} \text{ and } s_{n+1}^- = s_n \end{cases}$$

i.e.,

$$\mathbf{M}_n = \begin{cases} \mathbf{M}_{n+1}^- & \text{if } \tau_n > 0 \\ \mathbf{M}_{n+1}^+ & \text{if } \tau_n = 0 \end{cases}$$

(i.e., the 's_n-array' $\mathbf{M}_n$ gets copied as one of the segments $\mathcal{L}_{n+1}^+$ or $\mathcal{L}_{n+1}^-$ of the 's_{n+1} array' $\mathbf{M}_{n+1}$.)

From

$$\mathrm{f}M_i^{(n)} \cong \mathrm{f}M_0^{(n)} \cong \mathrm{f}M_0^{(n+1)} \cong \mathrm{f}M_j^{(n+1)}$$

for all $i < \ell_n$ and $j < \ell_{n+1}$ it follows that in fact there is a single 'trunk' $\sigma' \in \mathcal{S}(\omega)$ such that for all $n \in \omega$ and $i < \ell_n$,

$$\sigma_i^{(n)-} = \sigma_i^{(n)}\lceil\omega = \sigma'\lceil n.$$

Thus we get:

(6.2) **Lemma**

[1] For every $A \in \mathcal{U}(\omega+\omega)$,

(i) there exists $\sigma' \in \mathcal{S}(\omega)$ such that for all $n \in \omega, i < \ell_n$, $\sigma_i^{(n)-} = \sigma' \lceil n$, and $\sigma_i^{(n)} = \sigma_j^{(n)}$ whenever $\psi_n(i) = \psi_n(j)$;

(ii) for all $n \in \omega$, for all $i < \ell_n$, $\pi_i^{(n)}$ is compatible with σ', and for all $n, m \in \omega$, for all $i < \ell_n, j < \ell_m$, $\pi_i^{(n)} =^* \pi_j^{(m)}$;

(iii) for all $n \in \omega$, if $\pi_n = 0$ then $s_n = \sigma_{n+1}^+$ and $\langle \pi_i^{(n)} \rangle_{0 \leq i < \ell_n} = \langle \pi_i^{(n+1)} \rangle_{0 \leq i < \ell_{n+1}^+}$; and

(iv) for all $n \in \omega$, if $\pi_n > 0$ then $s_n = \sigma_{n+1}^-$ and $\langle \pi_i^{(n)} \rangle_{0 \leq i < \ell_n} = \langle \pi_i^{(n+1)} \rangle_{\ell_{n+1}^+ \leq i < \ell_{n+1}}$.

Moreover, the set of $\langle \sigma, \pi \rangle \in \mathcal{G}(\omega+1)$, $\langle \sigma^{(n)}, \pi^{(n)} \rangle \in \mathcal{G}(\omega+1)$ satisfying (i) through (iv) is uniquely determined by A.

[2] Every $A \in \mathcal{U}(\omega+\omega)$ is uniquely determined by set of $\langle \sigma, \pi \rangle \in \mathcal{G}(\omega+1)$, $\langle \sigma^{(n)}, \pi^{(n)} \rangle \in \mathcal{G}(\omega+1)$ satisfying (i) through (iv), and can be reconstructed from this set by applying the rank ω representation lemma to each $M_i^{(n)}$ and then to A/ω. •

Corollary (Crude rank $\omega+\omega$ representation lemma) Let

$$A(\sigma,\pi)[\mathbf{M}_n]_{n\in\omega} = A(\sigma, \pi, \langle \sigma_i^{(n)}, \pi_i^{(n)} \rangle_{n \in \omega, i \in \mathcal{L}_n}) \tag{6.3}$$

denote the chain described in the $\omega+\omega$ representation lemma, i.e., where $\langle \sigma, \pi \rangle, \langle \sigma_i^{(n)}, \pi_i^{(n)} \rangle \in \mathcal{G}(\omega+1)$ $(n \in \omega, i \in \mathcal{L}_n)$ satisfy conditions (i)-(iv) of the preceding lemma, and where $M_i^{(n)} = A(\sigma_i^{(n)}, \pi_i^{(n)})$. Then, every $A \in \mathcal{U}(\omega+\omega)$ is of either of the forms (6.3), for some $\langle \sigma, \pi \rangle, \langle \sigma_i^{(n)}, \pi_i^{(n)} \rangle \in \mathcal{G}(\omega+1)$ $(n \in \omega, i \in \mathcal{L}_n)$, or some array $\langle \mathbf{M}_n \rangle_{n\in\omega}$ satsfying the uniformity conditions (6.1), respectively.

This representation lemma simply summarizes what we get on the basis of rank ω representation and rank ω+finite approximation alone. Removing the redundancy in the $\mathbf{M}_n$ arrays will allow us to stream-line this 'representation' considerably.

[6.2] Tails of rank $\omega+\omega$

To compute the tails of the components $[x]_{\omega+n}^A$ and of A itself we use the 'usual' strategy (basing ourselves on the uniformity over limits lemma, but going back to the tail lemma for the calculations):

$$\mathrm{f}[x]_{\omega+n}^A = \mathrm{f}_x[x]_{\omega+n}^A = \mathrm{f}_x[x]_\omega^A + \sum_{i<n} S_{\omega+i}^A.\mathrm{N}$$

where S_α^A is the (isomorphism type) of the sucessor component of $[x]_\alpha^A$ in component $[x]_{\alpha+1}^A$. Since

$$\begin{aligned} [x]_{\omega+n+1}^A &= A(s_{n+1})[\mathbf{M}_{n+1}] \\ &= \begin{cases} A(s_{n+1}^-)[\mathbf{M}_{n+1}^-].\mathrm{Z}, & \text{if } s_{n+1}^+ \notin \mathcal{S} \\ A(s_{n+1}^+)[\mathbf{M}_{n+1}^+] + A(s_{n+1}^-)[\mathbf{M}_{n+1}^-].\mathrm{N}, & \text{else} \end{cases} \end{aligned}$$

it follows that the successor $\omega+n$ component of $[x]_{\omega+n}^A$ in $[x]_{\omega+n+1}^A$ is

$$S_{\omega+n}^A \cong A(s_{n+1}^-)[\mathbf{M}_{n+1}^-] \cong A(\sigma\lceil_{n+1})[\mathbf{M}_{n+1}^-].$$

(And $[x]^A_\omega = M^{(0)}_0 = A(\sigma^{(0)}, \pi^{(0)})$.) Thus (applying the ω tail lemma for the initial component),

$$\begin{aligned} \mathrm{f}_x A &= \mathrm{f}_x A(\langle 0\rangle)[A(\sigma^{(0)}_0, \pi^{(0)}_0)] + \sum_{n\in\omega} A_{\omega+n}.\mathrm{N} \\ &= \mathrm{f}A(\sigma^{(0)}_0, \pi^{(0)}_0) + \sum_{n\in\omega} A_{\omega+n}.\mathrm{N} \\ &= A(\sigma' \wedge \langle 0\rangle) + \sum_{n\in\omega} A(\sigma\lceil_{n+1})[\mathbf{M}^-_{n+1}].\mathrm{N} \end{aligned}$$

Thus, for $x \in A = A(\sigma, \pi)[\mathbf{M}_n]$ we get

$$\begin{aligned} \mathrm{f}_x A &= \mathrm{f}_x [x]^A_\omega + \sum_{n\in\omega} A_{\omega+n+1}\mathrm{N} \\ &= \mathrm{f}M^{(0)}_0 + \sum_{n\in\omega} A(s^{(0)-}_{n+1})[\mathbf{M}^-_{n+1}].\mathrm{N} \\ &= \sum_{n\in\omega} A(\sigma'\lceil_{n+1})\mathrm{N} + \sum_{n\in\omega} A(\sigma\lceil_{n+1})[\mathbf{M}^-_{n+1}].\mathrm{N} \end{aligned}$$

where $\{s^0_{n+1}\}_{n\in\omega}$ derives from $\sigma^{(0)}_0$, and where σ' is the common initial ω segment of all the $\sigma^{(n)}_i$ (i.e., $\sigma' = \sigma^{(n)}_i\lceil\omega$, for all n, i).

(6.3) **Lemma** (Crude rank $\omega+\omega$ tail lemma) For every $A \in \mathcal{U}(\omega+\omega)$, there are $\sigma \in \mathcal{S}(\omega+1)$ and $\langle \sigma^{(n)}, \pi^{(n)}\rangle \in \mathcal{G}(\omega+1)$, such that

$$\mathrm{f}A = \sum_{n\in\omega} A(\sigma^{(0)}\lceil_{n+1})\mathrm{N} + \sum_{n\in\omega} A(\sigma\lceil_{n+1})[\mathbf{M}^-_{n+1}]\mathrm{N},$$

and where $\sigma^{(0)}\lceil\omega = \sigma^{(n)}\lceil\omega$ for all n. •

Given $\langle\sigma, \pi\rangle \in \mathcal{G}(\omega+1)$ and the derived sequence $\{s_n\}_n$, define sequence $\{k_n\}_{n\in\omega}$ by $k_0 = 0$ and $k_{n+1} = k_n + \ell_{n+1} - \ell_n$. Set $k^{\sigma,\pi} = \sup_n k_n$.

Given a double array $\mathbf{M}_{n+1}$ deriving from a chain $A \in \mathcal{U}(\omega+\omega)$ define the sequence $\langle V_i\rangle_{i<k^{\sigma,\pi}}$ by $V_0 = M^{(0)}_0$ and supposing $\{V_i\}_{i<k_n}$ defined,

$$\mathbf{V}_{n+1} = \mathbf{V}_{[k_n, k_{n+1})} = \langle V_i\rangle_{k_n \le i < k_{n+1}} = \begin{cases} \langle\,\rangle & \text{if } \tau_n = \omega, \\ \mathbf{M}^-_{n+1} & \text{if } \tau_n = 0 \ (s_n = s^+_{n+1}), \\ \mathbf{M}^+_{n+1} & \text{if } \tau_n = 1 \ (s_n = s^-_{n+1}) \end{cases}$$

I.e., the V_i solve the following set of equations:

$$\begin{aligned} &V_0 = M^{(0)}_0, \\ &\mathbf{M}_{n+1} = \begin{cases} \mathbf{M}_n \ \wedge \ \mathbf{V}_{n+1} & \text{if } \tau_n = 0, \\ \mathbf{V}_{n+1} \ \wedge \ \mathbf{M}_n & \text{if } \tau_n = 1. \end{cases} \end{aligned}$$

The V_i, $i \in [k_n, k_{n+1})$ contain the bulding-blocks that are not already contained in the arrays $\mathbf{M}_k, k < n$.

Tails of rank $\omega+\omega$ represent a novel challenge: When in the rank $\omega + (< \omega)$ approximation the lower rank component gets embedded with $\tau_n = 1$ (i.e., there is an initial rank $\omega + n$ component in the rank

$\omega+n+1$ approximation not necessarily isomorphic to the rank $\omega+n$ approximation, the building-blocks of the former are not all 'visible' in the tail:

(6.4) **Corollary** (Tail revelation lemma)

$$\mathrm{f}A = \mathrm{f}A(\sigma^{(0)}, \pi^{(0)}) + \sum_{n\in\omega} S^A_{\omega+n}.\mathrm{N},$$

where the (isomorphism type of the) successor $\omega+n$ component $S^A_{\omega+n}$ of $A = A(\sigma, \pi, \langle \mathbf{M}_n : n \in \omega\rangle) \in \mathcal{U}(\omega+\omega)$, satisfies

$$S^A_{\omega+n} = A(s^-_{n+1})[\mathbf{M}^-_{n+1}] = \begin{cases} A(s^-_{n+1})[\mathbf{V}_{n+1}] & \text{if } \tau_n = 0, \\ A(s^-_{n+1})[\mathbf{M}_n] & \text{if } \tau_n \neq 0. \end{cases}$$

Where $S^A_{\omega+n} = A(s_n)[\mathbf{M}_n]$, the 'new' building blocks at level $n+1$ are 'hidden' from the tail, but they will be 'revealed' at the next higher level $m > n$ where $S^A_{\omega+m} = A(s_n)[\mathbf{M}_m]$ — because $\mathbf{V}_{n+1}$ is a sub-interval of building blocks of $\mathbf{M}_m$. As a consequence, there can be at most one level introducing new building blocks that may stay 'hidden' from the entire approximation to the tail, namely if for some largest n, $\tau_{\omega+n} = 1$ and for all $m > n$, $\tau_{\omega+m} = 1$. Anticipating that at any level there is in fact atmost one new building block, it follows that at most one such can stay permanently 'hidden' from the tail.

It's now time to prepare the grounds for a managable representation.

[6.3] Array reduction arithmetic

Set $w_n = w^{s_n}$ (where $w^s = \mathrm{ran}(\psi^s)$). By the ω+finite uniformity lemma, there is a sequence of sets $\{U^{(n)}_i : i \in w_n\} \subset \mathcal{U}(\omega)$ $(n \in \omega)$, such that for each n

$$(\forall i \in \ell_n)[M^{(n)}_i = U^{(n)}_{\psi_n(i)}].$$

Let $\mathbf{U}_n = \{\langle U^{(n)}_k, k\rangle : k \in w_n\}$ be the tagged short array for $[x]^A_{\omega+n}$, i.e., such that $[x]^A_{\omega+n} = A(s_n)\{\mathbf{U}_n\}$. By the copy lemma

$$\mathbf{U}_n = \begin{cases} (\mathbf{U}_{n+1})^+_{s_{n+1}} & \text{if } \tau_n = 0 \\ (\mathbf{U}_{n+1})^-_{s_{n+1}} & \text{if } \tau_n > 0 \end{cases}$$

We'd like to obtain a short array for $A \in \mathcal{U}(\omega+\omega)$ as a sequence $\mathbf{U} = \{U_k : k \in K\}$ (for some set K) that would allow to 'generate' the $\mathbf{U}_n$ perhaps resembling the way in which the s_n were generated from $\sigma \in \mathcal{S}(\omega+1)$ (and compatible τ). We shall first show that we can in fact obtain a 'canonically ordered' short array, and subsequently how to generate short arrays for the sub-components of A in the approximating chains $[x]^A_{\omega+n}$.

Set $\xi^s = \max_{i<\ell(s)} \psi^s(i)$ (and again, $\xi_n = \xi^{s_n}$). Define the *augmented weight sequence* by:

$$\widehat{\psi}^s = \langle \psi^s(0)+1\rangle \wedge \langle \psi^s(i) : 0 < i < \ell(s)\rangle = \langle \psi^s(0)+1\rangle \ \wedge \ \psi^s\lceil[1, \ell(s)).$$

(Also let $\widehat{w}^s = r\widehat{\psi}^s$; note that $|\widehat{w}^s| = |w^s| = \zeta^s$.)

From the corollaries to the weight induction lemma, 3.9, we have:

(6.5) **Lemma** (Weight sequence lemma, maximum weight lemma) For $s \in \mathcal{S}$,

(a)

$$\psi^s = \begin{cases} \psi^{s^-} & \text{if } s^+ \notin \mathcal{S} \\ \psi^{s^+} \wedge \psi^{s^-} & \text{if } s^+ \in \mathcal{S} \text{ and } s \neq s^- \wedge \langle 0 \rangle \\ \widehat{\psi}^{s^+} \wedge \psi^{s^-} & \text{if } s^+ \in \mathcal{S} \text{ and } s = s^- \wedge \langle 0 \rangle \end{cases}$$

(b)

$$\xi^s = \begin{cases} \xi^{s^-} & \text{if } s^+ \in \mathcal{S} \text{ and } \xi^{s^-} > \xi^{s^+}, \text{ or } s^+ \notin \mathcal{S} \\ \xi^{s^+} & \text{if } s^+ \in \mathcal{S} \text{ and } \xi^{s^-} \leq \xi^{s^+} \text{ and } s \neq s^- \wedge \langle 0 \rangle \\ \xi^{s^+} + 1 & \text{if } s^+ \in \mathcal{S} \text{ and } \xi^{s^-} \leq \xi^{s^+} \text{ and } s = s^- \wedge \langle 0 \rangle. \end{cases}$$

(c)

$$\xi^s \leq |s| - 1 \qquad \text{i.e.,}$$

$$\xi^s \leq n \text{ whenever } s \in \mathcal{S}(n+1)$$

(d)

$$\xi^{s \wedge \langle 0 \rangle} = \psi^{s \wedge \langle 0 \rangle}(0) = |s|$$

$$\xi^{s \wedge \langle b \rangle}, \psi^{s \wedge \langle b \rangle}(0) < |s| \text{ if } 0 < b \leq |s|$$

i.e.,

$$\xi^s = \psi^s(0) = |s| - 1 \text{ if } s = s^- \wedge \langle 0 \rangle$$

$$\xi^s, \psi^s(0) < |s| - 1 \text{ if } s = s^- \wedge \langle b \rangle \text{ for } 0 < b \leq |s|$$

Proof (a) and (b) follow from 3.9, (c) and (d) follow by induction. •

The maximum weight lemma almost seems to suggest that $\xi_{n+1} \in \{\xi_n, \xi_n + 1\}$. It may come as somewhat of a surprise that this does not hold. In fact, as the examples in preceding sections suggest quite the contrary. This is because the weight-incrementation of the branching nodes proceeds rather indiscriminately. Consider the example $\mathcal{T}(\langle 0, 1, 0 \rangle)$,


$\langle\langle 0,1,0\rangle; 0\rangle$
$\langle\langle 0,0\rangle; 1\rangle$
$\mathcal{T}(\langle 0,1\rangle)$
$\langle\langle 0\rangle; 2\rangle$
$\mathcal{T}(\langle 0\rangle)$

where it would have sufficed to 'copy', rather than increment, the weight from node $\langle 0, 0 \rangle$ to node $\langle 0 \rangle$, in order to distingish it in the weight-sequence ($\langle 1, 0, 0 \rangle$, rather than $\langle 2, 0, 0 \rangle$) from the two weight 0 nodes.

Nevertheless, something of this earlier intuition can be salvaged; we do have that at every stage at most one new 'weight-element' is introduced — and consequently at most one new rank ω 'building block' in the array of rank ω components. More precisely, what we will show is that

$$0 \leq \zeta_{n+1} - \zeta_n \leq 1,$$

where $\zeta^s = |w^s|$ and $\zeta_n = \zeta^{s_n}$. (Thus at every rank $\omega + n$ at most one 'new' type of ω component appears in the array of 'building blocks'.)

(6.6) **Lemma** (Plus one range lemma) For $s \in \mathcal{S}$ and $0 < b \leq |s|$,

$$w^{s\wedge\langle 0\rangle} \triangle w^{s\wedge\langle b\rangle} = w^{s\wedge\langle 0\rangle} \setminus w^{s\wedge\langle b\rangle} = \{\psi^{s\wedge\langle 0\rangle}(0)\}.$$

Proof By induction on $|s|$.

$|s| = 1$: Then $s \wedge \langle 0\rangle = \langle 0, 0\rangle$ and the only possiblility for b is $b = 1 = |s|$, so $s \wedge \langle b\rangle = \langle 0, 0\rangle$. Then $\psi^{s\wedge\langle 0\rangle} = \langle 1, 0\rangle$ and $\psi^{s\wedge\langle b\rangle} = \psi^s = \langle 0\rangle$, and the conclusion holds. Since this only covers the case $b = |s|$, we give the case $|s| = 2$ as well.

$|s| = 2$: There are two cases for s and for each of these two cases for b:

$s = \langle 0, 0\rangle$: Then $s \wedge \langle 0\rangle = \langle 0, 0, 0\rangle$, so $\psi^{s\wedge\langle 0\rangle} = \psi^{\langle 0,0,0\rangle} = \langle 2, 0, 1, 0\rangle$. For $b = 1$, $\psi^{s\wedge\langle 1\rangle} = \psi^{\langle 0,0,1\rangle} = \langle 0, 1, 0\rangle$, so

$$w^{s\wedge\langle 0\rangle} \triangle w^{s\wedge\langle b\rangle} = w^{s\wedge\langle 0\rangle} \setminus w^{s\wedge\langle b\rangle} = \{0, 1, 2\} \setminus \{0, 1\} = \{2\} = \{\psi^{s\wedge\langle 0\rangle}(0)\},$$

as required. For $b = 2$, $\psi^{s\wedge\langle 2\rangle} = \psi^{\langle 0,0,2\rangle} = \psi^{\langle 0,0\rangle} = \langle 1, 0\rangle$, and the same calculation applies as before.

$s = \langle 0, 1\rangle$: Then $s \wedge \langle 0\rangle = \langle 0, 1, 0\rangle$, so $\psi^{s\wedge\langle 0\rangle} = \psi^{\langle 0,1,0\rangle} = \langle 2, 0, 0\rangle$. For $b = 1$, $\psi^{s\wedge\langle 1\rangle} = \psi^{\langle 0,1,1\rangle} = \langle 0, 0\rangle$, so

$$w^{s\wedge\langle 0\rangle} \triangle w^{s\wedge\langle b\rangle} = w^{s\wedge\langle 0\rangle} \setminus w^{s\wedge\langle b\rangle} = \{0, 2\} \setminus \{0\} = \{2\} = \{\psi^{s\wedge\langle 0\rangle}(0)\},$$

as required. For $b = 2$, $\psi^{s\wedge\langle 2\rangle} = \psi^{\langle 0,1,2\rangle} = \psi^{\langle 0\rangle} = \langle 0\rangle$, and the calculation is as before.

For the induction step, distinguish cases as follows:

Case $0 < b < |s|$. Then

$$\psi^{s\wedge\langle 0\rangle} = \widehat{\psi}^{s\wedge\langle 0\rangle^+} \ \wedge \ \psi^{s\wedge\langle 0\rangle^-} = \widehat{\psi}^{s^-\wedge\langle 0\rangle} \ \wedge \ \psi^s.$$

Since $0 < b < |s|$, also

$$\psi^{s\wedge\langle b\rangle} = \psi^{s\wedge\langle b\rangle^+} \ \wedge \ \psi^{s\wedge\langle b\rangle^-} = \psi^{s^-\wedge\langle b\rangle} \ \wedge \ \psi^s.$$

The respective 'second' half segments being identical, and since $\psi^{s\wedge\langle 0\rangle}(0) = \psi^{s^-\wedge\langle 0\rangle}(0) + 1$,

$$\begin{aligned} w^{s\wedge\langle 0\rangle} \triangle w^{s\wedge\langle b\rangle} &= \widehat{w}^{s^-\wedge\langle 0\rangle} \triangle w^{s^-\wedge\langle b\rangle} \\ &= w^{s^-\wedge\langle 0\rangle} \setminus \{\psi^{s^-\wedge\langle 0\rangle}(0)\} \cup \{\psi^{s^-\wedge\langle 0\rangle}(0) + 1\}) \triangle w^{s^-\wedge\langle b\rangle}. \end{aligned}$$

Now (e.g., by the weight sequence lemma, (c), $\psi^{s^-\wedge\langle 0\rangle}(0) + 1 \notin w^{s^-\wedge\langle b\rangle}$. Observe that for sets A, B and elements $a \in A \setminus B$ and $\tilde{a} \notin B$,

$$(A \setminus \{a\} \cup \{\tilde{a}\}) \setminus B = (A \setminus B) \setminus \{a\} \cup \{\tilde{a}\},$$

and if also $B \setminus A = \emptyset$, then also $B \setminus (A \setminus \{a\} \cup \{\tilde{a}\}) = \emptyset$, and so

$$(A \setminus \{a\} \cup \{\tilde{a}\}) \triangle B = (A \setminus B) \setminus \{a\} \cup \{\tilde{a}\}.$$

Thus, by inductive hypothesis,

$$\begin{aligned} &w^{s^-\wedge\langle 0\rangle} \setminus \{\psi^{s^-\wedge\langle 0\rangle}(0)\} \cup \{\psi^{s^-\wedge\langle 0\rangle}(0) + 1\}) \triangle w^{s^-\wedge\langle b\rangle} \\ &= w^{s^-\wedge\langle 0\rangle} \setminus w^{s^-\wedge\langle b\rangle} \setminus \{\psi^{s^-\wedge\langle 0\rangle}(0)\} \cup \{\psi^{s^-\wedge\langle 0\rangle}(0) + 1\} \\ &= w^{s^-\wedge\langle 0\rangle} \triangle w^{s^-\wedge\langle b\rangle} \setminus \{\psi^{s^-\wedge\langle 0\rangle}(0)\} \cup \{\psi^{s^-\wedge\langle 0\rangle}(0) + 1\} \\ &= \{\psi^{s^-\wedge\langle 0\rangle}(0) + 1)\}, \\ &= \{\psi^{s\wedge\langle 0\rangle}(0)\}, \end{aligned}$$

as required.

Case $0 < b = |s|$. Then $\psi^{s\wedge\langle b\rangle} = \psi^s$, so

$$\begin{aligned} w^{s\wedge\langle 0\rangle} \triangle w^{s\wedge\langle b\rangle} &= \widehat{w}^{s^-\wedge\langle 0\rangle} \triangle w^s \\ &= w^{s^-\wedge\langle 0\rangle} \setminus \{\psi^{s^-\wedge\langle 0\rangle}(0)\} \cup \{\psi^{s^-\wedge\langle 0\rangle}(0)+1\}) \triangle \widehat{w}^s \\ &= \{\psi^{s^-\wedge\langle 0\rangle}(0)+1)\}, \end{aligned}$$

again, as required. •

(6.7) **Corollary** For $s \in \mathcal{S}$ and $0 < a, b \leq |s|$,

(i) $w^{s\wedge\langle a\rangle} \triangle w^{s\wedge\langle b\rangle} = \emptyset$.

(ii) $|\widehat{w}^{s\wedge\langle 0\rangle} \setminus w^{s\wedge\langle b\rangle}| = |\widehat{w}^{s\wedge\langle 0\rangle} \triangle w^{s\wedge\langle b\rangle}| = |w^{s\wedge\langle 0\rangle} \setminus w^{s\wedge\langle b\rangle}| = |w^{s\wedge\langle 0\rangle} \triangle w^{s\wedge\langle b\rangle}| \leq 1$.

(iii) $0 \leq \zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} \leq 1$ and $0 \leq \zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle b\rangle} \leq 1$

Proof (i) $w^{s\wedge\langle a\rangle} \triangle w^{s\wedge\langle b\rangle} = (w^{s\wedge\langle 0\rangle} \setminus \{\psi^{s\wedge\langle 0\rangle}(0)\}) \triangle (w^{s\wedge\langle 0\rangle} \setminus \{\psi^{s\wedge\langle 0\rangle}(0)\}) = \emptyset$.

(ii) By the plus one range lemma, $w^{s\wedge\langle 0\rangle} \setminus w^{s\wedge\langle b\rangle} = \{\psi^{s\wedge\langle 0\rangle}(0)\} = \{\xi^{s\wedge\langle 0\rangle}\}$, and by the preceding lemma $\xi^{s\wedge\langle 0\rangle} > \xi^{s\wedge\langle b\rangle}$, so $\widehat{w}^{s\wedge\langle 0\rangle} \setminus w^{s\wedge\langle b\rangle} = \{\widehat{\psi}^{s\wedge\langle 0\rangle}(0)\} = \{\xi^{s\wedge\langle 0\rangle}+1\}$.

(iii) Distinguish cases, and apply the plus one range lemma to the ranges compared.

Case $a = b = 0$: $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} = |r(\widehat{\psi}^{s\wedge\langle 0\rangle} \wedge \psi^{s\wedge\langle 0\rangle})| - \zeta^{s\wedge\langle a\rangle} = 1$ and ditto $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle b\rangle} = 1$

Case $a > b = 0$: $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} = |r(\widehat{\psi}^{s\wedge\langle 0\rangle} \wedge \psi^{s\wedge\langle a\rangle})| - \zeta^{s\wedge\langle a\rangle} = 1$ and $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} = |r(\widehat{\psi}^{s\wedge\langle 0\rangle} \wedge \psi^{s\wedge\langle a\rangle})| - \zeta^{s\wedge\langle 0\rangle} = 1$

Case $a = 0 < b$: $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} = |r(\psi^{s\wedge\langle b\rangle} \wedge \psi^{s\wedge\langle 0\rangle})| - \zeta^{s\wedge\langle 0\rangle} = 0$ and $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} = |r(\psi^{s\wedge\langle b\rangle} \wedge \psi^{s\wedge\langle 0\rangle})| - \zeta^{s\wedge\langle b\rangle} = 1$

Case $0 < a, b$: $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} = |r(\psi^{s\wedge\langle b\rangle} \wedge \psi^{s\wedge\langle a\rangle})| - \zeta^{s\wedge\langle a\rangle} = 0$ and $\zeta^{s\wedge\langle a,b\rangle} - \zeta^{s\wedge\langle a\rangle} = |r(\psi^{s\wedge\langle b\rangle} \wedge \psi^{s\wedge\langle a\rangle})| - \zeta^{s\wedge\langle b\rangle} = 0$. •

By the preceding lemma,

Corollary For each n,

(i) $\zeta_{n+1} - \zeta_n \leq 1$.

(ii) w_{n+1} is an end-extension of w_n, since either $w_{n+1} = w_n$, or both $w_{n+1} \supseteq w_n$ and $\min(w_{n+1} \setminus w_n) = \xi_{n+1} > \xi_n$.

(6.8) **Lemma** (Plus minus lemma) For all $s \in \mathcal{S}(<\omega)$, where $s^+ \in \mathcal{S}$,

$$|w^{s^+} \triangle w^{s^-}| \leq 1$$

Proof By induction on $\mathcal{D}(s)$. Thus, suppose that for all $t \in \mathcal{D}(s)$ with $t^+ \in \mathcal{S}$, $|w^{t^+} \triangle w^{t^-}| \leq 1$. By weight sequence lemma, (a),

$$\psi^{s^+} = \begin{cases} \psi^{s^{+-}} & \text{if } s^{++} \notin \mathcal{S} \\ \psi^{s^{++}} \ \wedge \ \psi^{s^{+-}} & \text{if } s^{++} \in \mathcal{S} \text{ and } s^+ \neq s^{--} \wedge \langle 0\rangle \\ \widehat{\psi}^{s^{++}} \ \wedge \ \psi^{s^{+-}} & \text{if } s^{++} \in \mathcal{S} \text{ and } s = s^{--} \wedge \langle 0\rangle, \end{cases}$$

and

$$\psi^{s^-} = \begin{cases} \psi^{s^{--}} & \text{if } s^{-+} \notin \mathcal{S} \\ \psi^{s^{-+}} \wedge \psi^{s^{--}} & \text{if } s^{-+} \in \mathcal{S} \text{ and } s^- \neq s^{--} \wedge \langle 0 \rangle \\ \widehat{\psi}^{s^{-+}} \wedge \psi^{s^{--}} & \text{if } s^{-+} \in \mathcal{S} \text{ and } s^- = s^{--} \wedge \langle 0 \rangle. \end{cases}$$

Let $s = t \wedge \langle a, b \rangle$ $(0 \leq a, b \leq |t|)$. By the preceding,

$$\psi^{s^+} = \psi^{t \wedge \langle b \rangle} = \begin{cases} \psi^t & \text{if } b = |t| \\ \psi^{t^- \wedge \langle b \rangle} \wedge \psi^t & \text{if } 0 < b < |t| \\ \widehat{\psi}^{t^- \wedge \langle 0 \rangle} \wedge \psi^t & \text{if } b = 0, \end{cases}$$

and

$$\psi^{s^-} = \psi^{t \wedge \langle a \rangle} = \begin{cases} \psi^t & \text{if } a = |t| \\ \psi^{t^- \wedge \langle a \rangle} \wedge \psi^t & \text{if } 0 < a < |t| \\ \widehat{\psi}^{t^- \wedge \langle 0 \rangle} \wedge \psi^t & \text{if } a = 0. \end{cases}$$

We shall make heavy use of $s^{+-} = s^{--}$, which we refer to below as t. Write $A \approx B$ for $|A| = |B|$ and $A \preceq B$ for $|A| \leq |B|$, etc.

Case 1 $s^{++} \notin \mathcal{S}$ (i.e., $b = |t|$).

Sub-case 1.1 $s^{-+} \notin \mathcal{S}$ (i.e., $a = |t|$): Then

$$w^{s^+} \bigtriangleup w^{s^-} = w^t \bigtriangleup w^t = \emptyset.$$

Sub-case 1.2 $s^{-+} \in \mathcal{S}$ and $s^- \neq s^{--} \wedge \langle 0 \rangle$ (i.e., $0 < a \leq |t|$):

$$\begin{aligned} w^{s^+} \bigtriangleup w^{s^-} &= w^t \bigtriangleup r(\psi^{t^- \wedge \langle a \rangle} \wedge \psi^t) \\ &= w^{t^- \wedge \langle a \rangle} \setminus w^t \preceq 1, \end{aligned}$$

by the induction hypothesis appied to $s^- = t \wedge \langle a \rangle$.

Sub-case 1.3 $s^{-+} \in \mathcal{S}$ and $s^- = s^{--} \wedge \langle 0 \rangle$ (i.e., $a = 0$): Then

$$\begin{aligned} w^{s^+} \bigtriangleup w^{s^-} &= w^t \bigtriangleup r(\widehat{\psi}^{s^{-+}} \wedge \psi^{s^{--}}) \\ &= w^t \bigtriangleup r(\langle \psi^t(0) + 1 \rangle \wedge \psi^t \lceil [1, \ell(t)) \wedge \psi^t) \\ &= \{\psi^t(0) + 1\} \approx 1. \end{aligned}$$

Case 2 $s^{++} \in \mathcal{S}$ and $s^+ \neq s^{--} \wedge \langle 0 \rangle$ (i.e., $0 < b < |t|$).

Sub-case 2.1 $s^{-+} \notin \mathcal{S}$ (i.e., $a = |t|$):

$$\begin{aligned} w^{s^+} \bigtriangleup w^{s^-} &= r(\psi^{t^- \wedge \langle b \rangle} \wedge \psi^t) \bigtriangleup w^t \\ &= w^{t^- \wedge \langle b \rangle} \setminus w^t \preceq 1 \end{aligned}$$

by induction hypothesis applied to $s^+ = t \wedge \langle b \rangle$.

Sub-case 2.2 $s^{-+} \in \mathcal{S}$ and $s^- \neq s^{--} \wedge \langle 0 \rangle$ (i.e., $0 < a < |t|$): By the preceding corollary,

$$\begin{aligned} w^{s^+} \bigtriangleup w^{s^-} &= r(\psi^{t^- \wedge \langle b \rangle} \wedge \psi^t) \bigtriangleup r(\psi^{t^- \wedge \langle a \rangle} \wedge \psi^t) \\ &\subseteq w^{t^- \wedge \langle b \rangle} \bigtriangleup w^{t^- \wedge \langle a \rangle} = \emptyset, \end{aligned}$$

using $(A \cup C) \triangle (B \cup C) = (A \triangle B) \setminus C \subseteq A \triangle B$.

Sub-case 2.3 $s^{-+} \in \mathcal{S}$ and $s^- = s^{--} \wedge \langle 0 \rangle$ (i.e., $a = 0$): By the preceding lemma,

$$w^{s^+} \triangle w^{s^-} = w^{t^- \wedge \langle 0 \rangle} \triangle w^{t^- \wedge \langle b \rangle} \approx 1$$

by the lemma.

Case 3 $s^{++} \in \mathcal{S}$ and $s^+ = s^{--} \wedge \langle 0 \rangle$ (i.e., $b = 0$).

Sub-case 3.1 $s^{-+} \notin \mathcal{S}$ (i.e., $a = |t|$):

$$\begin{aligned} w^{s^+} \triangle w^{s^-} &= r(\widehat{\psi}^{s^{++}} \wedge \psi^{s^{+-}}) \triangle w^t \\ &= r(\langle \psi^{t^- \wedge \langle 0 \rangle}(0) + 1 \rangle \ \wedge \ \psi^{t^- \wedge \langle 0 \rangle} \lceil [1, \ell(t^- \wedge \langle 0 \rangle) \ \wedge \ \psi^t) \triangle w^t \\ &= \widehat{w}^{t^- \wedge \langle 0 \rangle} \triangle w^t \approx 1 \end{aligned}$$

by part (ii) of the preceding corollary.

Sub-case 3.2 $s^{-+} \in \mathcal{S}$ and $s^- \neq s^{--} \wedge \langle 0 \rangle$: This case is 'symmetric' to case 2.2

Sub-case 3.3 $s^{-+} \in \mathcal{S}$ and $s^- = s^{--} \wedge \langle 0 \rangle$: Then $s^+ = s^-$, and the result is immediate. •

(6.9) **Corollary** (Determination lemma) Let $s = t \wedge \langle a, b \rangle \in \mathcal{S}$. Then either

(i) $b = |t| + 1$ and ($s^+ \notin \mathcal{S}$ and) $w^s = w^{s^-}$; or
(ii) $0 < a, b \leq |t|$ and $w^{s^+} = w^{s^-}$; or
(iii) $a = 0 < b \leq |t|$ and $w^{s^+} \subset r(\psi^{s^-} \lceil [1, \ell^-))$; or
(iv) $b = 0 < a \leq |t|$ and $w^{s^-} \subset r(\psi^{s^+} \lceil [1, \ell^+))$, hence also $w^{s^-} \subset r(\widehat{\psi}^{s^+} \lceil [1, \ell^+))$, or
(v) $0 = a = b$ and $w^{s^+} = w^{s^-}$, and hence $\widehat{w}^{s^+} \lceil [1, \ell^+) = w^{s^-} \lceil [1, \ell^-)$.

Proof (i) Follows since in this case $\ell = \ell^-$ and $\psi^s = \psi^{s^-}$.

(ii) Follows by subcase 2.2 of the preceding lemma.

(iii) By induction over $|t|$: For $t = \langle 0 \rangle$ the only case to consider is $s = \langle 0, 0, 1 \rangle$. Then

$$\begin{aligned} \psi^{s^+} &= \psi^{\langle 0,1 \rangle} = \psi^{\langle 0 \rangle} = \langle 0 \rangle \\ \psi^{s^-} &= \psi^{\langle 0,0 \rangle} = \langle 1, 0 \rangle \end{aligned}$$

so $w^{s^+} = \{0\} \subset \{0\} = r(\psi^{s^-} \lceil [1, 2))$.

So suppose the result holds for all $u \in \mathcal{S}(\leq n)$ for $n \geq 1$ and let $t \in \mathcal{S}(n+1)$. Then

$$\begin{aligned} \psi^{s^+} &= \psi^{t \wedge \langle b \rangle}) = \begin{cases} \psi^t & \text{if } b = |t| \\ \psi^{t^- \wedge \langle b \rangle} \wedge \psi^t & \text{else} \end{cases} \\ \psi^{s^-} &= \psi^{t \wedge \langle 0 \rangle} = \widehat{\psi}^{t^- \wedge \langle 0 \rangle} \wedge \psi^t \end{aligned}$$

By inductive hypothesis, $w^{t^- \wedge \langle b \rangle} \subset r(\psi^{t^- \wedge \langle 0 \rangle} \lceil [1, \ell^{-+}))$, where $\ell^{-+} = \ell(s^{-+}) = \ell(t^- \wedge \langle 0 \rangle)$, so the result follows.

(iv) The argument is almost completely symmetric to the one for (iii) and proceeds again by induction over $|t|$.

For $t = \langle 0 \rangle$ the only case to consider is $s = \langle 0, 1, 0 \rangle$. Then

$$\begin{aligned} \psi^{s^+} &= \psi^{\langle 0,0 \rangle} = \langle 1, 0 \rangle \\ \psi^{s^-} &= \psi^{\langle 0,1 \rangle} = \psi^{\langle 0 \rangle} = \langle 0 \rangle \end{aligned}$$

so $w^{s^-} = \{0\} \subset \{0\} = r(\psi^{s^+}\lceil[1,2))$.

Supposing the result holds for all $u \in \mathcal{S}(\leq n)$ for $n \geq 1$ and let $t \in \mathcal{S}(n+1)$. Then

$$\psi^{s^-} = \psi^{t\wedge\langle a\rangle}) = \begin{cases} \psi^t & \text{if } a = |t| \\ \psi^{t^-\wedge\langle a\rangle} \wedge \psi^t & \text{else} \end{cases}$$
$$\psi^{s^+} = \psi^{t\wedge\langle 0\rangle} = \widehat{\psi}^{t^-\wedge\langle 0\rangle} \wedge \psi^t$$

By inductive hypothesis, $w^{t^-\wedge\langle a\rangle} \subset r(\psi^{t^-\wedge\langle 0\rangle}\lceil[1,\ell^{-+}))$, where $\ell^{-+} = \ell(s^{-+}) = \ell(t^- \wedge \langle 0\rangle)$, so the result follows.

(v) Follows again from subcase 3.3 of the preceding lemma. •

Note that all this implies that $w^{s\wedge\langle a\rangle} = w^{s\wedge\langle b\rangle}$ iff $a = b = 0$ or $0 < a, b \leq |s|$.

Corollary For $s = t \wedge \langle a, b\rangle \in \mathcal{S}$ such that $s^+ \in \mathcal{S}$ one of the following applies:

(i) if $a = b = 0$, then $\zeta^s = \zeta^{s^+} + 1 = \zeta^{s^-} + 1$ and $\xi^s = \xi^{s^+} + 1 = \xi^{s^-} + 1$;

(ii) if $a > 0 = b$, then $\zeta^s = \zeta^{s^+} = \zeta^{s^-} + 1$ and $\xi^s = \xi^{s^+} + 1 > \max\{\xi^{s^+}, \zeta^{s^-}, \}$;

(iii) if $a = 0 < b$, then $\zeta^s = \zeta^{s^-} = \zeta^{s^+} + 1$ and $\xi^s = \xi^{s^-} + 1 > \max\{\xi^{s^+}, \zeta^{s^-}, \}$;

(iv) if $a, b > 0$, then $\zeta^s = \zeta^{s^+} = \zeta^{s^-}$ and $\xi^s = \xi^{s^+} = \xi^{s^-}$.

For $s = t \wedge \langle a, b\rangle \in \mathcal{S}$ such that $s^+ i \notin \mathcal{S}$

(v) $\zeta^s = \zeta^{s^-}$ and $\xi^s = \xi^{s^-}$.

(6.10) **Corollary and Definition** (Free position, first form) Let $s = t \wedge \langle a, b\rangle \in \mathcal{S}(< \omega)$. Then one of the following applies:

(i) $a = b = 0$, so $s^+ = s^- = t \wedge \langle 0\rangle$, hence
$\psi^s = \widehat{\psi}^{t\wedge\langle 0\rangle} \wedge \psi^{t\wedge\langle 0\rangle}$,
$\widehat{\psi}^{t\wedge\langle 0\rangle} \triangle \psi^{t\wedge\langle 0\rangle} = \{\psi^{t\wedge\langle 0\rangle}(0) + 1, \psi^{t\wedge\langle 0\rangle}(0)\} = \{\psi^s(0), \psi^{s^-}(0)\} = \{\psi^s(0), \psi^{s^+}(0)\}$.
Set $\phi^s = \{0, \ell(s^+)\}$.

(ii) $a = 0 < b < |s| - 1$, so $s^+ = t \wedge \langle b\rangle$, $s^- = t \wedge \langle 0\rangle$, hence
$\psi^s = \psi^{t\wedge\langle b\rangle} \wedge \psi^{t\wedge\langle 0\rangle}$,
$\psi^{t\wedge\langle b\rangle} \triangle \psi^{t\wedge\langle 0\rangle} = \{\psi^{t\wedge\langle 0\rangle}(0)\} = \{\psi^s(\ell(s^+))\}$.
Set $\phi^s = \{\ell(s^+)\}$.

(iii) $a = 0 < b = |s| - 1$, so $s^+ \notin \mathcal{S}$, hence
$\psi^s = \psi^{s^-}$.
Set $\phi^s = \emptyset$.

(iv) $b = 0 < a \leq |s| - 2$, so $s^+ = t \wedge \langle 0\rangle$, $s^- = t \wedge \langle a\rangle$, hence
$\psi^s = \widehat{\psi}^{t\wedge\langle 0\rangle} \wedge \psi^{t\wedge\langle a\rangle}$,
$\widehat{\psi}^{t\wedge\langle 0\rangle} \triangle \psi^{t\wedge\langle a\rangle} = \{\psi^{t\wedge\langle 0\rangle}(0) + 1\} = \{\psi^s(0)\}$.
Set $\phi^s = \{\ell(s^+)\}$.

(v) $0 < a, b \leq |s| - 2$, so $s^+ = t \wedge \langle b\rangle$, $s^- = t \wedge \langle a\rangle$, hence
$\psi^s = \psi^{t\wedge\langle b\rangle} \wedge \psi^{t\wedge\langle a\rangle}$,
$\psi^{t\wedge\langle b\rangle} \triangle \psi^{t\wedge\langle a\rangle} = \emptyset$.
Set $\phi^s = \emptyset$.

(vi) $0 < a \leq |s| - 2 < b = |s| - 1$, so $s^+ \notin \mathcal{S}$, hence

$\psi^s = \psi^{s^-}$.

Set $\phi^s = \emptyset$. •

Contracting this somewhat:

(6.11) **Corollary** (Free position, second form) For $s = t \wedge \langle a, b \rangle \in \mathcal{S}$ such that $s^+ \in \mathcal{S}$ one of the following applies:

(i) $a = b = 0$, $\psi^s = \widehat{\psi}^{s^+} \wedge \psi^{s^-}$ and $\phi^s = \{0, \ell_n\}$.

(ii) $a > 0 = b$, $\psi^s = \widehat{\psi}^{s^+} \wedge \psi^{s^-}$ and $\phi^s = \{0\}$.

(iii) $a = 0 < b$, $\psi^s = \psi^{s^+} \wedge \psi^{s^-}$ and $\phi^s = \{\ell_n\}$.

(iv) $0 < a, b$, $\psi^s = \psi^{s^+} \wedge \psi^{s^-}$ and $\phi^s = \emptyset$.

For $s = t \wedge \langle a, b \rangle \in \mathcal{S}$ such that $s^+ \notin \mathcal{S}$ one of the following applies:

(v) $\psi^s = \psi^{s^-}$ and $\phi^s = \emptyset$. •

Summary

(i) Call $i \in \ell(s)$ *free* for s if $\psi^s(i) \notin r(\psi^s\lceil(\ell(s) \setminus \{i\}))$.

(ii) Call $i \in \ell(s)$ *free* for s *relative to* T (or simply free for T, when s is clear from the context), where T is a subset of $\ell = [0, \ell)$ not containing i, if $\psi^s(i) \notin r(\psi^s\lceil T)$. (Thus, 'free for s' is 'free for s relative to $\ell \setminus \{i\}$'.)

Thus, i is free over s_n just if $\psi_n(i) \neq \psi_n(j)$ for all $j \in \ell_n$, $j \neq i$, hence just if the element M_i^n is not forced by the ω-uniformity conditions to be isomorphic to any of the other elements in the nth array, $\langle M_i^{(n)} : i \in \ell_n \rangle$. Similarly, $i \in \mathcal{L}_n^+$ is free (over s_n) relative to $T = \mathcal{L}_n^-$ if $\psi_n(i) \neq \psi_n(j)$ for all $j \in \mathcal{L}_n^-$, i.e., just if the element M_i^n is not forced by the ω-uniformity condition to be isomorphic to any of the elements in, $\langle M_i^{(n)} : i \in \mathcal{L}_n^- \rangle$.

What the determination lemma shows is that

(i) if $s_{n+1}(n+1) = n+1$, then $\psi_{n+1}(i) = \psi_n(i)$, and the $M_i^{(n+1)}$ are determined at the previous level.

(ii) if $0 < s_{n+1}(n), s_{n+1}(n+1) \leq n-1$, then no $i \in \mathcal{L}_{n+1}^+$ is free (for s_{n+1} relative to $\mathcal{L}_{n+1}^-$ (and vice versa with the roles of '+' and '−' interchanged), so the $M_i^{(n+1)}$ in the 'plus' half of $\mathcal{L}_{n+1}$ are determined by the $M^{(n+1)}$ in the 'minus' half, and vice versa; but one of these halfs is already determined by the previous level, so none of them are 'new'.

(iii) if $0 = s_{n+1}(n) < s_{n+1}(n+1) \leq n-1$, then no $i \in \mathcal{L}_{n+1}^+$, (in the 'plus' half) is free relative to $\mathcal{L}_{n+1}^- \setminus \{\ell_{n+1}^+\}$ (i.e., the 'minus' half without its first element).

(iv) if $0 = s_{n+1}(n+1) < s_{n+1}(n) \leq n-1$, then no $i \in \mathcal{L}_{n+1}^-$, (in the 'minus' half) is free relative to $\mathcal{L}_{n+1}^+ \setminus \{0\}$ (i.e., the 'plus' half without its first element).

(v) if $0 = s_{n+1}(n+1) = s_{n+1}(n)$, then no $i \in \mathcal{L}_{n+1}^- \setminus \{\ell_{n+1}^+\}$ is free relative to $\mathcal{L}_{n+1}^+ \setminus \{0\}$ and vice versa.

[7] Short representations

In this section and the next we give the stream-lined representations promised earlier.

[7.1] Short representations of rank $\omega + \omega$

A 'new' array element in the approximation of $A \in \mathcal{U}(\omega + \omega)$ by the chain of components of some arbitrary (but fixed) 'basepoint' x in A, may only be introduced into the 'level' $n+1$ array of rank ω building blocks (i.e., the array used for the rank $\omega + n$ approximation $[x]^A_{\omega+n}$, $\langle \mathbf{M}_n \rangle_{n\in\omega}$), if $\phi_n = \phi^{s_n} \neq \emptyset$ (free position lemma), in which case the latter also specifies the possible positions of the 'new' element in the interval $[0, \ell_n)$. However, a 'new' element can appear in position $0 \in \phi_{n+1}$ only if $\tau_n > 0$, since for $\tau_n = 0$ the level n array $\mathbf{M}_n$ gets copied into the $n+1$ array $\mathbf{M}^+_{n+1}$. in which case, the 'new' element can therefore only appear in the 'other' half of the array, at position $\ell^+_{n+1} = \ell_n$. (Obviously, the 'new' element may happen to be isomorphic with an 'existing' one; the ν_n indicate the positions of ω components not necessarily isomorphic to any of $M^{(k)}_i$, $k \leq n$, $i < \ell_k$, hence potentially 'new', compared with the arrays for the components of rank $\leq \omega + n$ containing the 'basepoint')

Thus, the 'new' position at level $n+1$ (rank $\omega + n + 1$) in the construction of a rank $\omega + \omega$ chain is therefore unqiuely determined whenever it exists, and this will determine the 'canonical ordering' of the short array for such A.

(7.1) **Definition and Lemma** Let $\{s_n\}_n$ be derived from $\langle \sigma, \pi \rangle \in \mathcal{G}(\omega + 1)$ Set

$$\nu_0 := \{0\}.$$

Suppose $s_{n+1} = t_{n+1} \wedge \langle a_{n+1}, b_{n+1} \rangle$ such that $s^+_{n+1} \in \mathcal{S}$, i.e., $\tau_n < \omega$. Then one of the following applies, where we write ν_n for ν^{s_n}:

(i) $a_{n+1} = b_{n+1} = 0$; then $\psi_{n+1} = \widehat{\psi}^+_{n+1} \wedge \psi^-_{n+1}$. Set

$$\nu_{n+1} := \begin{cases} \{\ell^+_{n+1}\} = \{\ell_n\} & \text{if } \tau_n = 0 \\ \{0\} & \text{else (if } \tau_n = 1). \end{cases}$$

(ii) $a_{n+1} > b_{n+1} = 0$; then, again, $\psi_{n+1} = \widehat{\psi}^+_{n+1} \wedge \psi^-_{n+1}$. Set

$$\nu_{n+1} := \begin{cases} \emptyset & \text{if } \tau_n = 0 \\ \{0\} & \text{else (if } \tau_n = 1). \end{cases}$$

(By the determination lemma, $w^{t \wedge \langle a \rangle} \subset w^{t \wedge \langle 0 \rangle} \lceil [1, \ell(t \wedge \langle 0 \rangle))$.)

(iii) $a_{n+1} = 0 < b_{n+1}$; then $\psi_{n+1} = \psi^+_{n+1} \wedge \psi^-_{n+1}$. Set

$$\nu_{n+1} := \begin{cases} \{\ell^+_{n+1}\} = \{\ell_n\} & \text{if } \tau_n = 0 \\ \emptyset & \text{else (if } \tau_n = 1). \end{cases}$$

(iv) $a_{n+1}, b_{n+1} > 0$; then, again $\psi_{n+1} = \psi^+_{n+1} \wedge \psi^-_{n+1}$. Set

$$\nu_{n+1} := \emptyset. \bullet$$

Corollary

$$\nu_0 = \{0\} \qquad \text{(note, } \phi^{\langle 0\rangle} = \phi_0 = \{0\}\text{)}$$

$$\nu_{n+1} = \begin{cases} \emptyset & \text{if } \phi_{n+1} = \emptyset \\ \{0\} & \text{if } \phi_{n+1} \supseteq \{0\} \text{ and } \tau_n = 1 \\ \{\ell^+_{n+1}\} & \text{if } \phi_{n+1} \supseteq \{\ell^+_{n+1}\} \text{ and } \tau_n = 0 \end{cases}$$

Note that $\nu_0 \neq \emptyset$ and that $\nu_{n+1} = \emptyset$ whenever $\tau_n = \omega$. Set

$$H = \{n \in \omega : \nu_n \neq \emptyset\}$$

$$K = |H|$$

Let $\{\eta_k\}_{k<K}$ enumerate H, i.e.,

$$\eta_0 = 0$$

$$\eta_{k+1} = \min\{n > \eta_k : \nu_n \neq \emptyset\}.$$

By convention, if $K < \omega$, $\eta_k = \omega$, for all $k \geq K$. Set $\mathrm{supp}(\eta) = \{k : \eta_k < \omega\}$, and $\mathrm{ran}(\eta)' = \mathrm{ran}(\eta\lceil\mathrm{supp}(\eta)) = \mathrm{ran}(\eta) \cap \omega$. Observe that H η and K are determined by $\langle\sigma, \tau\rangle$.

(7.2) **Definition** (Short arrays – compressed version) Define the *(compressed) short array* for $A = A(\sigma, \pi, \langle\sigma^{(n)}_i, \pi^{(n)}_i\rangle_{n\in\omega, 0\leq i<\ell_n}) \in \mathcal{U}(\omega + \omega)$ by

$$U_k := U^{(\eta_k)}_{\nu_{\eta_k}}.$$

where the $U^{(n)}_k$, $k \in w_n$ are defined to satisfy $U^{(n)}_{\psi_n(i)} = M^{(n)}_i = A(\sigma^{(n)}_i, \pi^{(n)}_i)$ via the ω-uniformity conditions. (Here, by abuse of notation, $U^{(n)}_{\{k\}} = U^{(n)}_k$.)

Expanding this,

$$U_0 = U^{(0)}_0$$

$$U_{k+1} = \begin{cases} U^{(\eta_{k+1})}_{\psi_{\eta_{k+1}}(\ell^+_{\eta_{k+1}})} & \text{if } \tau_{\eta_{k+1}-1} = 0 \\ U^{(\eta_{k+1})}_{\psi_{\eta_{k+1}}(0)} & \text{if } \tau_{\eta_{k+1}-1} = 1 \end{cases}$$

which in turn gives

$$U_0 = M^{(0)}_0,$$

$$U_{k+1} = \begin{cases} M^{(\eta_{k+1})}_{\ell^+_{\eta_{k+1}}} = M^{(\eta_{k+1})}_{\ell_{\eta_{k+1}-1}} & \text{if } \tau_{\eta_{k+1}-1} = 0 \\ M^{(\eta_{k+1})}_0 & \text{if } \tau_{\eta_{k+1}-1} = 1 \end{cases}$$

Define a left-inverse of η on $\mathrm{ran}(\eta)'$, by setting

$$\kappa_n = \max\{k : \eta_k \leq n\}.$$

Thus, $\kappa_n = k$ whenever n is the kth integer such that $\nu_n \neq \emptyset$. Note that $K = \mathrm{ran}(\kappa) = \mathrm{supp}(\eta)$.

Thus:

(i) The ν_{n+1} specify where at the level $n+1$ array a potentially ’new’ element $U_{\kappa_n} = M \in \mathcal{U}(\omega)$ $(\kappa_n \leq n)$ can get ’added’ to the array of building blocks — either as $M_0^{(n+1)}$ if $\tau_n = 1$, or as $M^{(n+1)}_{\ell^+_{n+1}} = M^{(n+1)}_{\ell_n}$ if $\tau_n = 0$.

(ii) Furthermore, level n array $\mathbf{M}_n$ gets ’copied’ as $\mathbf{M}^-_{n+1}$ if $\tau_n = 1$, and as $\mathbf{M}^+_{n+1}$ if $\tau_n = 0$.

(iii) The remainder of the level $n+1$ array (i.e., the blocks not in the ’copied from level n’-half — hence in the ’half’ containing the ’new’ position) is determined by the uniformity condition from the level $\omega + n$ array $\mathbf{M}_n$.

It remains to lay out the details on how the full double array $\langle \mathbf{M}_n \rangle_n$ is derived from the short array $\langle U_k \rangle_{k \in K}$ and vice versa.

Remark For all $k \in \operatorname{ran}(\kappa) = \operatorname{dom}(\eta)$,

(i) $\tau_{\eta_k} < \omega$.

(ii) $\kappa_{\eta_k} = k$;

(iii) $\kappa_m \leq k$ iff $m \leq \eta_k$;

(iv) if $k+1 < \operatorname{ran}(\kappa)$, then $\kappa_m = k$ iff $m \in [\eta_k, \eta_{k+1})$,
if $k+1 = \operatorname{ran}(\kappa) < \omega$, then $\kappa_m = k$ iff $m \in [\eta_k, \omega)$.

Proof (i) holds, since by definition (e.g., (free position lemma,(vi)). $\{n : \nu_n \neq \emptyset\} \subseteq \{n : \tau_n < \omega\}$. (ii) follows from (iii), and (iii) follows from (iv), so we show (iv): But (iv) is true for $k = 0$ $(\eta_0 = 0 = \kappa_0)$. Supposing (iv) true for all $l \leq k$, we get $\kappa_m = \kappa_{\eta_k} = k$ for $m \in [\eta_k, \eta_{k+1})$ if $k+1 \in \operatorname{ran}(\kappa)$, while $\kappa_m = \kappa_{\eta_k} = k$ for $m \in [\eta_k, \omega)$, if $k+1 = \operatorname{ran}(\kappa) < \omega$. •

(7.3) **Lemma**

(i) For all n, $\xi_{n+1} > \xi_n$ iff $\zeta_{n+1} = \zeta_n + 1$ iff $\nu_{n+1} \neq \emptyset$.

(ii) For all $n \in \operatorname{ran}(\eta)'$, $\kappa_n = \zeta_n - 1$ and so $\operatorname{dom}(\eta) = \operatorname{ran}(\kappa) = \sup_{n \in \omega} \zeta_n - 1$;

(iii)
$$\operatorname{dom}(\eta) = \operatorname{ran}(\kappa) = \begin{cases} \zeta_n & \text{if } n \text{ is minimal such that for all } m \geq n,\ \nu_m = \emptyset \\ \omega & \text{if } \nu_n \neq \emptyset \text{ for infinitely many } n, \end{cases}$$

Proof (i) Fix n and set $s_{n+1} = s \wedge \langle a, b \rangle$. We may assume $\tau_n < \omega$ as $\tau_n = \omega$ implies that $\xi_{n+1} = \xi_n$ and $\nu_{n+1} = \emptyset$. Thus, now, $s^+_{n+1} = s \wedge \langle b \rangle$ and $s^-_{n+1} = s \wedge \langle a \rangle$. From the definition $\xi_{n+1} > \xi_n$ iff $\zeta_{n+1} = \zeta_n + 1$.

Set $w^\pm_n = w^{s^\pm_n}$ and $\xi^\pm_n = \xi^{s^\pm_n}$.

Case $a = b = 0$**.** Then $s_{n+1} = s \wedge \langle 0, 0 \rangle$, and $w^+_{n+1} = w^{s \wedge \langle 0 \rangle} = w^-_{n+1}$. and $\xi^+_{n+1} = \xi^{s \wedge \langle 0 \rangle} = \xi^-_{n+1}$.

Subcase $\tau_n = 0$**.** Then
$s_n = s^+_{n+1} = s^-_{n+1}$,
$w_{n+1} = \widehat{w}^+_{n+1} \cup w^-_{n+1} = \widehat{w}^{s \wedge \langle 0 \rangle} \cup w^{s \wedge \langle 0 \rangle} = \widehat{w}_n \cup w_n = \{\psi_n(0) + 1\} \cup w_n \supset w_n$,
$\nu_{n+1} = \{\ell^+_{n+1}\} = \{\ell_n\}$.

Subcase $\tau_n = 1$**.** Then
$s_n = s^-_{n+1} = s^+_{n+1}$,
$w_{n+1} = \{\psi_n(0) + 1\} \cup w_n \succ w_n$ (as before),
$\nu_{n+1} = \{0\}$.

Case $a > b = 0$**.** Then $s_{n+1} = s \wedge \langle a, 0\rangle$, and $w^+_{n+1} = w^{s\wedge\langle 0\rangle} \supset w^-_{n+1}$.

Subcase $\tau_n = 0$**.** Then

$s_n = s^+_{n+1} = s \wedge \langle 0\rangle$,

$\zeta_{n+1} = |\widehat{w}^+_{n+1} \cup w^-_{n+1}| = |\widehat{w}^{s\wedge\langle 0\rangle} \cup w^{s\wedge\langle a\rangle}| = |w^{s\wedge\langle 0\rangle} \setminus \{\psi^{s\wedge\langle 0\rangle}(0)\} \cup \{\widehat{\psi}^{s\wedge\langle 0\rangle}(0)\} \cup w^{s\wedge\langle a\rangle}| = |w^{s\wedge\langle 0\rangle} \cup w^{s\wedge\langle a\rangle}| = \zeta^{s\wedge\langle 0\rangle} = \zeta_n$ (since $\{\psi^{s\wedge\langle 0\rangle}(0), \psi^{s\wedge\langle 0\rangle}(0)+1\} \cap w^{s\wedge\langle a\rangle} = \emptyset$ and $w^{s\wedge\langle 0\rangle} \supset w^{s\wedge\langle a\rangle}$),

$\nu_{n+1} = \emptyset$.

Subcase $\tau_n = 1$**.** Then

$s_n = s^-_{n+1} = s \wedge \langle a\rangle$,

$\zeta_{n+1} = |w^{s\wedge\langle 0\rangle} \cup w^{s\wedge\langle a\rangle}| > |w^{s\wedge\langle a\rangle}| = \zeta_n$ (following the preceding case),

$\nu_{n+1} = \{0\}$.

Case $a = 0 < b$**.** Then $s_{n+1} = s \wedge \langle 0, b\rangle$,

Subcase $\tau_n = 0$**.** Then

$s_n = s^+_{n+1} = s \wedge \langle b\rangle$,

$\zeta_{n+1} = |w^+_{n+1} \cup w^-_{n+1}| = |w^{s\wedge\langle b\rangle} \cup w^{s\wedge\langle 0\rangle}| = |w^{s\wedge\langle 0\rangle}| > |w^{s\wedge\langle b\rangle}| = \zeta_n$,

$\nu_{n+1} = \{\ell^+_{n+1}\} = \{\ell_n\}$.

Subcase $\tau_n = 1$**.** Then

$s_n = s^-_{n+1} = s \wedge \langle 0\rangle$,

$\zeta_{n+1} = |w^{s\wedge\langle 0\rangle}| = \zeta_n$ (again, following the preceding case),

$\nu_{n+1} = \emptyset$.

Case $a, b > 0$**.** In that case $w^{s\wedge\langle a\rangle} = w^{s\wedge\langle b\rangle}$ by the plus one range corollary, so $\xi_{n+1} = \xi^{s\wedge\langle a,b\rangle} = \xi^{s\wedge\langle a\rangle} = \xi^{s\wedge\langle b\rangle} = \xi_n$, but also $\nu_{n+1} = \emptyset$.

(ii) We have $s_0 = \langle 0\rangle$, $w^{s\langle 0\rangle} = \{0\}$ and $\kappa_0 = 0$. By the preceding remark, part (iii), $\kappa_{n+1} = \kappa_n$ if $\nu_{n+1} = \emptyset$, else $\kappa_{n+1} = \kappa_n + 1$, hence $\kappa_{n+1} = \kappa_n$ if $\zeta_{n+1} = \zeta_n$, else $\kappa_{n+1} = \kappa_n + 1$ and $\zeta_{n+1} = \zeta_n + 1$.

(iii) Follows from (ii). •

The preceding lemma permits the following alternative indexing of the short array, which will be exploited in the next section.

7.4 **Definiton** (Short arrays – uncompressed version) Define the *uncompressed short array* $\langle V_k\rangle_{k\in X}$ for $A \in \mathcal{U}(\omega+\omega)$ as above by

$$V_{\xi_{\eta_k}} = U^{(\eta_k)}_{\nu_{\eta_k}},$$

where $X = \{\xi_n : n \in H'\}$.

Remark The double arrays $\langle \mathbf{M}^{(n)}\rangle_{n\in\omega} = \langle M^{(n)}_i\rangle_{i\in\mathcal{L}_n, n\in\omega}$ and $\langle \mathbf{U}^{(n)}\rangle_{n\in\omega} = \langle U^{(n)}_k\rangle_{k\in w_n, n\in\omega}$, and the short arrays $\mathbf{U} = \langle U_k\rangle_{k<K}$ and $\mathbf{V} = \langle V_k\rangle_{k\in X} \subseteq \mathcal{U}(\omega)$ deriving from a chain $A \in \mathcal{U}(\omega+\omega)$ as given above solve the following set of equations:

$$M^{(0)}_0 = U^{(0)}_0 = U_0 = V_0$$

$$U_{\kappa_{n+1}} = V_{\xi_{n+1}} = \begin{cases} M^{(n+1)}_{\ell_n} = U^{(n+1)}_{\psi_{n+1}(\ell_n)} & \text{if } \nu_{n+1} \neq \emptyset \text{ and } \tau_n = 0 \\ M^{(n+1)}_0 = U^{(n+1)}_{\psi_{n+1}(0)} & \text{if } \nu_{n+1} \neq \emptyset \text{ and } \tau_n = 1 \end{cases}$$

and

$$\mathbf{M}_{n+1} = \begin{cases} \mathbf{M}^{(n)} \ \wedge \ \langle U_{\kappa_{n+1}} \rangle \ \wedge \ \mathbf{M}^{(n+1)}_{(\ell_n, \ell_{n+1})} & \text{if } \nu_{n+1} \neq \emptyset \text{ and } \tau_n = 0, \\ \langle U_{\kappa_{n+1}} \rangle \ \wedge \ \mathbf{M}^{(n+1)}_{(0, \ell^+_{n+1})} \ \wedge \ \mathbf{M}^{(n)} & \text{if } \nu_{n+1} \neq \emptyset \text{ and } \tau_n = 1. \end{cases}$$

Note that we may pass from the uncompressed to the compressed short array by applying the order-preserving mapping ('collapse') from X to K, $\xi_n \mapsto \kappa_n$ (effectively, a bounded search operation).

Note that generally $V_{\xi_{n+1}}$ and $M^{(n+1)}_{\xi_{n+1}} = U^{(n+1)}_{\xi_{n+1}}$ refer to elements in different positions of the level $n+1$ array (and may differ).

Call $\mathbf{M}^{(+)} \in {}^{\ell^+}\mathcal{U}(\omega)$ and $\mathbf{M}^{(-)} \in {}^{\ell^-}\mathcal{U}(\omega)$ *s-compatible* if $\mathbf{M} = \mathbf{M}^{(+)} \wedge \mathbf{M}^{(-)}$ satisfies the uniformity condition for s.

(7.5) **Lemma** $\mathbf{M}^{(+)} \in {}^{\ell^+}\mathcal{U}(\omega)$ and $\mathbf{M}^{(-)} \in {}^{\ell^-}\mathcal{U}(\omega)$ are s-compatible if the following apparently weaker condition holds:

$$(\forall i < \ell^+)(\forall k < \ell^-)[\psi^s(i) = \psi^s(\ell^+ + k) \ \rightarrow \ M^{(+)}_i \cong M^{(-)}_k], \tag{7.1}$$

where $\mathbf{M}^{(\pm)} = \langle M^{(\pm)}_k \rangle_{k < \ell^\pm}$, and where, as usual, $\ell = \ell(s)$, $\ell^\pm = \ell(s^\pm)$.

Proof For the 'easy' direction note that for $i \in \mathcal{L}^+$ and $j = \ell^+ + k \in \mathcal{L}^-$, if $\psi^s(i) = \psi^s(j)$, then by the uniformity conditions, $M^{(+)}_i = M_i \cong M_j = M^{(-)}_k$, as required.

So suppose $\mathbf{M} = \mathbf{M}^{(+)} \wedge \mathbf{M}^{(-)}$ satisfies condition (7.1) for s and let $i, j \in \mathcal{L}$. We have to show that either $\psi^s(i) \neq \psi^s(j)$ or that $M_i \cong M_j$.

Fix $t \in \mathcal{S}$ and $a, b \leq |t|$ such that $s = t \wedge \langle a, b \rangle$. Since s^+ is assumed in $\mathcal{S}$, either $b = 0$ and $\psi^s = \widehat{\psi}^{s^+} \wedge \psi^{s^-}$ or $b > 0$ and $\psi^s = \psi^{s^+} \wedge \psi^{s^-}$.

Thus, if $0 < i < \ell^+$ and $j = \ell^+ + k \in \mathcal{L}^-$, then by assumption $\psi(i) = \psi(j)$ implies that $M_i = M^{(+)}_i \cong M^{(-)}_k = M_j$. Since we may assume $i < j$, we are left with the cases

(i) $i = 0$,$j < \ell$,

(ii) $0 < i, j < \ell^+$, and

(iii) $\ell^+ \leq i, j < \ell$.

Case $i = 0$:

Subcase $b = 0$ and $\psi^s = \widehat{\psi}^{s^+} \wedge \psi^{s^-}$: Then for $k \leq \ell^-$ and $j = \ell^+ + k$,

$$\psi(0) = \psi^+(0) + 1 = \psi^{t \wedge \langle 0 \rangle}(0) + 1 = |t| + 1 > \psi^{t \wedge \langle a \rangle}(k) = \psi^{s^-}(k) = \psi^s(j),$$

by weight sequence lemma. Likewise, for $0 < j' < \ell^+$,

$$\psi(0) = \psi^+(0) + 1 = \psi^{t \wedge \langle 0 \rangle}(0) + 1 = |t| + 1 > \psi^{t \wedge \langle 0 \rangle}(j') = \psi^{s^+}(j') = \psi^s(j').$$

(Thus in this case the antecedent $\psi(0) = \psi(j)$ never applies, except for $0 = j$, when $M_0 \cong M_j$ holds trivially.)

Subcase $b > 0$ and $\psi^s = \psi^{s^+} \wedge \psi^{s^-}$: Thus, for $k \leq \ell^-$, $j = \ell^+ + k$

$$\psi^s(0) = \psi^{s^+}(0) = \psi^{t \wedge \langle b \rangle}(0),$$
$$\psi^s(j) = \psi^{s^-}(k) = \psi^{t \wedge \langle a \rangle}(k),$$

So if $\psi^s(0) = \psi^s(j)$, then by (7.1)

$$M_0 = M_0^{(+)} \cong M_k^{(-)} = M_j,$$

as required.

For $0 < j' < \ell^+$,

$$\psi^s(j') = \psi^{s^+}(j') = \psi^{t \wedge \langle b \rangle}(j').$$

By the plus one range lemma,

$$w^{s^+} = w^{t \wedge \langle b \rangle} = w^{t \wedge \langle b \rangle} = w^{s^-},$$

so fix $k'' \in \ell^-$ such that $\psi^{s^+}(0) = \psi^{s^-}(k'')$ and let $j'' = \ell^+ + k''$, so $\psi^s(j') = \psi^s(0) = \psi^{s^+}(0) = \psi^{s^-}(k'') = \psi^s(j'')$. So if $\psi^s(0) = \psi^s(j')$, then in fact $\psi^s(0) = \psi^s(j'') = \psi^s(j')$, so by (7.1) (applied twice to the second hypothesis),

$$M_0 = M_0^{(+)} \cong M_{k''}^{(-)} \cong M_{k'}^{(+)} = M_{j'},$$

as required.

Case $0 < i, j < \ell^+$: Then

$$\psi^s(i) = \psi^{s^+}(i) = \psi^{t \wedge \langle 0 \rangle}(i),$$
$$\psi^s(j) = \psi^{s^+}(j) = \psi^{t \wedge \langle 0 \rangle}(j),$$

Again, by plus one range lemma,

$$\psi^{t \wedge \langle 0 \rangle}(i) = \psi^{t \wedge \langle 0 \rangle}(j) \in w^{t \wedge \langle a \rangle},$$

so fix $k < \ell^-$ such that $\psi^{t \wedge \langle 0 \rangle}(i) = \psi^{t \wedge \langle 0 \rangle}(j) = \psi^{t \wedge \langle a \rangle}(k)$. By (7.1),

$$M_i = M_i^{(+)} = M_k^{(-)} = M_j^{(+)} = M_j.$$

Case $\ell^+ \leq i, j < \ell$. Then

$$\psi^s(i) = \psi^{s^-}(i) = \psi^{t \wedge \langle a \rangle}(i),$$
$$\psi^s(j) = \psi^{s^-}(j) = \psi^{t \wedge \langle a \rangle}(j),$$

Again, by plus one range lemma,

$$\psi^{t \wedge \langle a \rangle}(i) = \psi^{t \wedge \langle a \rangle}(j) \in w^{t \wedge \langle b \rangle},$$

so fix $k < \ell^+$ such that $\psi^{t \wedge \langle a \rangle}(i) = \psi^{t \wedge \langle a \rangle}(j) = \psi^{t \wedge \langle b \rangle}(k)$. By two applications of(7.1),

$$M_i = M_i^{(-)} = M_k^{(+)} = M_j^{(-)} = M_j. \bullet$$

Make the obvious modifications to the above terminology when dealing with 'shifted' or truncated sequences. E.g.: Call $\mathbf{M}^{(+)} \in {}^{\mathcal{L}(s)^+}\mathcal{U}(\omega)$ and $\mathbf{M}^{(-)} \in {}^{\mathcal{L}(s)^-}\mathcal{U}(\omega)$ *s-compatible* if $\mathbf{M}^{(+)}$ and $\widetilde{\mathbf{M}} \in {}^{\ell(s^-)}\mathcal{U}(\omega)$ are compatible, where $\widetilde{M_i} = M_{i-\ell^+}^{(-)}$.

Let $\mathcal{J}^\pm \subseteq \mathcal{L}^\pm$. Call $\mathbf{M}^{(+)} \in {}^{\mathcal{J}^+}\mathcal{U}(\omega)$ and $\mathbf{M}^{(-)} \in {}^{\mathcal{J}^-}\mathcal{U}(\omega)$ *s-compatible* if

$$(\forall i \in \mathcal{J}^+)(\forall j \in \mathcal{J}^-)[\psi^s(i) = \psi^s(j) \ \rightarrow \ M_i^{(+)} \cong M_j^{(-)}],$$

i.e., if $\mathbf{M} = \widetilde{\mathbf{M}}^{(+)} \wedge \widetilde{\mathbf{M}}^{(-)}$ satisfies the uniformity conditions for s restricted to $\mathcal{J}^+ \cup \mathcal{J}^-$, where $\widetilde{\mathbf{M}}^{(\pm)}$ are arbitrary extensions of $\mathbf{M}^{(\pm)}$ onto $\mathcal{J}^\pm$. Etc.

Let $\mathcal{J} \subseteq \mathcal{L}^+$. Call $\widetilde{\mathbf{M}} \in {}^{\mathcal{L}^+}\mathcal{U}(\omega)$ *s-determinable on* $\mathcal{J}$ from $\mathbf{M}^{(-)} \in {}^{\mathcal{L}^-}\mathcal{U}(\omega)$, if $r(\psi^+\lceil\mathcal{J}) \subseteq w^-$, and for all $i \in \mathcal{J}$, $\widetilde{\mathbf{M}}_i = \mathbf{M}^{(-)}_j$ for *every* j such that $\psi^+(i) = \psi^-(j)$. Likewise for $\mathcal{J} \subseteq \mathcal{L}^-$. Call $\widetilde{\mathbf{M}} \in {}^{\mathcal{L}^-}\mathcal{U}(\omega)$ *s-determinable on* $\mathcal{J}$ from $\mathbf{M}^{(+)} \in {}^{\mathcal{L}^+}\mathcal{U}(\omega)$, if $r(\psi^-\lceil\mathcal{J}) \subseteq w^+$, and for all $i \in \mathcal{J}$, $\widetilde{\mathbf{M}}_i = \mathbf{M}^{(-)}_j$ for *every* j such that $\psi^-(i) = \psi^+(j)$. If an $\widetilde{\mathbf{M}}$ is given as $\widetilde{\mathbf{M}} = \widetilde{\mathbf{M}}_{\mathcal{J}}$ for $\mathcal{J} \subseteq \mathcal{L}^+$ or $\mathcal{J} \subseteq \mathcal{L}^-$, we shall simply say $\mathbf{M}_{\mathcal{J}}$ is s-determinable from $\mathbf{M}^{(-)}$, respectively $\mathbf{M}^{(+)}$. If, e.g., $\mathbf{M}^{(-)} \in {}^{\mathcal{L}^-}\mathcal{U}(\omega)$, satisfies the uniformity conditions (on $\mathcal{L}^-$), then it suffices for $\widetilde{\mathbf{M}} \in {}^{\mathcal{L}^+}\mathcal{U}(\omega)$ to be s-determinable on $\mathcal{J}$ from $\mathbf{M}^{(-)}$ that $r(\psi^+\lceil\mathcal{J}) \subseteq w^-$, and for all $i \in \mathcal{J}$, $\widetilde{\mathbf{M}}_i = \mathbf{M}^{(-)}_j$ there is *some* j such that $\psi^+(i) = \psi^-(j)$.

(7.6) **Lemma** (Array equivalence lemma) Let $A = A(\sigma, \pi)[\mathbf{M}_n]_{n\in\omega} \in \mathcal{U}(\omega + \omega)$. Then $\mathbf{M}_0 = \langle U_0 \rangle$ and for each n, one of (i), (ii) or (iii) applies:

(i) **Case** $\tau_n = \omega$:

Then $\mathbf{M}_{n+1} = \mathbf{M}_n$.

(ii) **Case** $\tau_n = 1$:

Subcase $b_{n+1} = s_{n+1}(n+1) = 0$:

Then for $\widetilde{\mathbf{M}}_{(0,\ell^+_{n+1})}$ s_{n+1}-determinable from $\mathbf{M}_n$,

$$\mathbf{M}_{n+1} = \langle U_{\kappa_{n+1}} \rangle \ \wedge \ \widetilde{\mathbf{M}}_{(0,\ell^-_{n+1})} \ \wedge \ \mathbf{M}_n.$$

Subcase $b_{n+1} > 0$.

Then $\mathbf{M}_{n+1}$ is determined by $\mathbf{M}_n$ in the sense that

$$\mathbf{M}^-_{n+1} = \mathbf{M}_n \text{ and } \mathbf{M}^+_{n+1} \text{ is } s_{n+1}\text{-determinable from } \mathbf{M}^-_{n+1}.$$

(iii) **Case** $\tau_n = 0$:

Subcase $a_{n+1} = s_{n+1}(n) = 0$:

Then for $\widetilde{\mathbf{M}}_{(0,\ell^-_{n+1})}$ s_{n+1}-determinable from $\mathbf{M}_n$,

$$\mathbf{M}_{n+1} = \mathbf{M}_n \ \wedge \ \langle U_{\kappa_{n+1}} \rangle \ \wedge \ \widetilde{\mathbf{M}}_{(0,\ell^-_{n+1})}.$$

Subcase $a_{n+1} > 0$.

Then $\mathbf{M}_{n+1}$ is determined by $\mathbf{M}_n$ in the sense that

$$\mathbf{M}^+_{n+1} = \mathbf{M}_n \text{ and } \mathbf{M}^-_{n+1} \text{ is } s_{n+1}\text{-determinable from } \mathbf{M}^+_{n+1}$$

Proof The proof consists merely in pulling together earlier observations and contains nothing new. By definition $M^{(0)}_0 = U_0 = V_0$.

If $\tau_n = \omega$, then $s_{n+1} = s_n$, and per force $\mathbf{M}_{n+1} = \mathbf{M}_n$.

So suppose for the remainder that $\tau_n \in \omega$. Let $s = s_{n+1} = t \wedge \langle a, b\rangle$ (and $\ell = \ell_{n+1} = \ell(s_{n+1})$, $\ell^{\pm} = \ell^{\pm}_{n+1} = \ell(s^{\pm}_{n+1})$, etc.).

Case $a = b = 0$. Thus $\nu_n \subseteq \{0, \ell^+\}$, and that $\nu_n \neq \emptyset$. Thus, let $k + 1 = \kappa_{n+1}$ and $l + 1 = \xi_{n+1}$ and note that $\kappa_{n+1} > k \geq \kappa_m$ for $m \leq n$, $\xi_{n+1} > l \geq \xi_m$ for $m \leq n$ so that $U_{k+1} = V_{l+1}$ has not already been 'used' at any previous level in the 'approximation' to A by the chain of components $A(s_n)[\mathbf{M}_n]$. Note also that now $s^+ = t \wedge \langle 0\rangle = s^-$, $\ell^+ = \ell_n = \ell^-$, etc.

Subcase $\tau_n = 0$: Then $\nu_n = \{\ell^+\}$ and

$$\mathbf{M}_{n+1} = \mathbf{M}_n \ \wedge\ \langle U_{k+1}\rangle \ \wedge\ \widetilde{\mathbf{M}}_{(0,\ell^-_{n+1})},$$

where $\widetilde{\mathbf{M}}_{(0,\ell^-_{n+1})}$ is s_{n+1} determinable from $\mathbf{M}_n$ (since $w^+_{n+1} \setminus \{\psi_{n+1}(0)\} = w^{t\wedge\langle 0\rangle} \setminus \{\psi^{t\wedge\langle 0\rangle}(0)\} = w^-_{n+1} \setminus \{\psi_{n+1}(\ell_n)\}$).

Subcase $\tau_n = 1$: Then $\nu_n = \{0\}$ and

$$\mathbf{M}_{n+1} = \langle U_{k+1}\rangle \ \wedge\ \widetilde{\mathbf{M}}_{(0,\ell^+_{n+1})} \ \wedge\ \mathbf{M}_n,$$

where $\widetilde{\mathbf{M}}_{(0,\ell^+_{n+1})}$ is s_{n+1} determinable from $\mathbf{M}_n$ (for reasons as before).

Case $a > b = 0$. Thus $\nu_n \subseteq \{0\}$ and that $\nu_n = \emptyset$ iff $\tau_n = 0$. Also, $w^{s^-} = w^{t\wedge\langle a\rangle} \subseteq r(\psi^{t\wedge\langle 0\rangle}\lceil[1, \ell(t \wedge \langle 0\rangle)) = r(\psi^{s^+}\lceil[1, \ell^+))$.

Subcase $\tau_n = 0$: Then $s^+ = s_n$, $\ell^+ = \ell_n$ and $\nu_n = \emptyset$. By the preceding, $w^{s^-} \subseteq r(\psi_n\lceil[1, \ell_n))$, so every entry $\widetilde{M}^{n+1}_i, i < \ell^-$ is determined (up to isomorphism) an entry $M^{(n+1)}_i, i \in \mathcal{L}^+_{n+1} \setminus \{0\}$ via s_{n+1}-compatibility. Thus,

$$\mathbf{M}_{n+1} = \mathbf{M}_n \ \wedge\ \widetilde{\mathbf{M}}_{\mathcal{L}^-}$$

where the array $\widetilde{\mathbf{M}}_{\mathcal{L}^-}$ is s_{n+1}-determinable from $\mathbf{M}_n$ (since $w^{s^-} \subseteq r(\psi^{s^+}\lceil[1, \ell^+))$, implies that no position in $\mathcal{L}^- = [\ell^+, \ell)$ is 'free' relative to $\mathcal{L}^+ = [0, \ell^+)$.)

Subcase $\tau_n = 1$: Then $s^- = s_n$, $\ell^- = \ell_n$ and $\nu_n = \{0\}$. Let $k + 1 = \kappa_{n+1}$ and $l + 1 = \xi_{n+1}$ and note again that $U_{k+1} = V_{l+1}$ has not already been 'used' at any previous 'level' in the approximation of A. So,

$$\mathbf{M}_{n+1} = \langle U_{k+1}\rangle \ \wedge\ \widetilde{\mathbf{M}}_{(0,\ell^+_{n+1})} \ \wedge\ \mathbf{M}_n$$

where the array $\widetilde{\mathbf{M}}_{\mathcal{L}^+}$ is s_{n+1}-determinable from $\mathbf{M}_n$ (since $\widehat{w}^{s^+} \setminus w^{s^-} = \{\psi^s(0)\}$, i.e., no position in $(0, \ell^+)$ being 'free' relative to $\mathcal{L}^- = [\ell^+, \ell)$).

Case $a = 0 < b$. Thus $\nu_n \subseteq \{\ell^+\}$ and that $\nu_n = \emptyset$ iff $\tau_n = 1$. Also, $w^{s^+} = w^{t\wedge\langle b\rangle} \subseteq r(\psi^{t\wedge\langle 0\rangle}\lceil[1, \ell(t \wedge \langle 0\rangle))) = r(\psi^{s^-}\lceil[1, \ell^-)))$.

Subcase $\tau_n = 0$: Then $s^+ = s_n$, $\ell^+ = \ell_n$ and $\nu_n = \{\ell^+\}$. Let $k + 1 = \kappa_{n+1}$, and $l + 1 = \xi_{n+1}$, and note again that $U_{k+1} = V_{l+1}$ is not already 'used' at any previous 'level'. Also, $w_n = r(\psi^-\lceil[1, \ell^-)$, so all the entries $\widetilde{M}^{n+1}_i, i \in \mathcal{L}^- \setminus \{\ell^+_{n+1}\}$ are determined (up to isomorphism) by the entries in $M^{(n+1)}_i, i < \ell^+_{n+1}$ via s_{n+1}-compatibility. Thus,

$$\mathbf{M}_{n+1} = \mathbf{M}_n \ \wedge\ \langle U_{k+1}\rangle \ \wedge\ \widetilde{\mathbf{M}}_{(\ell^+_{n+1},\ell^-_{n+1})}$$

where the array $\widetilde{\mathbf{M}}_{(\ell^+_{n+1},\ell_{n+1})}$ is s_{n+1}-determinable from $\mathbf{M}_n$.

Subcase $\tau_n = 1$: Then $s^- = s_n$, $\ell^- = \ell_n$ and $\nu_n = \emptyset$. Now, $w^{s^+} = w^{t\wedge\langle b\rangle} \subseteq w^{t\wedge\langle 0\rangle} = w^{s^-}$, so all the entries $\widetilde{M}_i^{n+1}, i < \ell^+$ are determined (up to isomorphism) by the entries in $M_i^{(n+1)}, i \in \mathcal{L}_{n+1}^-$ via s_{n+1}-compatibility. So,

$$\mathbf{M}_{n+1} = \widetilde{\mathbf{M}}_{\mathcal{L}^+} \ \wedge\ \mathbf{M}_n$$

where the array $\widetilde{\mathbf{M}}_{\mathcal{L}^+}$ is s_{n+1}-determinable from $\mathbf{M}_n$.

Case $0 < a, b$. Thus $\nu_n = \emptyset$ but also $w^{s^+} = w^{s^-}$, which implies that each 'half' of $\mathbf{M}_{n+1}$ is determined by the other via s_{n+1} compatibility.

Subcase $\tau_n = 0$: Then $s^+ = s_n$, $\ell^+ = \ell_n$, and

$$\mathbf{M}_{n+1} = \mathbf{M}_n \ \wedge\ \widetilde{\mathbf{M}}_{\mathcal{L}^-}$$

where the array $\widetilde{\mathbf{M}}_{\mathcal{L}^-}$ is s_{n+1}-determinable from $\mathbf{M}_n$.

Subcase $\tau_n = 1$: Then $s^- = s_n$, $\ell^- = \ell_n$, and

$$\mathbf{M}_{n+1} = \widetilde{\mathbf{M}}_{\mathcal{L}^+} \ \wedge\ \mathbf{M}_n$$

where the array $\widetilde{\mathbf{M}}_{\mathcal{L}^+}$ is s_{n+1}-determinable from $\mathbf{M}_n$ •

The preceding lemma can be read in two ways:

Given $A = A(\sigma,\pi)[\mathbf{M}_n]_{n\in\omega} \in \mathcal{U}(\omega+\omega)$, the preceding proof describes a procedure to derive the short arrays $\mathbf{U} = \langle U_k\rangle_{k\in K}$ or $\mathbf{V} = \langle V_l\rangle_{l\in X} \subseteq \mathcal{U}(\omega)$ from the double array $\langle \mathbf{M}_n\rangle_{n\in\omega}$ (and $\langle\sigma,\pi\rangle \in \mathcal{G}(\omega+1)$).

Conversely, given $\mathbf{U} = \langle U_k\rangle_{k\in K}$ or $\mathbf{V} = \langle V_l\rangle_{k\in X} \subseteq \mathcal{U}(\omega)$, the proof gives a procedure to derive the double-array $\langle \mathbf{M}_n\rangle_{n\in\omega}$ from $\mathbf{U}$, respectively $\mathbf{V}$ (and $\langle\sigma,\pi\rangle$).

Note the following 'procedural' definition of the short arrays:

(i) Set $U_0 = V_0 = M_0^{0)}$.

(ii) Suppose $\langle U_i\rangle_{i<k+1}$ is defined, let $n = \eta_{k+1}$, and set

$$U_{k+1} = \begin{cases} M_0^{(n+1)} & \text{if } \tau_n = 1 \\ M_{\ell_n}^{(n+1)} & \text{if } \tau_n = 0. \end{cases}$$

(ii) Suppose $\langle V_l\rangle_{l\in w_n}$ is defined, let n' minimal $\geq n$ such that $\nu_{n'+1} \neq \emptyset$ and set for $l'+1 = \xi_{n+1} =$ the unique element in $w_{n'+1} \setminus w_n$

$$V_{l'+1} = \begin{cases} M_0^{(n'+1)} & \text{if } \tau_{n'} = 1 \\ M_{\ell_{n'}}^{(n'+1)} & \text{if } \tau_{n'} = 0. \end{cases}$$

Thus, a double array $\mathbf{M} = \langle M_i^{(n)}\rangle_{n\in\omega, i\in\ell_n}$ and the 'short' arrays $\mathbf{U} = \langle U_k\rangle_{k\in K}$ or $\mathbf{V} = \langle V_l\rangle_{l\in X}$ related as in the array equivalence lemma can be used interchangably in the characterisation of $A \in \mathcal{U}(\omega+\omega)$. Set

$$A(\sigma,\pi)\{U_k\}_{k\in K} = A(\sigma,\pi)\{V_l\}_{l\in X} = A(\sigma,\pi)[\mathbf{M}_n]_{n\in\omega} \tag{7.2}$$

whenever $\langle \mathbf{M}^{(n)}\rangle_{n\in\omega}$ reduces to (and is derivable from) $\langle U_k\rangle_{k\in K}$, or $\langle V_l\rangle_{l\in X}$, and call the first two expression a *short representation.*

(7.7) **Corollary** (Short rank $\omega+\omega$ representation lemma)

(a) Every $A \in \mathcal{U}(\omega+\omega)$ has a representation of the short forms (7.2).

(b) For every short array $\mathbf{U} \in {}^K\mathcal{U}\omega)$, respectively $\mathbf{V} \in {}^X\mathcal{U}\omega)$, the chain constructed as $A(\sigma,\pi)\{U_k\}_{k\in K})$, respectively $A(\sigma,\pi)\{V_l\}_{l\in X})$ is in $\mathcal{U}(\omega+\omega)$. •

[7.2] Generating local short arrays

It remains to exhibit a procedure to extract (via $\langle\sigma,\pi\rangle \in \mathcal{G}(\omega+1)$) the 'local' tagged short arrays $\mathbf{U}_n = \langle\langle U_k^{(n)}, k\rangle : k \in w_n\rangle$, from the 'global' short K-array $\mathbf{U}$, respectively X-array $\mathbf{V}$, for $A \in \mathcal{U}(\omega+\omega)$. (Here, $U^{(n)}_{\psi_n(i)} = M_i^{(n)}$ for $i \in \mathcal{L}_n$.) This will permit us to define $A = A(\sigma,\pi)\{\mathbf{U}\}$, respectively $A = A(\sigma,\pi)\{\mathbf{V}\}$, just if for some $x \in A$ and all n, $[x]^A_{\omega+n} = A(s_n)\{\mathbf{U}_n\}$, and vice-versa derive the short representation from the short 'local' approximations — but without having to take the detour through the $\mathbf{M}_n$ arrays.

We agreed to call a sequence $\mathbf{U} = \langle U_k : k \in I\rangle$, (where I is an initial segment of ω) a short array if for all $k, k' \in K$, $\mathbf{U}_k \in \mathcal{U}(\omega)$ and $\mathrm{f}U_k \cong \mathrm{f}U_{k'}$. Call the short array $\mathbf{U}$ *compatible* with $\langle\sigma,\pi\rangle \in \mathcal{G}(\omega+1)$ if $I = K$, and call it *realisable* if for some $A \in \mathcal{U}(\omega+\omega)$, and some compatible $\langle\sigma,\pi\rangle \in \mathcal{G}(\omega+1)$, $A = A(\sigma,\tau)\{\mathbf{U}\}$. (By the short $\omega+\omega$ representation theorem, every short array is realisable with every compatible $\langle\sigma,\pi\rangle \in \mathcal{G}(\omega+1)$.) We also said we could dispense with tagging the local short arrays if we can provide a 'canonical' choice of paths through $\mathcal{T}(s)$. We do not propose such a choice 'unifomly' for all $s \in \mathcal{S}(<\omega)$, but there is a natural such choice for each set $\{s_n\}_n$ derived from $\langle\sigma,\pi\rangle \in \mathcal{U}(\omega)$, namely the 'embedding' path $P_n = P_n(\sigma,\pi)$ determined by the $\tau_m, m < n$, defined as follows:

(7.8) **Definition**

(i) The top node is in P_n: $\langle s_n; 0\rangle \in P_n$.

(ii) Suppose $\sigma = \langle t, k\rangle \in P_n$ and let $m = |t| - 1$. If $m = 1$, the definition of P_n is complete. Else set

$$\langle t^+, k+1\rangle \in P_n \text{ if } \tau_{m-1} = 0,$$
$$\langle t^-, 0\rangle \in P_n \text{ if } \tau_{m-1} > 0.$$

Immediately from the definition, for all $m < n$,

$$\langle s_m, k_m\rangle \in P_n \text{ for some } k_m \leq m.$$

This is true for $m = n$ with $k_0 = 0$. Supposing $\langle s_m, k_m\rangle \in P_n$ and $m > 0$,

$$\langle s_m^+, k+1\rangle \in P_n \text{ and } s_{m-1} = s_m^+ \text{ if } \tau_{m-1} = 0,$$
$$\langle s_m^-, 0\rangle \in P_n \text{ and } s_{m-1} = s_m^- \text{ if } \tau_{m-1} > 0,$$

so $\langle s_{m-1}, k_{m-1}\rangle \in P_n$ for $k_{m-1} = k_m + 1$ if $\tau_{m-1} = 0$, else $k_{m-1} = 0$. Hence we are justified in calling the paths P_n 'determined by $\langle\sigma,\tau\rangle$.

Observe that the canonical ordering imposed on the local arrays by taking this path is an initial segment of the ordering imposed by 'novelty' on the short array for A. We can therefore define local array (untagged) short array as the initial segment of the (global) short array $\mathbf{U}$ 'used' at the given level:

(7.9) **Definition**

$$\mathbf{U}'_n = \langle U_k : k \leq \kappa_n\rangle = \langle U^{(\eta_k)}_{\nu_{\eta_k}} : k \leq \kappa_n\rangle$$

I.e., $\mathbf{U}'_n = \mathbf{U}\lceil\kappa_n$.

Since $\kappa_n = \zeta_n - 1$ (lemma 7.3), we can pass from the untagged to the tagged array by ordering $w_n = \{v_0 < \ldots, v_{\kappa_n}\}$ and setting $\langle U_{v_k}, v_k\rangle = \langle U^{(\eta_k)}_{\nu_{\eta_k}}, v_k\rangle$. I.e., $U^{(n)}_{\psi_n(i)} = U_k$ just if $\psi_n(i)$ is the kth largest element in w_n. Being able to dispense the tags, we shall now use '$\mathbf{U}_n$' to refer to the untagged local short arrays.

(7.10) **Definition** Define $\mathbf{U}^{\pm}_n$ (relative to σ, π and $\mathbf{U}$), so that $\mathbf{U}^{\pm}_n$ is the 'untagged' version of the original tagged $(\mathbf{U}_n)^{\pm}_{s_n}$ of section 3. (We may assume $n > 0$.)

If $\nu_n = \emptyset$, set $\mathbf{U}^-_n = \mathbf{U}^+_n = \mathbf{U}_n$.
If $\nu_n = \{0\}$ and $\tau_{n-1} = \ell^+_n$ set $\mathbf{U}^-_n = \mathbf{U}_n\lceil\kappa_{n-1}$ and $\mathbf{U}^+_n = \mathbf{U}_n$.
If $\nu_n = \{\ell^+_n\}$ and $\tau_{n-1} = 0$ set $\mathbf{U}^+_n = \mathbf{U}_n\lceil\kappa_{n-1}$ and $\mathbf{U}^-_n = \mathbf{U}_n$.

In set notation:
If $\nu_n = \{0\}$ and $\tau_{n-1} = \ell^+_n$, ran $\mathbf{U}^-_n =$ ran $\mathbf{U}_n \setminus \{U_{\nu_n}\}$ and ran $\mathbf{U}^+_n =$ ran $\mathbf{U}_n$.
If $\nu_n = \{\ell^+_n\}$ and $\tau_{n-1} = 0$, ran $\mathbf{U}^+_n =$ ran $\mathbf{U}_n \setminus \{U_{\nu_n}\}$ and ran $\mathbf{U}^-_n =$ ran $\mathbf{U}_n$.

(7.11) **Corollary**

(1) The sequence of string $\mathbf{U}_n$ over $\mathcal{U}(\omega)$ is *generated by (derived from)* $\langle\sigma, \pi\rangle$ and $\mathbf{U}$ in the follwing sense:

$$\mathbf{U}_0 = \{U_0\}$$

and for all n,

$$\mathbf{U}_{n+1} = \begin{cases} \mathbf{U}_n & \text{if } \nu_{n+1} = \emptyset \\ \mathbf{U}_n \wedge \langle U_{\nu_{n+1}}\rangle & \text{if } \nu_{n+1} \neq \emptyset. \end{cases}$$

(2) Let $\{\mathbf{U}_n\}_{n\in\omega}$ where $\mathbf{U}_n = \langle U_k\rangle_{k\leq\kappa_n}$, be the sequence generated by $\langle\sigma, \pi\rangle \in \mathcal{G}(\omega+1)$ and $\mathbf{U} = \langle U_k\rangle_{k\in K} \in {}^K\mathcal{U}(\omega)$, Then for each n, $\langle U_k : k \in w_{n+1}\rangle$ is an end-extension $\langle U_k : k \in w_n\rangle$, and the 'full' arrays $\mathbf{M}_n = \mathbf{M}_{\mathcal{L}_n} = \langle M^{(n)}_i\rangle_{i\in\mathcal{L}_n}$ are determined by $M^{(n)}_i = U_k$ just if $\psi_n(i)$ is the kth largest element in w_n.

(3) If $A = A(\sigma, \pi)\{\mathbf{U}\}$ and $\mathbf{U}_n$ are derived from $\langle\sigma, \pi\rangle \in \mathcal{G}(\omega+1)$ and $\mathbf{U} \in {}^K\mathcal{U}(\omega)$, then for some $x \in A$ and all n, $[x]^A_{\omega+n} = A(s_n)\{\mathbf{U}_n\} = A(s_n)[\mathbf{M}_n]$, and the approximating chain of sub-components can be generated by the following variant of Pastjin's uniform chain constructor:

$$A(0)\{U_0\} = U_0,$$
$$A(s_n)\{\mathbf{U}_n\} = \begin{cases} A(s^-_n)\{\mathbf{U}^-_n\}.\mathrm{Z} = A(s^-_n)\{\mathbf{U}_n\}.\mathrm{Z} & \text{if } s_n(n) = n \\ A(s^+_n)\{\mathbf{U}^+_n\} + A(s^-)\{\mathbf{U}^-_n\}.\mathrm{N} & \text{if } s_n(n) < n. \end{cases}$$

Note that the $\mathbf{U}_n$ 'argument' does not get truncated in the '$\nu^s = \emptyset$' cases arising both under $s(n) = n$ and $s(n) < n$. (Of course in the '$s(n) = n$' case ν_n is always $\emptyset$.) •

[8] Short representations of tails

Passing from the short array for a uniform chain of rank $\omega+\omega$ to that of its tail involves an additional twist, hinted at in the tail revelation lemma.

[8.1] Short representations of rank $\omega+\omega$

The preceding shows how to replace the arrays $\mathbf{M}_n$ in the description of $A \in \mathcal{U}(\omega+\omega)$. Now

$$\mathrm{f}A(\sigma,\pi)[\mathbf{M}_n]_{n\in\omega} = A(\mathrm{f}\sigma,\mathrm{f}\pi)[\mathrm{f}_{s_n}\mathbf{M}_n]_{n\in\omega}$$

and we would like similarly to replace the arrays $\mathrm{f}_{s_n}\mathbf{M}_n$ in the description of tails $\mathrm{f}A \in \mathcal{U}_0(\omega+\omega)$ (subsequently referred to as 'tail'-arrays). But note that the tail-arrays 'telescope' (since for tails, $\tau_n \equiv 0$); i.e., $\mathrm{f}_{s_{n+1}}\mathbf{M}_{n+1} = \mathrm{f}_{s^{\pm}_{n+1}}\mathbf{M}^{\pm}_{n+1} \ \wedge\ \mathbf{M}^{-}_{n+1}$ where the 'plus half', $\mathrm{f}_{s^{\pm}_{n+1}}\mathbf{M}^{\pm}_{n+1}$ belongs to the preceding level. Thus the building blocks 'new' in the tail array at level $n+1$ are determined by the 'minus half', $\mathbf{M}^{-}_{n+1}$. This is also clear by evaluating the expression for tail-chains (section 6):

$$\mathrm{f}A(\sigma,\pi)[\mathbf{M}_n]_{n\in\omega} \cong \mathrm{f}M_0^{(0)} + \sum_{n\in\omega} A(\sigma\lceil n+1)[\mathbf{M}^{-}_{n+1}].\mathrm{N}$$

Now compare how the arrays $\langle\mathbf{M}_{n+1}\rangle$ and $\langle\mathbf{M}^{-}_{n+1}\rangle$ relate to $\langle\mathbf{M}_n\rangle$ and $\mathbf{U}$ (via $\langle\sigma,\pi\rangle$). Here, the 'range' of an array is the set of possibly non-isomorphic building-blocks. Namely,

$$\mathbf{M}_0 = \langle M_0^{(0)}\rangle = \langle U_0\rangle$$

$$\operatorname{ran}\mathbf{M}_{n+1} = \begin{cases} \operatorname{ran}\mathbf{M}_n & \text{if } \nu_{n+1}=\emptyset \\ \{U_{\psi_{n+1}(0)}\}\cup\operatorname{ran}\mathbf{M}_n & \text{if } \nu_{n+1}\neq\emptyset \text{ and } \tau_n=1 \\ \operatorname{ran}\mathbf{M}_n\cup\{U_{\psi_{n+1}(\ell_n)}\} & \text{if } \nu_{n+1}\neq\emptyset \text{ and } \tau_n=0 \end{cases}$$

while

$$\mathrm{f}\mathbf{M}_0 = \langle \mathrm{f}M_0^{(0)}\rangle = \langle \mathrm{f}U_0\rangle$$

$$\operatorname{ran}\mathbf{M}^{-}_{n+1} = \begin{cases} \operatorname{ran}\mathbf{M}_n & \text{if } \tau_n>0 \text{ or } \nu_{n+1}=\emptyset \\ \{U_{\psi_{n+1}(\ell_n)}\}\cup\operatorname{ran}\mathbf{M}_n & \text{if } \tau_n=0 \text{ and } \nu_{n+1}\neq\emptyset \end{cases}$$

Thus, in the 'tail case' one of the following holds:

(i) $\nu_{n+1}=\emptyset$: In this case $\mathbf{M}^{-}_{n+1}$ is determined by s_{n+1} compatibility with $\mathbf{M}_n$. Or,

(ii) $\nu_{n+1}\neq\emptyset$, but $\tau_n=1$: In this case $\mathbf{M}^{-}_{n+1}=\mathbf{M}_n$; thus, in this case, the 'new' ω component is 'hidden', or 'invisible' to the tail. Or,

(iii) $\nu_{n+1}\neq\emptyset$ and $\tau_n=0$: Then $\mathbf{M}^{-}_{n+1}=\langle U_{\nu_{n+1}}\rangle\wedge\mathbf{M}^{(n)}_{(\ell^{+}_{n+1},\ell_{n+1})}$ where $U_{\nu_{n+1}}=U_{\kappa_{n+1}}$ and where the array $\mathbf{M}^{(n)}_{(\ell^{+}_{n+1},\ell_{n+1})}$ s_{n+1}-determinable from $\mathbf{M}_n$. In this case, therefore, the 'new' ω component is 'visible' to the tail.

Hence, a 'new' rank ω building block enters the $n+1$st 'tail' array only if $\tau_n=0$. If $\tau_n>0$, the 'new' element may have to be traced back several levels back, namely to the largest level $k<n$ such that $\tau_k=0$. E.g., if $n>0$ and $\tau_{n-1}=0$, $M^{(n)}_{\ell_{n-1}}=U_{\xi_n}$. If $n>0$ and $\tau_{n-1}>0$ also, then the last 'new' ω component was introduced at an even earlier level. More precisely, a ’new’ element U_k enters the ’tail’ array, essentially $\mathbf{M}^-_{n+1}$, just if either

(i) $\tau_n = 0$ and $n = \eta_k$ or

(ii) n is minimal such that for some $m < n$, $\tau_m = 1$ and $m = \eta_k$, $\tau_{n'} = 0$ for $n' \in (m, n)$, and $\tau_n = 1$

(and while $\tau_{n'} = 0$ in the second scenario, some ’later new’ elements may have entered the tail-array, namely $U_{k'}$ where $n' = \eta_{k'}$ for $n' \in (m, n)$).

(8.1) **Definition** ($\mathbf{W}_L = \mathrm{f}\mathbf{U}_K$)

We define the map that completes the bottom arrow in the following diagramme:

$$\begin{tikzcd}
A(\sigma,\pi)[\mathbf{M}_n]_n \arrow[r, "\mathrm{f}"] \arrow[d] & A(\mathrm{f}\sigma,\mathrm{f}\pi)[\mathrm{f}_{s_n}\mathbf{M}_n]_n \arrow[d] \\
\langle \mathbf{M}_n \rangle_{n\in\omega} \arrow[r] \arrow[d] & \langle \mathrm{f}_{s_n}\mathbf{M}_n \rangle_{n\in\omega} \arrow[d] \\
\langle U_k \rangle_{k\in K} \arrow[r] & \langle W_l \rangle_{l\in L}
\end{tikzcd}$$

I.e., set $\mathbf{W}_L = \mathrm{f}\mathbf{U}_K$ whenever

$\mathbf{U}_K = \langle U_k \rangle_{k\in K}$ is the short array for some $A(\sigma,\pi)[\mathbf{M}_n]_n$, and

$\mathbf{W}_L = \langle W_l \rangle_{l\in L}$ is the short array for $\mathrm{f}A = A(\mathrm{f}\sigma,\mathrm{f}\pi)[\mathrm{f}_{s_n}\mathbf{M}_n]_n$,

where $\mathrm{f}\sigma = \sigma\lceil\omega \wedge \langle 0\rangle$, where $\mathrm{f}\pi = \langle 0^\omega\rangle$, and where $\mathrm{f}_{s_n}(\mathbf{M}_n)$ is as defined in section 3.

Outline: Set $W_0 = \mathrm{f}U_0$.

While $\tau_n = 0$ ($n = 0, 1, 2, \ldots$), set $W_l = U_k$ whenever n is such that $\nu(n+1) \neq \emptyset$ (by implication, $\nu(n+1) = \{\ell_n\} = \{\ell^+_{n+1}\}$), and l is minimal such that W_l is not yet defined.

For the first $n_1 \geq 0 = n_0 + 1$ (i.e., we set $n_0 = -1$) such that $\tau_{n_1} > 0$, set $W_l = U_0$ (l as before).

While $\tau_n = 0$ ($n = n_1 + 1, n_1 + 2, \ldots$), set $W_l = U_k$ whenever n is such that $\nu_n \neq \emptyset$ (and l as before).

For the second $n_2 > n_1$ such that $\tau_{n_1} > 0$, set $W_l = U_{k'}$ (l as before), provided it was the case that $n_1 + 1 = \eta_{k'}$ (i.e., $\nu(n_1 + 1) \neq \emptyset$ - else W_l remains undefined). Etc.

Detail: Let $\{n_i : i \in I\}$ be the set of n such that $n = 0$ or $n > 0$ and $\tau_{n-1} > 0$. (Here I is an initial segment of ω, i.e., $I = \omega$ or $I \in \omega$.) Set $n_I = \omega$ if I is finite (i.e., $I \in \omega$). Thus, $\omega = \bigcup_{i\in I}[n_i, n_{i+1})$.

Set $U_k^{(n)} :=\uparrow$ (to mean: ’is undefined’), if $n < \eta_k$, and set $U_k^{(n)} := U_k$, if $n \geq \eta_k$ ($k \in K$). Say $U_k^{(n)} :=\downarrow$ (to mean ’is defined’), if $U_k^{(n)} = U_k \neq\uparrow$. Let $\overline{k} = \max\{k : \eta_k < n\} = \max\{k : U_k^{(n-1)} \downarrow\}$. (Note $\overline{k}(0) = \max\emptyset = 0$.)

Define $W_0^{(0)} := W_0 := \mathrm{f}U_0$ and $W_l^{(0)} :=\uparrow$ for all $l > 0$. Let $\overline{l} = \max\{l : W_l^{(n-1)} \downarrow\}$. (So, $\overline{l}(0) = 0$.)

Let now $n > 0$.

Suppose $n \in (n_i, n_{i+1})$ for some i. If $n \neq \eta_k$, for any k, set $\overline{l}(n) := \overline{l}(n-1)$ and $W_l^{(n)} := W_l^{(n-1)}$ for all l. Else, for the unique k such that $n = \eta_k$, set $\overline{l}(n) := \overline{l}(n-1) + 1$, $W_{\overline{l}(n)}^{(n)} := U_k$ and $W_l^{(n)} := W_l^{(n-1)}$ for all $l \neq \overline{l}(n)$.

Suppose $n = n_i$ for some i (since $n > 0$, $i > 0$). If $n_{i-1} \neq \eta_k$, for any k, set $\overline{l}(n) := \overline{l}(n-1)$ and $W_l^{(n)} := W_l^{(n-1)}$. Else, for the unique k such that $n_{i-1} = \eta_k$, set $\overline{l}(n) := \overline{l}(n-1) + 1$, $W_{\overline{l}(n)}^{(n)} := U_k$ and $W_l^{(n)} := W_l^{(n-1)}$ for all $l \neq \overline{l}(n)$.

End of definition.

(8.2) **Corollary** For all n, $\overline{l}(n) \in \{\overline{k}(n), \overline{k}(n)+1\}$ and for all $i \in I$, if $n_i \neq \eta_k$, for any k, then for all $n \in [n_i, n_{i+1})$, $\overline{l}(n) = \overline{k}(n)+1$; if $n_i = \eta_k$ for some unique k, then for all $n \in [n_i, n_{i+1})$, $\overline{l}(n) = \overline{k}(n)$.

Proof By induction.

For $n = 0$, $\overline{l}(0) = 0 = \overline{k}(0)$.

Let $n > 0$, and suppose the first claim is true for all $n' < n$, and if $n = n_{i+1}$, $i \in I$, the second claim is true for i.

If $n \in (n_i, n_{i+1})$, then from the definiton, $\overline{l}(n) - \overline{l}(n-1) = \overline{k}(n) - \overline{k}(n-1)$, so the claim continues to hold, while the second claim holds by continuing the induction over $n'' \in (n_i, n_{i+1})$.

Suppose then $n = n_i$ for some i, and we may assume $i > 0$. If either both n_i and n_{i-1} are in $\{\eta_k\}_k$ or neither is, then again $\overline{l}(n) - \overline{l}(n-1) = \overline{k}(n) - \overline{k}(n-1)$, for all $n \in (n_i, n_{i+1})$ by hypothesis, and for $n = n_i$ by definition, so the induction proceeds as in the previous paragraph. Else one of the following holds:

Either $n_{i-1} \neq \eta_k$, for any k, while $n_i = \eta_{k'}$ for some (unique) k'; then $\overline{l}(n_i) = \overline{l}(n_i - 1) = \overline{k}(n_i - 1) + 1 = \overline{k}(n_i)$.

Or $n_{i-1} = \eta_k$, for some (unique) k, while $n_i \neq \eta_{k'}$ for any k'; then $\overline{l}(n_i) = \overline{l}(n_i - 1) + 1 = \overline{k}(n_i - 1) + 1 = \overline{k}(n_i) + 1$. •

(8.3) **Corollary**

$$\mathrm{f}A(\sigma,\pi)\{U_i\}_{i\in K} = A(\sigma\lceil\omega\wedge\langle 0\rangle)\{W_i\}_{i\in L} = \mathrm{f}M_0^{(0)} + \sum_{n\in\omega} A(\sigma\lceil n+1)[\mathbf{M}^-_{n+1}].\mathrm{N}.$$

The next lemma tells where (and if) a given building block U_k appears in the tail.

(8.4) **Lemma** (Tail relocation lemma) For $0 < k < K$, let $n_k = \eta_k - 1$ (being thus also the least n such that $\kappa_{n+1} = k$).

(i) Either for all $0 < k < K$,

(a) $\tau_{n_k} = 0$, in which case $U_k = M_{\ell_{n_k}}^{(n_k+1)}$ gets inserted into $\mathbf{W}_L$ at 'stage n_k' (i.e., as W_k or W_{k+1}); or

(b) $\tau_{n_k} = 1$, but there is some $m \geq n_k + 1$ such that again $\tau_m = 1$, in which case $U_k = M^{n_k+1})_0 = M_0^{(m)} = M_{\ell_m}^{(m+1)}$ gets inserted into $\mathbf{W}_L$ at stage m, for the least such m.

(ii) Or there is a unique $0 < k < K$ such that $\tau_{n_k} = 1$, in which case $U_k = M_0^{(n_k+1)}$, but for all $m \geq n_k + 1$, $\tau_m \in \{0, \omega\}$, and the copy-definition $U_k = M_0^{(m)} = M_{\ell_m}^{(m+1)}$ is never made, whence U_k does not get inserted into $\mathbf{W}_L$ (although this may happen to some other elements $U_{k'}$ isomorphic to U_k.)

Proof Suppose (i) fails and let k_0 be the least k such that (i) fails for k, i.e., $\tau_{n_{k_0}} = 1$, but there is no $m \geq n_{k_0}$ such that $\tau_m = 1$, i.e., for all $m \geq n_{k_0}$, $\tau_m \in \{0, \omega\}$. For any $k \in (k_0, K)$ we have $\tau_{n_k} \neq \omega$, hence by the assumption that (i) fails at k_0, $\tau_{n_k} = 0$; so these k all satisfy (i) via the first disjunct. Thus, k_0 uniquely satisfies (ii). •

Remark (Scenarios for $\langle W_l : l \in L\rangle$)

(a) Suppose first that for all m, $\tau_m = 0$. In this case U_0 always gets 'copied' as $M_0^{(m+1)}$ into the 'plus' half of the level m building blocks, hence never appears (as $M_{\ell_m}^{(m+1)}$) in a 'tail'-array $\mathbf{M}^-_{m+1}$, while every U_k

for $k > 0$ gets ’inserted’ (as $M^{(n_k+1)}_{\ell_{n_k}}$) in the ’tail’-array $\mathbf{M}^-_{n_k+1}$. at the same level at which it appears in the full array $\mathbf{M}_{n_k+1}$. In particular, $W_0 = \mathrm{f}U_0$, and $W_{k+1} = U_{k+1}$ for $k \in K$, and also $L = K$. U_0 never appears in the tail array.

Note that if $\tau_m = \omega$, then $\mathrm{f}_{s_{m+1}}\mathbf{M}_{m+1} = \mathrm{f}_{s^-_{m+1}}\mathbf{M}_{m+1} \wedge \mathbf{M}_{m+1}$, so U_0 does get copied into the tail-array at level $m+1$. If $\tau_m = 1$ for some least m, then $U_0 = M^{(m)}_0$ gets copied as $M^{(m+1)}_{\ell^+_{m+1}}$ into the ’minus half’ of the level $m+1$ building blocks, hence also into the tail-array $\mathrm{f}_{s_{m+1}}\mathbf{M}_{m+1} = \mathrm{f}_{s^+_{m+1}}\mathbf{M}^+_{m+1} \wedge \mathbf{M}^-_{m+1}$ of level $m+1$.

(b) So suppose that $\tau_m = 1$ for some $m \in \omega$. Let m_0 be the least m such that $\tau_m = 1$; then for some $k = k_1 \in K$, m_0 also is the least n such that $\kappa_n = k_1$. Then for all $n \leq m_0$ such that $\tau_n < \omega$ we have in fact $\pi_n = 0$, and so $U_k = M^{(n_k+1)}_{\ell^+_{n_k+1}}$ for $k \in (0, k_1)$, while level m_0+1 represents the earliest opportunity to copy U_0 into the ’minus half’, and hence the tail-array, $U_0 = M^{(m_0+1)}_{\ell^+_{m_0+1}}$. (The extreme case $m_0 = k_1 = 0$ is possible, in which case the prescription for $k \in (0, k_1)$ will be vacuous.)

(b.1) Suppose next and that the first case of the tail relocation lemma applies, i.e., that for all $k \in L$, (i) holds. Let $k_{2i}, i = 1, 2\ldots$ such that $\tau_{n_{k_{2i}}} = 1$ and $m_i, i = 1, 2, \ldots$ such that m_i is the least $m \geq n_{k_{2i}} + 1$ such that $\tau_{m_i} = 1$. Let $k_{2i+1}, i = 1, 2\ldots$ such that $\kappa_{m_i} = k_{2i+1}$. Then $U_k = M^{(n_k+1)}_{\ell^-_{n_k+1)}}$ for $k \in (k_{2i}, k_{2i+1})$, while $U_{k_{2i}} = M^{(m_i+1)}_{\ell^-_{m_i+1}}$. Thus, $W_k = U_{k+1}$ for $k \in \bigcup_i [k_{2i}, k_{2i+1})$, where $k_0 := 0$, and $W_{k_{2i+1}} = U_{k_{2i}}$; in particular $W_{k_1} = U_{k_0} = U_0$.

If $K = n+1 \in \omega$, then n is maximal such that $\nu_{\eta_n} \neq \emptyset$. If $n = m_{i^0}$ for some last i^0, the definition is again complete, with $\{W_k : k \in K\} = \{U_k : k \in K\}$. Else $m_{i^0} < n$ and $\tau_{n_k} = 0$ for all $k \in (k_{2i^0+1}, K)$ so $U_k = M^{(n_k+1)}_{\ell^-_{n_k+1)}}$ for all such k, and so also $W_k = U_k$ for these k.

If $K = \omega$, k_i is defined for each $i \in \omega$, W_k is defined for each $k \in \omega$, $\{W_k : k \in \omega\} = \{U_k : k \in \omega\}$, and $L = K = \omega$.

(b.2) Finally, still assuming that $\tau_m = 1$ for some least m, suppose that the second case of tail relocation lemma applies, i.e., for some unique $k = k^0 \in K$ (ii) holds.

If $k^0 = 0$ then $U_0 = M^{(1)}_1$, while for all m, $U_1 = M^{(m+1)}_0$, and $U_{m+2} = M^{(m+1)}_{\ell^-_{m+1}}$, so $W_0 = U_0$ and for all $k \geq 1$, $W_k = U_{k+1}$, while U_1 never appears in the tail array. In particular, $L = K = \omega$.

So suppose $k_0 > 0$. For all $k < k_0$ case (i) of tail relocation lemma applies, so the definition starts as under (b.1), i.e., there exists a sequence $k_i, i = 0, \ldots, i^0$ in (k, k^0) such that $W_k = U_{k+1}$ for $k \in \bigcup_i [k_{2i}, k_{2i+1})$, where $k_0 := 0$ and $W_{k_{2i+1}} = U_{k_{2i}}$, and if $k_{2i^0+1} < k^0$, then $W_k = U_k$ for $k \in (k_{2i^0+1}, k^0)$. The remainder of the definition continues as in the case $k^0 = 0$, i.e., $U_{k^0+1} = M^{(k^0+1)}_0 = M^{(m+1)}_0$ for all $m \geq k^0 + 1$, while for all $k > k^0$ $U_k = M^{(n_k+1)}_{\ell^-_{n_k+1}}$. Thus, for all $k > k^0$ $W_k = U_{k+1}$, while U_{k^0} never gets inserted into $\mathbf{W}_L$, and also $L = K$.

[8.2] Tracking new building blocks in the tail array

We close this section by formalising a few observations about how new building blocks appear in tail arrays.

(8.5) **Definition**

$$\delta_0(\kappa_0) := \delta_0(0) = \nu_0$$

$$\delta_{n+1}(\kappa_{n+1}) := \begin{cases} \nu_{n+1} & \text{if } \nu_n \neq \emptyset \text{ and hence } \kappa_{n+1} = \kappa_n + 1. \\ \delta_n(\kappa_n) & \text{if } \nu_n = \emptyset \text{ and hence } \kappa_{n+1} = \kappa_n \end{cases}$$

and for $k < \kappa_{n+1}$

$$\delta_{n+1}(k) := \begin{cases} \delta_n(k) & \text{if } \tau_n = \omega \\ \delta_n(k) \cup \{i \in \mathcal{L}^-_{n+1} : (\forall j \in \delta(k))[\psi_n(i) = \psi_n(j)]\} & \text{if } \tau_n = 0 \\ \{i \in \mathcal{L}^+_{n+1} : (\forall j \in \delta(k))[\psi_n(i) = \psi_n(j)]\} \cup \{\ell(s_n) + i : i \in \delta_n(k)\} & \text{if } \tau_n = 1. \end{cases}$$

Remark $M^{(n)}_i \cong M^{(n')}_{i'}$ whenever for some $k \leq \kappa_{\max\{n,n'\}}$ $i, i' \in \delta_{\max\{n,n'\}}(k)$ and k is minimal for the property that $(\exists n'')[i, i' \in \delta_{n''}(k)]$.

(8.6) **Definition**

$$\chi(k) := \inf\{n \in \omega : \delta_n(k) \cap \mathcal{L}^-_{n+1} \neq \emptyset\}.$$

That is, $\chi(k)$ is the least level n at which the element $U_k \in \mathbf{U}$ gets 'copied' as $M^{(n)}_i$ in a position $i \in \mathcal{L}^-_{n+1}$ of the 'tail-array' $\mathbf{M}^-_{n+1}$, if there is, else $\chi(k) = \omega$.

(8.7) **Lemma** $\chi(k) < \omega$ iff for some n such that $\kappa_n \geq k$, $\tau_n \in (0, \omega)$.

Proof Fix U_k and let $n_k = \eta_k$ be the least n such that $\kappa_{n+1} = k$. Then either $k = n = 0$, or $\tau_{n_k} \in \{0, 1\}$ and $\ell_{n_k+1} = \ell^+_{n_k+1} + \ell^-_{n_k+1}$. Let i_k such that $U_k = M^{(n_k)}_{i_k}$. If $\tau_{n_k} = 0$, then $i_k = \ell_{n_k} = \ell^+_{n_k+1}$, and $\chi(k) = n_k + 1$. So suppose $\tau_{n_k} = 1$ and $i_k = 0$. If for all $m \geq n_k + 1$, $\tau_m = 0$, or $\tau_m = \omega$, then for no m with $\kappa_m \geq k$ is $\pi_m > 0$. and $\chi(k) = \omega$, since for all such m, $\mathbf{M}^-_{m+1} = \mathbf{M}_m$, i.e., for all $m \geq n_k + 1$, $U_k = M^m_0 = M^{m+1}_0$. Otherwise, there is a least $m \geq n_k + 1$ such that $\pi_m \in (0, \omega)$, and for this m, $\kappa_m \geq k$ and $\mathbf{M}^-_{m+1} = \mathbf{M}_m$ so in particular $U_k = M^m_0 = M^{m+1}_{\ell^+_{m+1}}$. •

[9] Extensions

We conclude the paper by pushing the construction principle exhibitd in section 7 to 'higher ranks'.

[9.1] Rank $\omega \cdot \omega = \omega^2$

Climbing through levels $\omega{\cdot}n$ is now fairly straight-forward: Write γ for $\langle \sigma, \pi \rangle \in \mathcal{G}(\omega)$. By 5.12, every $A \in \mathcal{U}(\omega)$ has a representation of form $A(\gamma)$, for some $\gamma \in \mathcal{G}(\omega)$; by 7.7 combined with 5.12, every $A \in \mathcal{U}(\omega + \omega)$ has a representation of form $A(\gamma)\{A(\gamma_k)\}_{k \in K}$ for $K = K(\gamma) \subseteq \omega$ and some $\gamma, \gamma_k \in \mathcal{G}(\omega)$ (only depending on A, and therefore satsfying $\mathrm{f}\sigma_k = \mathrm{f}\sigma_{k'}$, but otherwise arbitrary). By the same token, every $A \in \mathcal{U}(\omega + \omega + \omega)$ is of form

$$A(\gamma)\{A(\gamma_k)\{A(\gamma_{kl})\}_{l \in K(\gamma_k)}\}_{k \in K(\gamma)}, \tag{a}$$

for $\gamma, \gamma_k, \gamma_{kl}$ (again, only depending on A, and therefore satsfying $\mathrm{f}\sigma_k = \mathrm{f}\sigma_{k'}$ and $\mathrm{f}\sigma_{kl} = \mathrm{f}\sigma_{kl'}$ but otherwise arbitrary). Let's abbreviate (a) as

$$A(\gamma; \gamma_k; \gamma_{kl}) = A(\gamma; \gamma_k; \gamma_{kl})_{k \in K(\gamma), l \in K(\gamma_k)}$$

Formally, by induction,

(9.1) **Definition**

$$\begin{aligned} A(\gamma; \gamma_{k_1}; \gamma_{k_1 k_2}; \ldots; \gamma_{k_1 \ldots k_n}) &:= A(\gamma; \gamma_{k_1}; \gamma_{k_1 k_2}; \ldots; \gamma_{k_1 \ldots k_n})_{k_1 \in K(\gamma), k_2 \in K(\gamma_{k_1}), \ldots, k_n \in K(\gamma_{k_1 \ldots k_{n-1}})} \\ &:= A(\gamma)\{A(\gamma_{k_1}; \gamma_{k_1 k_2}; \ldots; \gamma_{k_1 \ldots k_n})_{k_2 \in K(\gamma_{k_1}), \ldots, k_n \in K(\gamma_{k_1 l \ldots k_{n-1}})}\}_{k_1 \in K(\gamma)} \end{aligned}$$

Generalizing the preceding observations:

(9.2) **Corollary** Every $A \in \mathcal{U}(\omega \cdot n)$ is of form

$$A(\gamma; \gamma_{k_1}; \gamma_{k_1 k_2}; \ldots; \gamma_{k_1 \ldots k_n}) = A(\gamma; \gamma_{k_1}; \gamma_{k_1 k_2}; \ldots; \gamma_{k_1 \ldots k_n})_{k_1 \in K(\gamma), k_2 \in K(\gamma_{k_1}), \ldots, k_n \in K(\gamma_{k_1 \ldots k_{n-1}})}$$

for some $\gamma, \gamma_{k_1}, \ldots, \gamma_{k_1 \ldots k_n}$ (only depending on A, and therefore satisfying $\mathrm{f}\sigma_{k_1} = \mathrm{f}\sigma_{k_1'}$ $\mathrm{f}\sigma_{k_1 k_2} = \mathrm{f}\sigma_{k_1 k_2'}$ $\ldots$ $\mathrm{f}\sigma_{k_1 \ldots k_{n-1} k_n} = \mathrm{f}\sigma_{k_1 \ldots k_{n-1} k_n'}$ ..., but otherwise arbitrary). ∘

Now let $x \in A \in \mathcal{U}(\omega^2)$; For each n,

$$[x]^A_{\omega \cdot n} = A(\gamma^{(n)}; \gamma^{(n)}_{k_1}; \gamma^{(n)}_{k_1 k_2}; \ldots; \gamma^{(n)}_{k_1 \ldots k_n})_{k_1 \in K(\gamma^{(n)}), k_2 \in K(\gamma^{(n)}_{k_1}), \ldots, k_n \in K(\gamma^{(n)}_{k_1 \ldots k_{n-1}})}$$

for some $\gamma^{(n)}, \gamma^{(n)}_{k_1}, \gamma_{k_1 k_2}, \ldots, \gamma^{(n)}_{k_1 \ldots k_n} \in \mathcal{G}(\omega)$, such that for each n there is some $\theta(n) := h \in K(\gamma^{n+1})$, such that

$$\begin{aligned} &A(\gamma^{(n)}; \gamma^{(n)}_{k_1}; \gamma^{(n)}_{k_1 k_2}; \ldots; \gamma^{(n)}_{k_1 \ldots k_n})_{k_1 \in K(\gamma^{(n)}), k_2 \in K(\gamma^{(n)}_{k_1}), \ldots, k_n \in K(\gamma^{(n)}_{k_1 \ldots k_{n-1}})} \\ &\quad = A(\gamma^{(n+1)}_h; \gamma^{(n+1)}_{h k_2}; \ldots; \gamma^{(n+1)}_{h, k_2 \ldots k_n})_{k_2 \in K(\gamma^{(n+1)}_h), \ldots, k_n \in K(\gamma^{(n+1)}_{h, k_2 \ldots k_{n-1}})} \end{aligned} \tag{9.1}$$

and hence such that

$$\begin{aligned} \gamma^{(n)} &= \gamma^{(n+1)}_h \\ \gamma^{(n)}_{k_1} &= \gamma^{(n+1)}_{h k_1} \\ &\ldots \\ \gamma^{(n)}_{k_1 \ldots k_n} &= \gamma^{(n+1)}_{h k_1 \ldots k_n} \end{aligned} \tag{9.2}$$

Define an 'embedding thread' between the ω-limit levels:

(9.3) **Definition** Let θ be the function defined via $\theta(n) = h$ just if (9.1), and hence (9.2), holds. Let

$$A(\gamma^{(0)};\gamma^{(1)};\gamma^{(1)}_{k_1^{(1)}};\gamma^{(2)};\gamma^{(2)}_{k_1^{(2)}};\gamma^{(2)}_{k_1^{(2)}k_2^{(2)}}\gamma^{(3)};\gamma^{(3)}_{k_1^{(3)}};\gamma^{(3)}_{k_1^{(3)}k_2^{(3)}};\gamma^{(3)}_{k_1^{(3)}k_2^{(3)}k_3^{(3)}};\ldots) \tag{9.3}$$

be the unique $A \in \mathcal{U}(\omega^2)$ such that

$$\begin{aligned}
&\exists h_0\ \gamma^{(0)} = \gamma^{(1)}_{h_0};\\
&\exists h_1\ \gamma^{(1)} = \gamma^{(2)}_{h_1}\\
&\quad \&\ \langle \gamma^{(1)}_{k_1^{(1)}} : k_1^{(1)} \in K(\gamma^{(1)})\rangle = \langle \gamma^{(2)}_{h_1 k_2^{(2)}} : k_2^{(2)} \in K(\gamma^{(2)}_{h_1})\rangle;\\
&\exists h_2\ \gamma^{(2)} = \gamma^{(3)}_{h_2}\\
&\quad \&\ \langle \gamma^{(2)}_{k_1^{(2)}} : k_1^{(2)} \in K(\gamma^{(2)})\rangle = \langle \gamma^{(3)}_{h_2 k_2^{(3)}} : k_2^{(3)} \in K(\gamma^{(3)}_{h_2})\rangle\\
&\quad \&\ \langle \gamma^{(2)}_{k_1^{(2)}k_2^{(2)}} : k_2^{(2)} \in K(\gamma^{(2)}_{k_1^{(2)}})\rangle = \langle \gamma^{(3)}_{h_2 k_2^{(3)} k_3^{(3)}} : k_3^{(3)} \in K(\gamma^{(3)}_{h_2 k_2^{(3)}})\rangle;\\
&\qquad \ldots
\end{aligned} \tag{9.4}$$

(9.4) **Corollary** (Rank ω^2 representation lemma) Every $A \in \mathcal{U}(\omega^2)$ has a representation of the form (9.3). •

For completeness, note that the nth block of γ tokens in representation (9.3) of A is

$$\ldots\gamma^{(n)}_{k_1^{(n)}};\gamma^{(n)}_{k_1^{(n)}k_2^{(n)}};\gamma^{(n)}_{k_1^{(n)}k_2^{(n)}k_3^{(n)}};\ldots;\gamma^{(n)}_{k_1^{(n)}\ldots k_n^{(n)}};\ldots \tag{9.3a}$$

and the 'embedding conditions' on the nth and $n+1$st block are:

$$\begin{aligned}
&\qquad \ldots\\
&\exists h_n\ \gamma^{(n)} = \gamma^{(n+1)}_{h_n}\\
&\quad \&\ \langle \gamma^{(n)}_{k_1^{(n)}} : k_1^{(n)} \in K(\gamma^{(n)})\rangle = \langle \gamma^{(n+1)}_{h_n k_2^{(n+1)}} : k_2^{(n+1)} \in K(\gamma^{(n+1)}_{h_n})\rangle\\
&\quad \&\ \ldots\\
&\quad \&\ \langle \gamma^{(n)}_{k_1^{(n)}\ldots k_{i+1}^{(n)}} : k_{i+1}^{(n)} \in K(\gamma^{(n)}_{k_1^{(n)}\ldots k_i^{(n)}})\rangle = \langle \gamma^{(n+1)}_{h_n k_2^{(n+1)}\ldots k_{i+1}^{(n+1)}} : k_{i+1}^{(n+1)} \in K(\gamma^{(n+1)}_{h_n k_2^{(n)}\ldots k_i^{(n+1)}})\rangle;\\
&\quad \&\ \ldots\\
&\quad \&\ \langle \gamma^{(n)}_{k_1^{(n)}\ldots k_n^{(n)}} : k_n^{(n)} \in K(\gamma^{(n)}_{k_1^{(n)}\ldots k_{n-1}^{(n)}})\rangle = \langle \gamma^{(n+1)}_{h_n k_2^{(n+1)}\ldots k_n^{(n+1)}} : k_n^{(n+1)} \in K(\gamma^{(n+1)}_{h_n k_2^{(n)}\ldots k_{n-1}^{(n+1)}})\rangle.\\
&\qquad \ldots
\end{aligned} \tag{9.4a}$$

The 'implicit' conditions on the $\gamma^{(n)}_{i_1\ldots i_n} = \langle \sigma^{(n)}_{i_1\ldots i_n}, \pi^{(n)}_{i_1\ldots i_n}\rangle$ are:

$$\mathrm{f}\sigma^{(n)}_{k_1\ldots k_{n-1}k_n} = \mathrm{f}\sigma^{(n)}_{k_1\ldots k_{n-1}k'_n}.$$

Let's call a system of $\gamma^{(n)}_{k_1^{(n)}\ldots k_n^{(n)}} \subset \mathcal{G}(\omega)$ satisfying the 'embedding conditions' (9.4) (as well as the 'implicit' conditions) a $\mathcal{G}(\omega^2)$ *system* and denote such by Γ. Corollary 9.4 says that every $A \in \mathcal{U}(\omega^2)$ is represented by a $\mathcal{G}(\omega^2)$ system. The following notation denotes a $\mathcal{G}(\omega^2)$ system, the arrows expressing the 'embedding conditions' (9.4):

$$\begin{tikzcd}[row sep=small, column sep=small]
\omega \; : & \gamma^{(0)} \arrow[dr] & & & & \\
\omega\cdot 2 \; : & \gamma^{(1)} \arrow[dr] & \gamma^{(1)}_{k_1^{(1)}} \arrow[dr] & & & \\
\omega\cdot 3 \; : & \gamma^{(2)} \arrow[dr] & \gamma^{(2)}_{k_1^{(2)}} \arrow[dr] & \gamma^{(2)}_{k_1^{(2)}k_2^{(2)}} \arrow[dr] & & \\
 & \cdots \arrow[dr] & \cdots & \cdots & \cdots \arrow[dr] & \\
\omega\cdot n \; : & \gamma^{(n)} \arrow[dr] & \gamma^{(n)}_{k_1^{(n)}} \arrow[dr] & \cdots & \gamma^{(n)}_{k_1^{(n)}\ldots k_n^{(n)}} \arrow[dr] & \\
 & \cdots & \cdots & \cdots & \cdots & \cdots
\end{tikzcd} \tag{9.5}$$

[9.2] Rank ω^3

Beyond ω^2 the pace begins to look slow, since we still only progress by ω steps, using the construction principle of 7.7. Schematically:

$$\begin{tikzcd}[row sep=small, column sep=small]
\omega^2 \; : & \Gamma^{(0)} \arrow[dr] & & & & \\
\omega^2+\omega \; : & \gamma^{(1)} \arrow[dr] & \Gamma^{(1)}_{k_1^{(1)}} \arrow[dr] & & & \\
\omega^2+\omega\cdot 2 \; : & \gamma^{(2)} \arrow[dr] & \gamma^{(2)}_{k_1^{(2)}} \arrow[dr] & \Gamma^{(2)}_{k_1^{(2)}k_2^{(2)}} \arrow[dr] & & \\
 & \cdots \arrow[dr] & \cdots & \cdots & \cdots \arrow[dr] & \\
\omega^2+\omega\cdot n \; : & \gamma^{(n)} \arrow[dr] & \gamma^{(n)}_{k_1^{(n)}} \arrow[dr] & \cdots & \Gamma^{(n)}_{k_1^{(n)}\ldots k_n^{(n)}} \arrow[dr] & \\
 & \cdots & \cdots & \cdots & \cdots & \cdots
\end{tikzcd} \tag{9.6}$$

where the embedding conditions for the $\gamma^{(n)}_{i_1\ldots i_n}$ translate verbatim into embedding conditions for the $\Gamma^{(n)}_{i_1\ldots i_n}$. (The 'implicit' conditions on the $\gamma^{(n)}_{i_1\ldots i_{n-1}}$ are as before; the 'implicit' conditions on the $\Gamma^{(n)}_{i_1\ldots i_n}$ are 'implicit' in the 'implicit' conditions of their constituent $\gamma^{(n)}_{i_1\ldots i_n}$.) Again, given $A\in\mathcal{U}(\omega^2+\omega^2)$ and $x\in A$, there is a system as in (9.6) such that for each n,

$$[x]^A_{\omega^2+\omega\cdot n} = A(\gamma^{(n)};\gamma^{(n)}_{k_1};\gamma^{(n)}_{k_1k_2};\ldots;\gamma^{(n)}_{k_1\ldots k_{n-1}};\Gamma^{(n)}_{k_1\ldots k_n})_{k_1\in K(\gamma^{(n)}),k_2\in K(\gamma^{(n)}_{k_1}),\ldots,k_n\in K(\gamma^{(n)}_{k_1\ldots k_{n-1}})},$$

so every $A\in\mathcal{U}(\omega^2+\omega^2)$ has a representation by a $\mathcal{G}(\omega^2\cdot 2)$ system of from (9.6). Now iterate. Replace the $\mathcal{G}(\omega^2)$ systems $\Gamma^{(n)}_{i_1\ldots i_n}$ in (9.6) by $\mathcal{G}(\omega^2\cdot 2)$ systems (again denoted $\Gamma^{(n)}_{i_1\ldots i_n}$) and obtain a $\mathcal{G}(\omega^2\cdot 3)$ system representing $A\in\mathcal{U}(\omega^2\cdot 3)$.

Formally, if (for some ordinal α) the concept of a $\mathcal{G}(\alpha)$ system is defined, a $\mathcal{G}(\alpha+\omega^2)$ system is a system for form

$$\begin{tikzcd}
\alpha & : & \Gamma^{(0)} \arrow[dr] & & & & \\
\alpha+\omega & : & \gamma^{(1)} \arrow[dr] & \Gamma^{(1)}_{k_1^{(1)}} \arrow[dr] & & & \\
\alpha+\omega\cdot 2 & : & \gamma^{(2)} \arrow[dr] & \gamma^{(2)}_{k_1^{(2)}} \arrow[dr] & \Gamma^{(2)}_{k_1^{(2)}k_2^{(2)}} \arrow[dr] & & \\
 & & \cdots \arrow[dr] & \cdots & \cdots \arrow[dr] & \cdots & \\
\alpha+\omega\cdot n & : & \gamma^{(n)} \arrow[dr] & \gamma^{(n)}_{k_1^{(n)}} \arrow[dr] & \cdots & \Gamma^{(n)}_{k_1^{(n)}\dots k_n^{(n)}} \arrow[dr] & \\
 & & \cdots & \cdots & \cdots & \cdots & \cdots
\end{tikzcd}$$

where each $\Gamma^{(n)}_{k_1^{(n)}\dots k_n^{(n)}}$ is a $\mathcal{G}(\alpha)$ system, and the arrows represent the embedding conditions (9.4):

$$\begin{aligned}
&\exists h_0\ \Gamma^{(0)} = \Gamma^{(1)}_{h_0};\\
&\exists h_1\ \gamma^{(1)} = \gamma^{(2)}_{h_1}\\
&\ \ \&\ \langle \Gamma^{(1)}_{k_1^{(1)}} : k_1^{(1)} \in K(\gamma^{(1)})\rangle = \langle \Gamma^{(2)}_{h_1 k_2^{(2)}} : k_2^{(2)} \in K(\gamma^{(2)}_{h_1})\rangle;\\
&\exists h_2\ \gamma^{(2)} = \gamma^{(3)}_{h_2}\\
&\ \ \&\ \langle \gamma^{(2)}_{k_1^{(2)}} : k_1^{(2)} \in K(\gamma^{(2)})\rangle = \langle \gamma^{(3)}_{h_2 k_2^{(3)}} : k_2^{(3)} \in K(\gamma^{(3)}_{h_2})\rangle\\
&\ \ \&\ \langle \Gamma^{(2)}_{k_1^{(2)} k_2^{(2)}} : k_2^{(2)} \in K(\gamma^{(2)}_{k_1^{(2)}})\rangle = \langle \Gamma^{(3)}_{h_2 k_2^{(3)} k_3^{(3)}} : k_3^{(3)} \in K(\gamma^{(3)}_{h_2 k_2^{(3)}})\rangle;\\
&\qquad \dots\\
&\exists h_n\ \gamma^{(n)} = \gamma^{(n+1)}_{h_n}\\
&\ \ \&\ \langle \gamma^{(n)}_{k_1^{(n)}} : k_1^{(n)} \in K(\gamma^{(n)})\rangle = \langle \gamma^{(n+1)}_{h_n k_2^{(n+1)}} : k_2^{(n+1)} \in K(\gamma^{(n+1)}_{h_n})\rangle\\
&\ \ \&\ \dots\\
&\ \ \&\ \langle \gamma^{(n)}_{k_1^{(n)}\dots k_{i+1}^{(n)}} : k_{i+1}^{(n)} \in K(\gamma^{(n)}_{k_1^{(n)}\dots k_i^{(n)}})\rangle = \langle \gamma^{(n+1)}_{h_n k_2^{(n+1)}\dots k_{i+1}^{(n+1)}} : k_{i+1}^{(n+1)} \in K(\gamma^{(n+1)}_{h_n k_2^{(n)}\dots k_i^{(n+1)}})\rangle;\\
&\ \ \&\ \dots\\
&\ \ \&\ \langle \Gamma^{(n)}_{k_1^{(n)}\dots k_n^{(n)}} : k_n^{(n)} \in K(\gamma^{(n)}_{k_1^{(n)}\dots k_{n-1}^{(n)}})\rangle = \langle \Gamma^{(n+1)}_{h_n k_2^{(n+1)}\dots k_n^{(n+1)}} : k_n^{(n+1)} \in K(\gamma^{(n+1)}_{h_n k_2^{(n)}\dots k_{n-1}^{(n+1)}})\rangle;\\
&\qquad \dots
\end{aligned} \tag{9.4b}$$

Iterating systems $\mathcal{G}(\omega^2\cdot n+\omega^2)$ through all n, we obtain a $\mathcal{G}(\omega^2\cdot\omega) = \mathcal{G}(\omega^3)$ system (comprising within it $\mathcal{G}(\omega^2\cdot n)$ systems as 'building-blocks'), and every $A \in \mathcal{U}(\omega^3)$ is represented by one such by the inductive reasoning of before, i.e., since each component $[x]^A_{\omega^2\cdot n}$ is represented by a constituent $\mathcal{G}(\omega^2\cdot n)$ system.

[9.3] Rank ω^ω

Now, we'd like to define a $\mathcal{G}(\alpha+\omega^3)$ as the iteration just described, starting with a $\mathcal{G}(\alpha)$ system $\Gamma^{(0)}_0$, Iterating yet again such systems $\mathcal{G}(\omega^3\cdot n+\omega^3)$ through all n, we should obtain a $\mathcal{G}(\omega^4)$ system. Iterating through all $\mathcal{G}(\omega^n)$ systems, we should obtain a $\mathcal{G}(\omega^\omega)$ system. *And so on.* But we are not ready to carry this out here, and will contend ourselves with merely having indicated how this 'iteration' might be continued through an initial segment of the ordinals.

(9.5) **Challenge** Devise a recursive system of notation for uniform chains of constructible rank.

References:

[P] Francis Pastijn, *A class of uniform chains*, in: S.M.Goberstein and P.M.Higgins [eds.] Semigroups and their applications.

Revision history:

Research for this paper was carried out between 1996 and 1999 by the first author. A first draft was completed December 2000. A substantially revised second draft, begun almost a year later, was completed January 2003. Some streamlining and improvements to the diagrammes resulted in the first submission to the 'arxiv' (December 2003). The ideas of the conclusion added for this final revision represent unfinished business. [Contact address: 94 Spencer Place, Leeds, LS7 4DT, UK]